

Table of Contents

List of figures	IV
List of tables	VI
List of abbreviations	VII
Summary	IX
Zusammenfassung	XI
1. Introduction	1
1.1 The grapevine <i>Vitis vinifera</i> ssp. <i>sativa</i> and viticulture	1
1.2 Grapevine diseases	2
1.2.1 Downy mildew	3
1.2.2 Life cycle of <i>P. viticola</i> causing downy mildew disease	4
1.3 The plant immune system	6
1.3.1 PTI: PAMP-Triggered Immunity	9
1.3.2 ETI: Effector Triggered Immunity	10
1.4 Current status of downy mildew resistance	12
1.4.1 Wild species: <i>Vitis coignetiae</i>	13
1.5 Grapevine resistance breeding	14
1.5.1 Artificial Intelligence-based phenotyping	14
1.5.1.1 The Convolutional Neural Networks	15
1.5.2 Genotyping	17
1.5.2.1 PIWI cultivars (ger. pilzwiderstandsfähige Rebsorten)	17
1.5.2.2 Marker-Assisted Selection	18
1.5.2.3 Pyramiding resistance loci	20
1.6 Research objectives	20
2 Materials and Methods	22
2.1 Materials	22
2.1.1 List of chemicals and kits	22
2.1.2 List of laboratory equipments	23
2.1.3 List of software and online tools	25
2.1.4 Plant material	26
2.1.5 <i>Plasmopara viticola</i> spore material	27

2.2	Methods	27
2.2.1	Phenotypic assessment	27
2.2.1.1	Leaf disc assay	27
2.2.1.2	High throughput imaging	28
2.2.1.3	Manual phenotyping: Downy Mildew	29
2.2.1.4	Manual Phenotyping: Leaf hair	29
2.2.1.5	Correlation of Phenotypic data: Downy Mildew vs. leaf hair	30
2.2.2	DNA extraction	30
2.2.3	Marker analysis	30
2.2.3.1	SSR markers.....	30
2.2.3.2	PCR.....	31
2.2.3.3	RNase H2-dependent amplicon sequencing markers	32
2.2.4	Genetic mapping	32
2.2.4.1	Simple Sequence Repeats based genetic map.....	32
2.2.4.2	RNase H2-dependent amplicon sequencing marker based genetic map	33
2.2.5	QTL analysis	34
2.2.5.1	Simple Sequence Repeats	34
2.2.5.2	RNase H2-dependent amplicon sequencing markers	34
2.2.6	Microscopy: Aniline blue staining	35
2.2.7	Growth and sporulation.....	35
2.2.8	Artificial Intelligence-based phenotyping.....	36
2.2.8.1	SCNN: Training and performance evaluation	36
2.2.9	Statistics	37
3	Results.....	38
3.1	SCNN-based downy mildew quantification	38
3.1.1	SCNN: Training results.....	38
3.1.2	Experts generated ground truth data.....	39
3.1.3	Overall performance evaluation.....	41
3.1.3.1	Classification pipeline: SCNN and ResNet	42
3.2	Summary of phenotypic data	44

3.2.1	Downy mildew resistance and morphological trait leaf hair (trichome) density	44
3.3	Genetic map	46
3.3.1	Initial map: SSR marker	46
3.3.2	Fine mapping: rhAmpSeq marker	51
3.4	QTL analysis	56
3.4.1	SSR marker data	56
3.4.2	rhAmpSeq marker data	58
3.4.2.1	QTL analysis: downy mildew resistance	58
3.4.2.2	QTL analysis: Leaf hair	59
3.4.3	Marker analysis to differentiate <i>Rpv8</i> , <i>Rpv12</i> and <i>Rpv32</i> locus	63
3.4.4	Correlation between downy mildew resistance and leaf hair density	64
3.5	Aniline blue staining	64
3.6	Comparison of different resistances: <i>Rpv8</i> , <i>Rpv12</i> , and <i>Rpv32</i>	65
4	Discussion	68
4.1	New resistance locus in ‘Morio Muskat’ x COxGT2 population	68
4.1.1	SCNN: An artificial intelligence based downy mildew quantification system	68
4.1.2	Phenotypic evaluation of downy mildew	71
4.1.3	Mapping and QTL analysis	73
4.1.4	Aniline blue staining	76
4.1.5	Growth and sporulation of <i>P. viticola</i> on different resistance loci carriers	78
4.1.6	Downy mildew vs. leaf hair: A weak negative correlation	79
5	Conclusion and future perspective	82
	References	84
	Annex I	113
	Annex II	114
	Annex III	115
	ResNet model development: Training and performance evaluation	115
	Annex IV	118
	Acknowledgments	133