

1


The Palladium/Norbornene-Catalyzed Annulation Chemistry: Rapid Access to Diverse Ring Structures

Mark Lautens and Xavier Abel-Snape

Davenport Research Laboratories, Department of Chemistry, University of Toronto, Toronto, Ontario M5S 3H6, Canada

1.1 Introduction

Palladium–norbornene (NBE) cooperative catalysis, commonly known as the Catellani reaction, constitutes a general and straightforward method to sequentially difunctionalize a haloarene at the *ortho* and *ipso* positions, with two different reagents [1–6]. These reagents are typically opposite in reactivity, as the first to react does so as an electrophile (**E**), which will functionalize the *ortho* site, while the second serves as a terminating reagent, which is often a nucleophile in character (**Nu**), which will add at the *ipso* position (Scheme 1.1). This sequence is made possible due to a unique combination of characteristics, including NBE's exceptional reactivity due to the strain, the resulting rigid framework that creates a transient directing group, and lack of accessible β -hydrogens, that prevent side reactions.




X: I, Br, OTf

Scheme 1.1 The general Catellani reaction.

The mechanism has been investigated in detail. Following oxidative addition into the C—X bond, the initial arylpalladium(II) species preferentially reacts with NBE via carbopalladation in order to release its ring strain rather than with the terminating reagent (Scheme 1.2). Every Catellani reaction subsequently generates a key intermediate, known as the arylnorbornyl palladacycle (**ANP**), which is typically formed after concerted metalation deprotonation (CMD) occurs at the *ortho* position in the presence of a base. The electrophile is then installed via one of two possible pathways: (i) oxidative addition to form a Pd(IV) intermediate followed by reductive elimination or (ii) dinuclear transmetalation [7–9]. Due to the steric congestion,

NBE is then extruded *via* β -C elimination, giving rise to a new *ortho*-functionalized arylpalladium(II) species that can now react with the terminating reagent.



Legend

O.A.: Oxidative addition

M.I.: Migratory insertion

C.M.D.: Concerted metalation-deprotonation

D.T.M.: Dinuclear transmetalation

R.E.: Reductive elimination

β -C Elim.: Beta-carbon elimination

TM.: Transmetalation

L.S.: Ligand substitution

Scheme 1.2 The general mechanism of the Catellani reaction.

In most cases, the aryl group bears one *ortho*-substituent to avoid di-*ortho*-functionalization with the electrophile or NBE-integrated side-products and ultimately, a lower yield of desired product [9]. This requirement is known as the *ortho* constraint. To tackle this issue, various modified NBE scaffolds have been developed to successfully employ *ortho*-unsubstituted aryl halides as substrates that give good to excellent yields [10–12].

Chapter 1 presents various cyclization methodologies harnessing Pd-NBE cooperative catalysis. The first section describes the most common way of forming rings, i.e. intramolecular cyclization, where two or all three out of the aryl halide, electrophile or terminating reagent are tethered to one another. The second section reports annulations involving sequential intermolecular *ortho*-functionalization


and ring closure steps with external reagents. Three-membered rings constitute the focus of section three, where their innate strain turns them into valuable electrophiles and terminating reagents upon ring opening, thereby forming five-membered rings. Reactions where NBE and its analog norbornadiene find themselves incorporated in the final annulated product instead of solely being used as transient directing groups are included in section four. The final section is comprised of reactions where the annulation step occurs after the catalytic cycle.

1.2 Intramolecular Cyclizations

1.2.1 Electrophile Tethered to Terminating Reagent

1.2.1.1 *Ortho* Alkylation


***Ips*o Heck Termination** The original 1997 report by Catellani described a reaction between an unsubstituted or *para* substituted aryl iodide, an alkyl halide, and a Heck acceptor [1]. The catalyst, known as the PNP complex, was a phenyl norbornyl palladium halide dimer prepared from phenyl mercuric chloride, NBE, and palladium chloride. In 1999, Pd(OAc)₂ in DMF was shown to be a suitable combination for reacting *ortho*-substituted aryl iodides [13]. In 2000, Lautens developed an annulative process and reported what have become the most widely used conditions, namely Pd(OAc)₂, phosphines, acetonitrile, and cesium carbonate (Scheme 1.3). In this example, the electrophile, i.e. the alkyl bromide, is tethered to the Heck acceptor providing access to fused ring systems [14].



Scheme 1.3 First annulative Catellani methodology.


This set of conditions paved the way for subsequent ring-forming processes, generating a variety of benzofused carbo- and heterocycles via *ortho*-alkylation and *ipso*-Heck termination under identical or modified conditions (Scheme 1.4) [15–19]. Some of these examples illustrate that heterocycles are tolerated, which was not possible until Lautens' report in 2006 [18].

Alkyl bromides were generally preferred over the analogous iodides likely due to potential side reactions, namely oxidative addition of Pd(0) into the C(sp³)-I bond followed by β-H elimination and reductive elimination to give the corresponding olefin and HI [13]. However, the iodides were ideal electrophiles in a β-fluoroalkylation process [20]. Alkyl tosylates were also found to be compatible electrophiles in the Catellani reaction [21].




Scheme 1.4 Examples of *ortho*-alkylation/*ipso*-Heck-termination annulative methodologies.

The Zhou group identified epoxides as alkylating reagents in a macrocyclization event using the potassium salt of 5-NBE-2-carboxylic acid **N**¹ (Scheme 1.5) [22].



Scheme 1.5 Macrocyclization using an epoxide as an alkylating reagent.


(Homo)allylic alcohols were suitable as the Heck acceptor, furnishing the corresponding carbonyl compounds via a redox-relay Heck cyclization (Scheme 1.6) [23].



Scheme 1.6 First methodology using (homo)allylic alcohols as the Heck acceptor.


Zhou was able to generate ring sizes ranging from five to seven [19, 24, 25]. Dong was also able to provide aldehyde-tethered rings using modified procedures (Scheme 1.7) [26, 27].

***Ips*o C–H Arylation** The first examples of annulative C–H arylation were reported by Lautens in 2005. The use of an unfunctionalized arene offers an attractive alternative to cross-coupling reactions where both arenes typically need a compatible



Scheme 1.7 Examples of *ortho*-alkylation/*ipso*-redox-relay Heck annulative methodologies.

functional group. Lautens showcased the power of C–H arylation by generating annulated indoles (Scheme 1.8) [28].




Scheme 1.8 Synthesis of annulated indoles via *ipso* C–H arylation.

This concept was generalized to include the synthesis of related hetero- and carbocycles (Scheme 1.9) [29–36].


***Ips*o Alkyne Insertion** Following *ortho*-alkylation and NBE extrusion, the resulting arylpalladium(II) species may undergo a migratory insertion relay step, followed by subsequent annulation reactions that increase molecular complexity.

Lautens reported reactions of alkyne-substituted alkyl halides that lead to *ipso*-alkyne insertion and C–H functionalization, leading to tetracyclic-fused pyrrole and indole derivatives. Carbopalladation of the alkyne precedes the C–H activation (Scheme 1.10a,b) [37, 38]. A related approach was reported a few years later to furnish tetrasubstituted helical alkenes (Scheme 1.10c) [39].

The vinyl-Pd(II) species can undergo an *exo*-migratory insertion across NBE or norbornadiene followed by C–H activation to incorporate the bicycle in the final product. This method provided a different kind of tetrasubstituted helical alkenes as a single diastereomer (Scheme 1.11a) [40]. Interestingly, using chiral bromoalkyl aryl alkynes resulted in moderate diastereoselectivities (Scheme 1.11b) [41]. It was proposed the R^4 substituent induces helical chirality upon *ipso*-alkyne insertion and the resulting major vinylpalladium(II) species is favored over the minor due




Scheme 1.9 Examples of *ortho*-alkylation/*ipso*-C–H arylation annulative methodologies.




Scheme 1.10 *Ipso*-alkyne insertion followed by C–H activation to form (a) tetracyclic-fused pyrrole derivatives (b) tetracyclic-fused indole derivatives (c) tetrasubstituted helical alkenes.

to 1,3-allylic strain between the pseudoequatorial R^4 substituent and R^3 -aryl ring [42, 43]. In both cases, NBE undergoes *exo*-insertion into the C–Pd(II) bond with its methylene group facing away from R^1 . Subsequently, C–H activation onto the alkyne-tethered arene occurs, followed by reductive elimination.




Scheme 1.11 Syntheses of tetrasubstituted helical alkenes (a) initial report (b) subsequent work using enantiomerically pure bromoalkyl aryl alkynes.

Nucleophilic attack on the vinyl-Pd(II) species can also occur. The Luan group reported two methodologies involving the dearomatization of indole and a phenol system, respectively, thereby forming a spiro palladacycle upon nucleophilic substitution (Scheme 1.12) [44, 45]. In a related report, Zhang, Liang, Li, and Quan synthesized indoles via a concerted C–N bond forming and N–S bond cleaving process following *ipso*-alkyne insertion [46].



Scheme 1.12 *Ipso*-alkyne insertion followed by dearomatization of (a) indoles (b) phenols.

Ipsa Dearomatization Using a similar bromoalkyl-tethered indoles with a free N–H group, a dearomatization step can occur in the presence of a base. The *ortho*-functionalized arylpalladium(II) intermediate undergoes a ligand substitution with the deprotonated indole at its 3-position to subsequently provide spiroindolenines (Scheme 1.13) [47].



Scheme 1.13 Synthesis of spiroindolenines.


Ipsa Enolate Termination A related carbon-based nucleophile can be generated as a metal-enolate. Zhou developed a three-component synthesis of 1,8-disubstituted tetralines from 2-substituted aryl iodides, aryl methyl ketones and 1-bromo-3-chloropropane (Scheme 1.14a). It was proposed the aryl methyl ketones reacted with 1-bromo-3-chloropropane under the basic conditions via an S_N2 reaction to form a bromoalkyl-tethered ketone prior to entering the catalytic cycle [48]. Liang synthesized spirodihydroindenones using a bromoalkyl-tethered cyclopentanone bearing an acidic α -proton (Scheme 1.14b) [49]. Zhou reported an enantioselective annulative process using a bromomethyl-tethered cyclohexanone that formed an enamine *in situ* with a chiral amino acid catalyst (Scheme 1.14c) [50].

Ipsa C–Alkyl Termination Alkyl nucleophiles using organometallic reagents are generally considered to be less successful in transition-metal-catalyzed reactions compared to aryl or vinyl nucleophiles due to the increased number of possible side-reactions that may occur, for instance β -H elimination and the more difficult transmetallation processes. As such, using an alkyl carbagermatrane as a fine-tuned organogermanium reagent, Xiao was able to construct carbocycles with ring sizes ranging from six to eight (Scheme 1.15) [51].


Ipsa C–N Termination A nucleophilic heteroatom can also be employed as a compatible *ipso* terminating reagent. Using brominated alkylamines, Lautens was able to furnish indolines and tetrahydroquinolines depending on the alkyl chain's length (Scheme 1.16) [52, 53]. It was established that a *para*-nitrophenyl group as R^4 was the optimal nitrogen-protecting group for the synthesis of indolines. Phenyl and ethoxy-carbonyl were the only other groups that were found to be compatible.

1.2.1.2 *Ortho* Arylation



Ipsa C–N Termination *Ortho*-arylation is usually conducted with a less reactive haloarene than the one meant to undergo sequential *ortho*- and *ipso*-functionalizations. Typically, the former is an aryl bromide and the latter, an aryl iodide. The



(a) ● Seven examples ● Up to 58% yield ● 0.2 mmol scale




(b) ● 15 examples ● Up to 74% yield ● Up to 4.0 mmol scale



(c) ● 10 examples ● Up to 95% yield ● Up to 97% ee

Scheme 1.14 Harnessing *in situ* generated enolates to synthesize (a) tetralines (b) spirodihydroindenones (c) bridged ketones.




● 18 examples ● Up to 89% yield ● 0.1 mmol scale

Scheme 1.15 Synthesis of carbocycles using alkyl carbagermatranes.


careful choice of different haloarenes ensures Pd(0) oxidatively adds into the more reactive C–I bond preferentially and that the **ANP** then reacts with the less reactive aryl bromide. Lautens showed aryl triflates can be used instead of aryl iodides [54], while aryl chlorides can also constitute the *ortho*-arylation reagent [54–56]. Two bromoarenes can also be used as coupling partners, although significant electronic differences make one more reactive than the other [57].

Catellani applied this reasoning in her synthesis of 6-phenanthridinones by reacting iodoarenes with 2-bromobenzamides (Scheme 1.17) [58]. Once *ortho*-arylation



Scheme 1.16 Synthesis of (a) indolines and (b) tetrahydroquinolines via *ortho*-alkylation and *ipso*-Buchwald–Hartwig coupling.

and NBE extrusion occurred, an *ipso*-C–N coupling took place to furnish the desired azacycles.




Scheme 1.17 Synthesis of 6-phenanthridinones.

Various N-heterocycles of different ring sizes were synthesized based on this method (Scheme 1.18) [54, 55, 57, 59–66].


***Ips*o C–O Termination** Similarly, C–O bond formation can occur using the appropriate terminating reagents (Scheme 1.19 [67–70]).

Various C=X Bonds as *Ips*o Heck Terminating Reagents Carbonyls can serve as *ipso*-terminating reagents, although their reactivity differs significantly depending on the functional group to which they belong and on the reaction conditions. Lautens was able to synthesize 9*H*-fluoren-9-ols from ketones as well as 9*H*-fluoren-9-ones from esters and aldehydes via direct addition to the carbonyl (Scheme 1.20) [56].

Using a chiral NBE derivative, Zhou was able to generate fluorenols enantioselectively (Scheme 1.21) [71].



Scheme 1.18 Subsequent examples of *ortho*-arylation/*ipso*-C–N termination annulative methodologies.



Scheme 1.19 Examples of *ortho*-arylation/*ipso*-C–O termination annulative methodologies.


2-Bromoarylaldehyde hydrazones were used in a denitrogenative synthesis of fluorenes (Scheme 1.22) [72]. It was determined the reaction pathway does not proceed via carbene insertion.

***Ips*o Enolate Termination** The Lautens group included three examples of enolates as terminating reagents in their work on carbonyls as *ipso* terminating reagents (Scheme 1.23) [56]. Tweaking the conditions by removing water and changing the solvent from DME to acetonitrile modulated the system's reactivity and favored enolate formation rather than the direct addition of the arylpalladium(II) intermediate to the ketone.


1.2.1.3 *Ortho* Acylation

***Ips*o Heck Termination** Using a mixed anhydride, Dong discovered how to *ortho*-acylate aryl iodides [73]. Subsequently, an extension of this method was developed to generate macrocycles via an *ipso*-Heck termination step (Scheme 1.24) [74].


Smaller ring systems were also accessible using an analogous reagent (Scheme 1.25) [11]. A mixed anhydride could also be generated *in situ* from




Scheme 1.20 *Ortho*-arylation/*ipso*-C=X termination methodologies for the synthesis of (a) 9*H*-fluoren-9-ols from ketones (b) 9*H*-fluoren-9-ones from esters (c) 9*H*-fluoren-9-ones from aldehydes.




Scheme 1.21 Enantioselective synthesis of fluorenols.



Scheme 1.22 Synthesis of fluorenes via denitrogenation.




Scheme 1.23 Synthesis of phenanthren-9-ols.




Scheme 1.24 Macrocycle formation via *ortho*-acylation and *ipso*-Heck termination.

the corresponding carboxylic acid and the Yamaguchi reagent [75]. Carbamoyl chlorides were developed as alternative *ortho*-acylating reagents, giving access to related carbocycles [76, 77].




Scheme 1.25 Subsequent examples of *ortho*-acylation/*ipso*-Heck termination annulative methodologies.

***Ips*o C–H Arylation** Jiao was the first to use carbamoyl chlorides as *ortho*-acylating reagents in Pd/NBE chemistry. These reagents were tethered to aryl rings, thereby leading to an intramolecular *ipso* C–H arylation termination, which furnished the corresponding phenanthridinones (Schemes 1.26, 1.27) [77].



Scheme 1.26 First use of carbamoyl chlorides in Pd/NBE chemistry.




Scheme 1.27 Subsequent examples of *ortho*-acylation/*ipso*-C–H arylation annulative methodologies.

Dong obtained a fluorenone as a side-product in one of the scope examples of their reaction using benzoic anhydride and an isopropyl carbonate anhydride derived from benzoic acid (Scheme 1.27) [73]. Lumb and Luan obtained a similar side-product during the optimization of their reaction using benzoic anhydride as well [78]. Dong reported a single example starting from an alkenyl triflate [76]. Chen and Zhu demonstrated fluorinated imidoyl chlorides could also be used as *ortho*-acylating reagents [79].

1.2.1.4 *Ortho* Amination and *Ips*o Heck Termination

A significant advance in Catellani methodology was the *ortho*-amination using Pd/ NBE cooperative catalysis, as reported by Dong in 2013, using *O*-benzoylhydroxylamines. This finding was the first time a heteroatom was introduced at the *ortho* position [80]. Since then, multiple reports have made use of this methodology to generate aminated arenes as well as N-heterocycles. For example, Zhou developed an amination reagent tethered to a silyl enol ether to make N-containing bridged scaffolds (Scheme 1.28) [81].




Scheme 1.28 Synthesis of N-containing bridged scaffolds via *ortho*-amination/*ipso*-Heck annulation.

The Liu and Dong groups concurrently published on a related topic in which C3,C4-disubstituted indoles were synthesized using near-identical conditions (Scheme 1.29) [82, 83].


Shortly after, Dong discovered that a C7-brominated NBE (**N**⁷) was key in generating tetrahydrobenzo[*b*]azepines using *ortho*-unfunctionalized aryl iodides as substrates (Scheme 1.30a) [12].

This methodology bypasses the long-standing “*ortho* constraint.” All but two products were accompanied by a minor regioisomer arising from reinsertion




● 15 examples (Liu) & 20 examples (Dong) ● Up to 83% yield ● Up to 1 mmol scale

Scheme 1.29 Synthesis of C3,C4-disubstituted indoles via *ortho*-amination/*ipso*-Heck termination.



(a) ● 28 examples ● Up to 78% yield ● 0.1 mmol scale



(b) ● 20 examples ● Up to 95% yield ● 0.1 mmol scale


Scheme 1.30 Synthesis of tetrahydrobenzo[b]azepines using (a) C7- and (b) C2-substituted NBEs.

of the Pd–H species wherein the alkene ends up inside the seven-membered ring in conjugation with the aryl ring. It was postulated that the absence of an *ortho*-substituent leads to a Pd–H reinsertion step, thereby furnishing a tertiary C(sp³)–Pd(II) intermediate that undergoes β–H elimination inside the ring. When using *ortho*-functionalized aryl iodides, no isomerized products were obtained, most likely due to the *ortho*-substituents' bulkiness (Scheme 1.30b). Optimization of this reaction demonstrated C2-substituted NBE **N**⁸ constituted the most appropriate bicyclic alkene to direct the transformation.

A crystal structure of the complex shown in Scheme 1.31 was obtained, wherein the bromine atom acts as an L ligand coordinating to d⁸ Pd, thereby reducing side reactions by stabilizing reaction intermediates. The bromo substituent is proposed to stabilize the β–C elimination transition state to promote NBE extrusion rather than a second *ortho*-C–H activation step, which would lead to *ortho*-difunctionalization. Additionally, the bromine's electronegativity hinders direct C–C reductive elimination from the **ANP** due to charge repulsion with the metalated aryl ring. Lastly, although the Br substituent is more sterically hindered than a hydrogen, it is not bulky enough to prevent the **ANP** from reacting with the electrophile.


1.2.1.5 *Ortho* Alkenylation and *Ipsso* Amination via Reversal of Regioselectivity

Interestingly, using oxime acetates derived from methyl ketones led to formal *ortho*-alkenylation and *ipso*-amination (Scheme 1.32) [84]. Following the **ANP**'s



Scheme 1.31 Br on a modified NBE's C7 acting as an L ligand coordinating to d^8 Pd.


oxidative addition into the oxime acetate's N–O bond, the resulting Pd–N-bound isomer tautomerizes to its corresponding Pd–C-bound isomer. This process subsequently leads to a C–C reductive elimination at the *ortho*-position and a C–N reductive elimination at the *ipso*-position.



Scheme 1.32 Indole formation via reversal of traditional regioselectivity of the electrophilic and terminating groups.

1.2.1.6 Pd(II)-Initiated Annulations

Aryl (pseudo)halides and palladium(0) complexes are not the only entry points for Catellani reactions. Alternatively, boronic esters and free (N–H) indoles can react with Pd(II) complexes to initiate the desired transformation, as first demonstrated by Bach for intermolecular reactions (Scheme 1.33) [85–88]. In the case of boronic esters, reductive elimination generates Pd(0) after the formation of the ring via a Heck reaction step. Therefore oxygen must be present in the reaction mixture to oxidize Pd(0) to Pd(II) to complete the catalytic cycle. As for free (N–H) indoles, protodepalladation occurs after formation of the ring to regenerate the active Pd(II) catalyst, thereby avoiding the zero oxidation state.




Scheme 1.33 Pd(II)-initiated annulative methodologies using (a) boronic acids and (b) free (N–H) indoles as entry points.

1.2.2 Aryl Halide Tethered to Terminating Reagent

1.2.2.1 *Ortho* Alkylation, *Ortho* Arylation and *Ortho* Amination

A second type of intramolecular cyclization involves functionalizing the haloarene *ortho*-C–H bond followed by *ipso*-annulation with an already existing functional group tethered to the haloarene. The biggest challenge to overcome is the direct *ipso*-annulation, which would occur if the *ortho*-functionalization via an intermolecular formation of the **ANP** were too slow relative to the intramolecular cyclization step.

Lautens was the first to successfully report this annulative strategy, which provided *ortho*-alkylated benzoxacycles (Scheme 1.34) [89]. Only one other example involving *ortho*-alkylation followed by an intramolecular Heck reaction has been reported [90].




Scheme 1.34 First methodology of *ortho*-functionalization with an external reagent followed by intramolecular *ipso*-termination.

Subsequent methodologies based on this type of cyclization have generated various carbo- and heterocycles (Scheme 1.35). Other methodologies employing *ortho*-alkylations have been paired with *ipso* C–alkyl [91] and C–N terminations [92]. *Ortho*-arylations have been combined with *ipso*-functionalizations such as Heck [93, 94]), C–N [92], C–H arylations [95, 96], and alkyne insertion [97]. As for *ortho*-aminations, these have been matched with *ipso*-Heck [90]), C–N [98], C–H arylation [99–103], and C–H alkylation terminations [102, 104].


1.2.3 Aryl Halide Tethered to Electrophile

1.2.3.1 *Ortho* Alkylation

The third type of intramolecular cyclization involves functionalizing the haloarene substrate's *ortho*-C–H bond with an already existing functional group tethered to the haloarene followed by intermolecular *ipso*-annulation. In a seminal report, Lautens used *meta*-substituted iodoarenes to functionalize both *ortho*-positions with two different electrophiles, one being the tethered alkyl halide and the other, an external alkyl halide (Scheme 1.36a) [105]. A Heck termination constituted the *ipso*-functionalizing step. Aryl iodides with substituents at the 2- and 5-positions relative to the C–I bond were also compatible. However no external alkyl halide was added due to there being only one available *ortho*-C–H bond (Scheme 1.36b).




Scheme 1.35 Subsequent examples of *ortho*-functionalization with an external reagent followed by intramolecular *ipso*-termination.



Scheme 1.36 First methodology of *ortho*-functionalization with a tethered electrophile followed by intermolecular *ipso*-termination. (a) Unsymmetrical di-*ortho*-alkylation (b) Mono *ortho*-alkylation.


This method was subsequently combined with the original intramolecular cyclization strategy using a difunctional ambident species (bromoenoate or bromoalkyl indole) (Scheme 1.37) [106].

Since then, various groups have devised variations on the method, using Heck couplings [53, 107–111] as well as different terminations, including C(sp²)-H



Scheme 1.37 Combining an iodoarene tethered to an electrophile with an electrophile tethered to a terminating group. (a) bromoenoate (b) bromoalkylindole.

activation [112]), cyanation [111, 113, 114], carbene insertion [115], Suzuki [111], and Sonogashira coupling [111], Miyaura borylation [111], and hydride reduction (Scheme 1.38) [111].




Scheme 1.38 Subsequent examples of intramolecular *ortho*-functionalization followed by intermolecular *ipso*-termination.

1.2.4 Aryl Halide Tethered to Electrophile and Terminating Reagent

1.2.4.1 *Ortho* Alkylation


The final type of intramolecular cyclization consists of making the electrophilic and terminating groups part of the same molecule as the substrate haloarene. Only two reports of such a strategy have been disclosed, both by Lautens (Scheme 1.39) [53, 108].

The use of enantioenriched substrates led to the corresponding enantioenriched products with an overall inversion of configuration at the stereocenter with either minimal or significant decrease in ee depending on the substrate class and reaction conditions. In Scheme 1.40a, a mere 2% decrease in ee was observed when converting the alkyl iodide substrate to the corresponding tricyclic product. On the other hand, in Scheme 1.40b, subjecting the alkyl bromide substrate to the reaction conditions resulted in a cyclized product with a net –14% ee difference. Mechanistic



Scheme 1.39 Intramolecular *ortho*-alkylation followed by intramolecular (a) *ipso*-Heck termination (b) *ipso* C–H arylation.

studies showed CsI formed under the reaction conditions can racemize the substrate's chiral center in MeCN (Scheme 1.40b), but not in DME (Scheme 1.40a) due to the salt's increased solubility in the former solvent. It was proposed that inversion of configuration takes place at the oxidative addition step from Pd(II) to Pd(IV) via an $\text{S}_{\text{N}}2$ mechanism, as reductive elimination from Pd(IV) to Pd(II) occurs via retention of configuration [116–118].



Scheme 1.40 Study on the erosion of enantiomeric excess using enantioenriched substrates in (a) DME and (b) MeCN as solvents.


1.3 Intermolecular Cyclizations Following Intermolecular *Ortho* Functionalization

Contrary to Section 1, which focused on annulations wherein at least two out of the aryl halide, electrophile and terminating reagent components are tethered to one another, Section 2 describes ring-closing methods in which all three constitute separate molecules that react sequentially.

1.3.1 *Ortho* Arylation

1.3.1.1 *Ipso* Alkyne Insertion and *Ipso* Formal Benzyne Insertion


Catellani synthesized phenanthrenes using two equivalents of an *ortho*-substituted aryl iodide, one of them acting as the electrophile, and an alkyne partner to close the ring via sequential carbopalladation and reductive elimination (Scheme 1.41) [119]. Kwong also developed a regioselective synthesis of phenanthrenes, this time from two different aryl halides that allowed greater molecular complexity in the final product. CO₂ was extruded as a result of decarboxylation of the *ortho*-bromoarylcarboxylic acid in the catalytic cycle [120].



Scheme 1.41 Synthesis of phenanthrenes from an alkyne and (a) two identical aryl iodides (b) two different aryl halides.

Triphenylenes were synthesized by Liang and Yang according to a similar method to Catellani's wherein two equivalents of an *ortho*-bromoarylcarboxylic acid were needed (Scheme 1.42) [121].

Luan synthesized spirocycles via a three-component reaction using bromonaphthols as electrophiles, which ultimately underwent a dearomatization process in order to close the catalytic cycle (Scheme 1.43) [97].




● 21 examples

● Up to 99% yield

● Up to 1 mmol scale


Scheme 1.42 Synthesis of triphenylenes from an aryl iodide and two *ortho*-bromoarylcarboxylic acids.



● 43 examples

● Up to 89% yield

● Up to 4 mmol scale



Scheme 1.43 Synthesis of spirocarbocycles from aryl iodides, bromonaphthols, and alkynes.


1.3.2 *Ortho* Amination

1.3.2.1 *Ips*o Alkyne Insertion

Zhang et al. synthesized indoles via an interesting process (Scheme 1.44). The final steps of the reaction first involve the formation of an intermediary indole quaternary ammonium salt, which is postulated to weaken the N–Me bond. A final $\text{S}_{\text{N}}2$ displacement with a benzoate anion gives rise to the desired indole and methyl benzoate as the by-product [122].

1.4 Cyclizations with Three-Membered Heterocycles as Both the Electrophile and Terminating Reagent


Three-membered heterocycles can serve as both the electrophile and the terminating agent. Their innate strain renders them reactive to the oxidative addition step with the ANP, thereby breaking one of the C–heteroatom bonds. A chemoselective C–C reductive elimination occurs to functionalize the haloarene's *ortho*-position. Following NBE extrusion, C–N or C–O reductive elimination takes place to furnish a new heterocyclic five-membered ring.



Scheme 1.44 Synthesis of indoles from aryl iodides, *O*-benzoylhydroxylamines, and alkynes.


1.4.1 2*H*-Azirines

Lautens reported the synthesis of indoles using 2*H*-azirines, which were added via slow addition to the reaction mixture and resulted in a 1:1 ratio relative to the aryl iodide (Scheme 1.45a) [123]. Excess 2*H*-azirine resulted in the formation of dihydroimidazoles (Scheme 1.45b).



Scheme 1.45 Synthesis of (a) indoles and (b) dihydroimidazoles from aryl iodides and 2*H*-azirines.

Coordination of the nitrogen lone pair to the **ANP**'s Pd(II) center weakens the 2*H*-azirine's bonds, which allows oxidative addition of the Pd(II) into the ligand's C–N single bond to occur to furnish the corresponding Pd(IV) complex (Scheme 1.46) [124–129]. Reductive elimination via a chemoselective C–C bond formation leads to an eight-membered palladacycle, which is followed by NBE extrusion to give rise to a more stable six-membered palladacycle. A 3*H*-indole is then formed by a final reductive elimination, thereby regenerating the active Pd(0) catalyst. Tautomerization to the more thermodynamically stable 1*H*-indole takes place when no excess 2*H*-azirine is present. However, exposure to another




Scheme 1.46 Proposed mechanism for the formation of indoles and dihydroimidazoles from the ANP and 2H-azirine(s).

equivalent of 2H-azirine leads to a dihydroimidazole via a postulated Pd(0)-catalyzed formal [3 + 2]-cycloaddition.

1.4.2 Aziridines


Similarly, Bi and Liang generated indolines from aryl iodides and aziridines (Scheme 1.47) [130]. These make analogous products to those previously reported by Lautens using bromoethylamines (Scheme 1.17) [52]. The latter initially thought the formation of aziridines from bromoethylamines by intramolecular nucleophilic displacement had to be avoided for the reaction to work. However, Bi and Liang's report suggests that the formation of the aziridine might have been the productive reaction pathway.



Scheme 1.47 Synthesis of indolines from aryl iodides and (a) 2-unsubstituted aziridines (b) 2-substituted aziridines.

1.4.3 Epoxides

Dong synthesized 2,3-dihydrobenzofurans from aryl iodides and epoxides (Scheme 1.48a) [131]. The same family of products was later generated by Cheng and Zhou under considerably different conditions (Scheme 1.48b) [132].



Scheme 1.48 Synthesis of 2,3-dihydrobenzofurans from aryl iodides and epoxides (a) conditions developed by Dong (b) conditions developed by Cheng and Zhou.


Both methodologies include an example of an enantiopure epoxide that leads to a stereoretentive annulation without any loss of ee in the product. Subsequently, Dong reported an enantioselective version of the reaction using racemic epoxides with an enantiopure chiral NBE derivative that furnished the corresponding 2,3-dihydrobenzofurans in moderate ee [133].

1.5 Norbornene/Norbornadiene-Integrated Cyclizations

1.5.1 Norbornene


Although NBE is usually used as a transient directing group, it can sometimes be integrated in the final product to form different types of ring systems. Some reports have even disclosed results showing that exceptions to the general rules in Catellani chemistry exist.

For example, NBE-incorporated adducts derived from $C(sp^2)$ – $C(sp^3)$ reductive elimination from the Pd(IV) intermediate were generated instead of products derived from the usual $C(sp^2)$ – $C(sp^2)$ reductive elimination (Scheme 1.49a) [134]. This outcome could be explained due to chelation of the amide moiety of the 2-bromophenylacetamide that generates steric strain in the transition state that would lead to $C(sp^2)$ – $C(sp^2)$ reductive elimination, thereby hindering this pathway. However, it was observed that adding water leads to $C(sp^2)$ – $C(sp^2)$ reductive elimination due to displacement of the $P(2\text{-furyl})_3$ ligand, thereby decreasing the steric strain in the corresponding transition state and favoring this pathway. Water also hampered NBE extrusion and rather led to a dearomatized product via sequential 5-*exo*-migratory insertion and β -H elimination (Scheme 1.49b).



Scheme 1.49 Synthesis of NBE-containing adducts due to a coordinating amide-moiety (a) general reaction conditions (b) study of the effect of added water on the selectivity of the reaction.


Similarly, dihydrodibenzoazepine derivatives were synthesized from aryl iodides, *ortho*-bromoanilines, and NBE (Scheme 1.50a) [135]. These compounds resulted from $C(sp^2)-C(sp^3)$ reductive elimination from the Pd(IV) intermediate due to the chelation of the *ortho*-bromoaniline's amino group to palladium. A subsequent report used aryl bromides instead of aryl iodides with potassium iodide as an additive (Scheme 1.50b) [136].



Scheme 1.50 Synthesis of NBE-containing azacycles following $C(sp^2)-C(sp^3)$ reductive elimination using (a) aryl iodides (b) aryl bromides.


Interestingly, while combining *ortho*-substituted aryl iodides with 2-bromo-NH-sulfoximines leads to the usual $C(sp^2)-C(sp^2)$ reductive elimination, aryl iodides containing electron-withdrawing groups at the *meta*-position led to NBE-containing adducts resulting from $C(sp^2)-C(sp^3)$ reductive elimination (Scheme 1.51a) [62]. Increasing the equivalents of 2-bromo-NH-sulfoximine led to adducts bearing two sulfoximine moieties (Scheme 1.51b).

Other examples involve various termination steps that outcompete NBE extrusion [137–147]. For example, these steps may involve nucleophilic substitution on the



Scheme 1.51 Synthesis of NBE- and sulfoximine-containing adducts using (a) one equivalent of 2-bromo-NH-sulfoximine (b) two equivalents of 2-bromo-NH-sulfoximine.

norbornylpalladium(II) intermediate or a migratory insertion into the C–Pd bond of that intermediate (Scheme 1.52).



Scheme 1.52 Subsequent examples of NBE-containing adducts.

As mentioned earlier (Section *Ips*o Alkyne Insertion), Lautens was able to incorporate NBE in the final product following *ortho* and *ipso* functionalization to provide tetrasubstituted helical alkenes (Scheme 1.11) [40, 41].

1.5.2 Norbornadiene


Although rarely used, norbornadiene can replace NBE and be a compatible partner in Catellani-type reactions. Both of their alkenyl hydrogens deviate in the *endo* direction relative to the C1–C2 plane, with an angle of 7° for NBE and about 2–4° for norbornadiene (Scheme 1.53) [148]. In both cases, the fact the alkene moiety is pyramidalized instead of flat is a result of the system attempting to

minimize torsional strain and therefore the distortion energy. This leads to a distortion-accelerated reaction upon submitting the bicyclic alkene to a compatible reagent.




Scheme 1.53 Structural comparison between NBE and norbornadiene.

In 2017, Luan reported the synthesis of aminated spirocarbocycles having integrated a norbornadiene fragment (Scheme 1.54) [149]. Following *ortho*-amination, palladium-induced phenol dearomatization outcompetes norbornadiene extrusion via ligand substitution on the metal.




Scheme 1.54 Synthesis of norbornadiene-fused spirocarbocycles via phenol dearomatization.

Incorporation of a retro-Diels–Alder step formally installs a molecule of acetylene into the product, ejecting a molecule of cyclopentadiene. Derat and Catellani demonstrated this concept in their synthesis of dibenzoazepines by replacing NBE with norbornadiene (Scheme 1.55a) [135]. Della Ca' used an aryl bromide instead of an aryl iodide and obtained analogous results (Scheme 1.55b) [136].




Scheme 1.55 Synthesis of dibenzoazepines following C(sp²)–C(sp³) reductive elimination using (a) aryl iodides (b) aryl bromides.

Lautens had shown that *ortho*-bromoanilines and *ortho*-chlorobenzamides react with norbornadiene in the presence of palladium, and undergo a retro Diels–Alder step to provide access to indoles and isoquinolinones [150]. The Liang group adapted this strategy and added in a Catellani step to construct C4-functionalized indoles via *ortho*-amination or *ortho*-glycosylation, with the C2 and C3 atoms being provided by norbornadiene or its tosylated derivative (Scheme 1.56) [151, 152]. Using 2-iodobiphenyls as substrates, Liang was able to generate 1-aminated phenanthrenes (Scheme 1.57) [153]. A concurrent synthesis of such products was also developed by Yang and Liang using similar conditions [154].



Scheme 1.56 Synthesis of C4-functionalized indoles via (a) *ortho*-amination (b) *ortho*-glycosylation.



Scheme 1.57 Synthesis of aminated phenanthrenes.


Finally, Lin and Kwong employed 2-haloarylcarboxylic acids and 8-bromo-1-naphthoic acid to provide the phenanthrene and benzo[4,5]cyclohepta[1,2,3-*de*]naphthalene derivatives, respectively (Scheme 1.58) [155].

1.6 One-Pot Postcatalytic Intramolecular Cyclizations

1.6.1 Postcatalytic Intramolecular Michael Additions


1.6.1.1 *Ortho* Alkylation

In some cases, the product of the catalytic reaction does not yet contain a ring, but subsequent cyclization can occur after the catalytic cycle. For example, using



Scheme 1.58 Synthesis of (a) phenanthrene derivatives and (b) heptagon-embedded aromatic compounds via *ortho*-arylation.

N-Cbz-bromoalkylamines and α -chloroamides as the electrophile and activated alkenes as the *ipso*-Heck acceptor, Ferraccioli synthesized six- and seven-membered N-heterocycles via a postcatalytic aza-Michael addition (Scheme 1.59) [156–158].




Scheme 1.59 Synthesis of N-heterocycles via a postcatalytic aza-Michael addition using activated alkenes and (a) N-Cbz-bromoalkylamines (b) α -chloroamides.

Epoxides and aziridines have also been used as electrophiles by Cheng and Zhou that, respectively, formed alcohols and amines as products after the catalytic cycle, respectively. Similarly to Ferraccioli's work, oxa- and aza-Michael additions then took place to form the corresponding oxa- and azacycles (Scheme 1.60) [22, 159, 160].

1.6.1.2 *Ortho* Arylation

Bromoarenes have also been used in this type of process to synthesize 6*H*-dibenzopyrans, dihydrophenanthridines, and dibenzazepines (Scheme 1.61) [161–164]. 6*H*-Dibenzopyrans were subsequently generated enantioselectively using a cinchona alkaloid cocatalyst [165].




Scheme 1.61 Synthesis of dibenzocycles using (a) *ortho*-bromophenols, (b) *ortho*-bromoanilines and (c) *ortho*-bromobenzylamines.

Interestingly, using methyl vinyl ketone as the *ipso*-terminating reagent leads to a retro-Mannich reaction to furnish the corresponding phenanthridines and imines (Scheme 1.62) [164, 166].


1.6.2 Postcatalytic Intramolecular Addition to Norbornyl Moiety

Finally, an interesting NBE-containing scaffold could be obtained using 3-iodochromones, α -bromoacetophenones and NBE (Scheme 1.63) [167]. Instead of



Scheme 1.62 Postcatalytic retro-Mannich reactions using MVK as the *ipso*-terminating reagent to form (a) phenanthridines and (b) imines.

the usual β -C elimination step resulting in the extrusion of NBE, a base-mediated β -H elimination step took place, which kept the norbornyl fragment in the product. Keto-enol tautomerization followed by intramolecular cyclization was proposed to deliver the spirocycle in a formal (2 + 3 + 1) annulation. A subsequent report using tetracyclododecene instead of NBE led to a formal (2 + 2 + 1) annulation [168].



Scheme 1.63 Postcatalytic intramolecular Michael addition onto an NBE moiety.

1.7 Summary

Since the first report by Catellani in 1997, the use of NBE to temporarily direct C–H functionalization has become of increasing interest in the synthesis of functionalized aromatic compounds. Lautens first introduced phosphine ligands to the catalytic system, which are now broadly used in this type of reaction. Lautens also reported the first annulative approach using the Catellani reaction. In the intervening two decades, many research groups have contributed to the reaction and to annulation strategies specifically. Ring formation can occur in many ways, including by tethering the electrophile or nucleophile, intermolecularly following *ortho* functionalization with an external electrophile, by using a three-membered ring as the coupling partner, by integrating NBE or norbornadiene in the final scaffold, and following of the catalytic cycle. The future of this field looks brighter with each passing year as more groups provide creative approaches that incorporate annulative Catellani reactions.

References

- 1 Catellani, M., Frignani, F., and Rangoni, A. (1997). A complex catalytic cycle leading to a regioselective synthesis of o,o'-disubstituted vinylarenes. *Angew. Chem. Int. Ed. Engl.* 36 (1–2): 119–122.
- 2 Catellani, M., Motti, E., Faccini, F., and Ferraccioli, R. (2005). New catalytic methods for the synthesis of selectively substituted aromatics through palladacycles. *Pure Appl. Chem.* 77 (7): 1243–1248.
- 3 Martins, A., Mariampillai, B., and Lautens, M. (2010). Synthesis in the key of Catellani: norbornene-mediated ortho C–H functionalization. In: *C–H Activation* (ed. J.Q. Yu and Z. Shi), 1–33. Berlin, Heidelberg: Springer Berlin Heidelberg.
- 4 Ye, J. and Lautens, M. (2015). Palladium-catalysed norbornene-mediated C–H functionalization of arenes. *Nat. Chem.* 7 (11): 863–870.
- 5 Della Ca', N., Fontana, M., Motti, E., and Catellani, M. (2016). Pd/Norbornene: a winning combination for selective aromatic functionalization via C–H bond activation. *Acc. Chem. Res.* 49 (7): 1389–1400.
- 6 Wang, J. and Dong, G. (2019). Palladium/norbornene cooperative catalysis. *Chem. Rev.* 119 (12): 7478–7528.
- 7 Catellani, M. (2002). Arene substitution involving temporary incorporation and removal of carbon tethers via carbopalladation and decarbopalladation. In: *Handbook of Organopalladium Chemistry for Organic Synthesis* (ed. E.I. Negishi), 1479–1489. John Wiley & Sons, Inc.
- 8 Cárdenas, D.J., Martín-Matute, B., and Echavarren, A.M. (2006). Aryl transfer between Pd(II) centers or Pd(IV) intermediates in Pd-catalyzed domino reactions. *J. Am. Chem. Soc.* 128 (15): 5033–5040.
- 9 Maestri, G., Motti, E., Della Ca', N. et al. (2011). Of the ortho effect in palladium/norbornene-catalyzed reactions: a theoretical investigation. *J. Am. Chem. Soc.* 133 (22): 8574–8585.
- 10 Wang, J., Li, R., Dong, Z. et al. (2018). Complementary site-selectivity in arene functionalization enabled by overcoming the ortho constraint in palladium/norbornene catalysis. *Nat. Chem.* 10 (8): 866–872. <https://doi.org/10.1038/s41557-018-0074-z>.
- 11 Liu, F., Dong, Z., Wang, J., and Dong, G. (2019). Palladium/norbornene-catalyzed indenone synthesis from simple aryl iodides: concise syntheses of pauciflorol F and acedinone A. *Angew. Chem. Int. Ed.* 58 (7): 2144–2148.
- 12 Liu, X., Wang, J., and Dong, G. (2021). Modular entry to functionalized tetrahydrobenzo[b]azepines via the palladium/norbornene cooperative catalysis enabled by a C7-modified norbornene. *J. Am. Chem. Soc.* 143 (26): 9991–10004.
- 13 Catellani, M. and Cugini, F. (1999). A catalytic process based on sequential ortho-alkylation and vinylation of ortho-alkylaryl iodides via palladacycles. *Tetrahedron* 55 (21): 6595–6602.
- 14 Lautens, M. and Piguel, S. (2000). A new route to fused aromatic compounds by using a palladium-catalyzed alkylation-alkenylation sequence. *Angew. Chem. Int. Ed.* 39 (6): 1045–1046.

- 15 Lautens, M., Paquin, J.F., Piguel, S., and Dahlmann, M. (2001). Palladium-catalyzed sequential alkylation – alkenylation reactions and their application to the synthesis of fused aromatic rings. *J. Org. Chem.* 66 (24): 8127–8314.
- 16 Lautens, M., Paquin, J.F., and Piguel, S. (2002). Palladium-catalyzed sequential alkylation-alkenylation reactions. Application to the synthesis of 2-substituted-4-benzoxepines and 2,5-disubstituted-4-benzoxepines. *J. Org. Chem.* 67 (11): 3972–3974.
- 17 Alberico, D., Paquin, J.F., and Lautens, M. (2005). Palladium-catalyzed sequential alkylation-alkenylation reactions: Application towards the synthesis of polyfunctionalized fused aromatic rings. *Tetrahedron* 61 (26): 6283–6297.
- 18 Mitsudo, K., Thansandote, P., Wilhelm, T. et al. (2006). Selectively substituted thiophenes and indoles by a tandem palladium-catalyzed multicomponent reaction. *Org. Lett.* 8 (18): 3939–3942.
- 19 Shang, Y., Wu, C., Gao, Q. et al. (2021). Diversity-oriented functionalization of 2-pyridones and uracils. *Nat. Commun.* 12 (1): 2988.
- 20 Chen, S., Liu, X., Li, K. et al. (2022). Modular approach for diversity-oriented synthesis of β -fluoroalkylated arenes via Pd/norbornene cooperative catalysis. *Chem. Asian J.* 17: e202200858.
- 21 Gao, Q., Liu, Z.S., Hua, Y. et al. (2020). Alkyl tosylates as alkylating reagents in the Catellani reaction. *Synthesis (Germany)* 52 (6): 834–846.
- 22 Cheng, H.G., Wu, C., Chen, H. et al. (2018). Epoxides as alkylating reagents for the Catellani reaction. *Angew. Chem. Int. Ed.* 57 (13): 3444–3448.
- 23 Liu, Z.S., Qian, G., Gao, Q. et al. (2018). Palladium/norbornene cooperative catalysis to access tetrahydronaphthalenes and indanes with a quaternary center. *ACS Catal.* 8 (6): 4783–4788.
- 24 Liu, Z.S., Qian, G., Gao, Q. et al. (2019). 5-Norbornene-2-carboxylic acid: another catalytic mediator for Catellani-type reactions. *Tetrahedron* 75 (12): 1774–1780.
- 25 Gao, S., Qian, G., Tang, H. et al. (2019). Three-step total synthesis of Ramelteon via a Catellani strategy. *ChemCatChem* 11 (23): 5762–5765.
- 26 Wang, J., Dong, Z., Yang, C., and Dong, G. (2019). Modular and regioselective synthesis of all-carbon tetrasubstituted olefins enabled by an alkenyl Catellani reaction. *Nat. Chem.* 11 (12): 1106–1112.
- 27 Wang, J., Zhou, Y., Xu, X. et al. (2020). Entry to 1,2,3,4-tetrasubstituted arenes through addressing the “meta constraint” in the palladium/norbornene catalysis. *J. Am. Chem. Soc.* 142 (6): 3050–3059.
- 28 Bressy, C., Alberico, D., and Lautens, M. (2005). A route to annulated indoles via a palladium-catalyzed tandem alkylation/direct arylation reaction. *J. Am. Chem. Soc.* 127 (38): 13148–13149.
- 29 Blaszykowski, C., Aktoudianakis, E., Bressy, C. et al. (2006). Preparation of annulated nitrogen-containing heterocycles via a one-pot palladium-catalyzed alkylation/direct arylation sequence. *Org. Lett.* 8 (10): 2043–2045.
- 30 Martins, A., Alberico, D., and Lautens, M. (2006). Synthesis of polycyclic heterocycles via a one-pot ortho alkylation/direct heteroarylation sequence. *Org. Lett.* 8 (21): 4827–4829.

- 31 Blaszykowski, C., Aktoudianakis, E., Alberico, D. et al. (2008). A palladium-catalyzed alkylation/direct arylation synthesis of nitrogen-containing heterocycles. *J. Org. Chem.* 73 (5): 1888–1897.
- 32 Laleu, B. and Lautens, M. (2008). Synthesis of annulated 2H-indazoles and 1,2,3- and 1,2,4-triazoles via a one-pot palladium-catalyzed alkylation/direct arylation reaction. *J. Org. Chem.* 73 (22): 9164–9167.
- 33 Martins, A. and Lautens, M. (2008). A palladium-catalyzed approach to polycyclic sulfur heterocycles. *J. Org. Chem.* 73 (22): 8705–8710.
- 34 Jafarpour, F. and Ashtiani, P.T. (2009). A novel one-step synthesis of imidazo[5,1- α]isoquinolines via a tandem Pd-catalyzed alkylation-direct arylation sequence. *J. Org. Chem.* 74 (3): 1364–1366.
- 35 Jafarpour, F. and Hazrati, H. (2010). Direct synthesis of dihydrobenzo[a]carbazoles via palladium-catalyzed domino annulation of indoles. *Adv. Synth. Catal.* 352 (2–3): 363–367.
- 36 Jafarpour, F., Otaredi-Kashani, A., and Behpajoo, S. (2013). Domino o-alkylation and heteroarylation of iodoarenes: one-pot synthesis of benzocyclohepta[b]indoles. *Synthesis (Stuttg.)* 45: 1094–1098.
- 37 Gericke, K.M., Chai, D.I., and Lautens, M. (2008). The versatile role of norbornene in C-H functionalization processes: concise synthesis of tetracyclic fused pyrroles via a threefold domino reaction. *Tetrahedron* 64 (26): 6002–6014.
- 38 Aureggi, V., Davoust, M., Gericke, K.M., and Lautens, M. (2009). A convenient synthesis of indolo- and pyrrolobenzazepines via a threefold norbornene-mediated domino reaction. *Synlett* 2009: 1004–1008.
- 39 Liu, H., El-Salfiti, M., and Lautens, M. (2012). Expedient synthesis of tetrasubstituted helical alkenes by a cascade of palladium-catalyzed C-H activations. *Angew. Chem. Int. Ed.* 51 (39): 9846–9850.
- 40 Gericke, K.M., Chai, D.I., Bieler, N., and Lautens, M. (2009). The norbornene shuttle: multicomponent domino synthesis of tetrasubstituted helical alkenes through multiple C-H functionalization. *Angew. Chem. Int. Ed.* 48 (8): 1447–1451.
- 41 Liu, H., El-Salfiti, M., Chai, D.I. et al. (2012). Modular and stereoselective synthesis of tetrasubstituted helical alkenes via a palladium-catalyzed domino reaction. *Org. Lett.* 14 (14): 3648–3651.
- 42 Francis, J. (1968). Allylic strain in six-membered rings. *Chem. Rev.* 68 (4): 375–413.
- 43 Hoffmann, R.W. (1989). Allylic 1,3-strain as a controlling factor in stereoselective transformations. *Chem. Rev.* 89 (8): 1841–1860.
- 44 Bai, L., Liu, J., Hu, W. et al. (2018). Palladium/norbornene-catalyzed C–H alkylation/alkyne insertion/indole dearomatization domino reaction: assembly of spiroindolenine-containing pentacyclic frameworks. *Angew. Chem. Int. Ed.* 57 (18): 5151–5155.
- 45 Nan, J., Yuan, Y., Bai, L. et al. (2018). Highly chemoselective construction of spiro[4.5]decane-embedded polycyclic scaffolds by a palladium/norbornene-catalyzed C–H activation/arene dearomatization reaction. *Org. Lett.* 20 (23): 7731–7734.

- 46 Zhang, B.S., Wang, F., Gou, X.Y. et al. (2021). Palladium-catalyzed synthesis of tricyclic indoles via a N-S bond cleavage strategy. *Org. Lett.* 23 (19): 7518–7523.
- 47 Wu, X.X., Shen, Y., Chen, W.L. et al. (2015). Palladium-catalyzed dearomative cyclization by a norbornene-mediated sequence: a route to spiroindolenine derivatives. *Chem. Commun.* 51 (94): 16798–16801.
- 48 Lei, C., Cao, J., and Zhou, J.S. (2016). Palladium-catalyzed arylation of ketones and acetonitrile with ortho alkylation of aryl rings: *de novo* synthesis of tetralines and benzocycloheptenes. *Org. Lett.* 18 (23): 6120–6123.
- 49 Zhang, B.S., Hua, H.L., Gao, L.Y. et al. (2017). Palladium-catalyzed arene C–H activation/ketone C–H functionalization reaction: route to spirodihydroindones. *Org. Chem. Front.* 4 (7): 1376–1379.
- 50 Ye, J. and Zhou, Q. (2022). Pd/NBE/chiral amino acid cooperative catalysis for enantioselective construction of all-carbon bridged ring systems. *Chinese J. Org. Chem.* 42 (1): 307.
- 51 Xie, X.Y., Jiang, W.T., and Xiao, B. (2021). Alkyl carbagermatrane enabled synthesis of seven-membered carbocycle-fused aromatics through Catellani strategy. *Synthesis (Germany)* 53 (16): 2819–2827.
- 52 Thansandote, P., Raemy, M., Rudolph, A., and Lautens, M. (2007). Synthesis of benzannulated N-heterocycles by a palladium-catalyzed C–C/C–N coupling of bromoalkylamines. *Org. Lett.* 9 (25): 5255–5258.
- 53 Rudolph, A., Rackelmann, N., Turcotte-Savard, M.O., and Lautens, M. (2009). Application of secondary alkyl halides to a domino aryl alkylation reaction for the synthesis of aromatic heterocycles. *J. Org. Chem.* 74 (1): 289–297.
- 54 Blanchot, M., Candito, D.A., Larnaud, F., and Lautens, M. (2011). Formal synthesis of nitidine and NK109 via palladium-catalyzed domino direct arylation/N-arylation of aryl triflates. *Org. Lett.* 13 (6): 1486–1489.
- 55 Candito, D.A. and Lautens, M. (2009). Palladium-catalyzed domino direct arylation/n-arylation: convenient synthesis of phenanthridines. *Angew. Chem. Int. Ed.* 48 (36): 6713–6716.
- 56 Bin, Z.Y., Mariampillai, B., Candito, D.A. et al. (2009). Exploiting the divergent reactivity of aryl-palladium intermediates for the rapid assembly of fluorene and phenanthrene derivatives. *Angew. Chem. Int. Ed.* 48 (10): 1849–1852.
- 57 Elsayed, M.S.A., Griggs, B., and Cushman, M. (2018). Synthesis of benzo[1,6]naphthyridinones using the Catellani reaction. *Org. Lett.* 20 (17): 5228–5232.
- 58 Ferraccioli, R., Carenzi, D., Rombolà, O., and Catellani, M. (2004). Synthesis of 6-phenanthridinones and their heterocyclic analogues through palladium-catalyzed sequential aryl-aryl and N-aryl coupling. *Org. Lett.* 6 (25): 4759–4762.
- 59 Della Ca', N., Sassi, G., and Catellani, M. (2008). A direct palladium-catalyzed route to selectively substituted carbazoles through sequential C–C and C–N bond formation: synthesis of carbazomycin A. *Adv. Synth. Catal.* 350 (14–15): 2179–2182.

- 60 Maestri, G., Larraufie, M.H., Derat, É. et al. (2010). Expeditious synthesis of phenanthridines from benzylamines via dual palladium catalysis. *Org. Lett.* 12 (24): 5692–5695.
- 61 Tsukano, C., Muto, N., Enkhtaivan, I., and Takemoto, Y. (2014). Synthesis of pyrrolophenanthridine alkaloids based on C(sp³)-H and C(sp²)-H functionalization reactions. *Chem. Asian J.* 9 (9): 2628–2634.
- 62 Zhou, H., Chen, W., and Chen, Z. (2018). Pd/norbornene collaborative catalysis on the divergent preparation of heterocyclic sulfoximine frameworks. *Org. Lett.* 20 (9): 2590–2594.
- 63 Chen, D.Z., Yang, B.J., He, X.L. et al. (2019). Design, synthesis and structure-activity relationship optimization of phenanthridine derivatives as new Wnt/ β -catenin signalling pathway agonists. *Bioorg. Chem.* 84: 285–294.
- 64 Liu, Z.S., Xie, P.P., Hua, Y. et al. (2021). An axial-to-axial chirality transfer strategy for atroposelective construction of C–N axial chirality. *Chem* 7 (7): 1917–1932.
- 65 Chen, D.Z., Fan, S.R., Yang, B.J. et al. (2021). Phenanthridine derivative host heat shock cognate 70 down-regulators as porcine epidemic diarrhea virus inhibitors. *J. Nat. Prod.* 84: 1175–1184.
- 66 Wang, Y.T., Long, X.Y., Ding, X. et al. (2022). Novel nucleocapsid protein-targeting phenanthridine inhibitors of SARS-CoV-2. *Eur. J. Med. Chem.* 227: 113966.
- 67 Motti, E., Della, C.N., Xu, D. et al. (2012). A sequential Pd/norbornene-catalyzed process generates o-biaryl carbaldehydes or ketones via a redox reaction or 6 h-dibenzopyrans by C–O ring closure. *Org. Lett.* 14 (22): 5792–5795.
- 68 Motti, E., Della Ca, N., Xu, D. et al. (2013). Competitive pathways in Pd-catalyzed synthesis of arylphenols. *Tetrahedron* 69 (22): 4421–4428.
- 69 Yamamoto, Y., Murayama, T., Jiang, J. et al. (2018). The vinylogous Catellani reaction: a combined computational and experimental study. *Chem. Sci.* 9 (5): 1191–1199.
- 70 Hua, Y., Liu, Z.S., Xie, P.P. et al. (2021). Kinetic resolution of tertiary benzyl alcohols via palladium/chiral norbornene cooperative catalysis. *Angew. Chem. Int. Ed.* 60 (23): 12824–12828.
- 71 Liu, Z.S., Hua, Y., Gao, Q. et al. (2020). Construction of axial chirality via palladium/chiral norbornene cooperative catalysis. *Nat Catal.* 3 (9): 727–733.
- 72 Fu, W.C. and Kwong, F.Y. (2020). A denitrogenative palladium-catalyzed cascade for regioselective synthesis of fluorenes. *Chem. Sci.* 11 (5): 1411–1417.
- 73 Dong, Z., Wang, J., Ren, Z., and Dong, G. (2015). Ortho C–H acylation of aryl iodides by palladium/norbornene catalysis. *Angew. Chem. Int. Ed.* 54 (43): 12664–12668.
- 74 Wang, J., Zhang, L., Dong, Z., and Dong, G. (2016). Reagent-enabled ortho-alkoxycarbonylation of aryl iodides via palladium/norbornene catalysis. *Chem* 1 (4): 581–591.

- 75 Xu, S., Jiang, J., Ding, L. et al. (2018). Palladium/norbornene-catalyzed ortho aliphatic acylation with mixed anhydride: selectivity and reactivity. *Org. Lett.* 20 (2): 325–328.
- 76 Wu, Z. and Dong, G. (2022). Rapid access to multisubstituted acrylamides from cyclic ketones via palladium/norbornene cooperative catalysis. *Angew. Chem. Int. Ed.* 61 (17): e202201239.
- 77 Li, X., Pan, J., Song, S., and Jiao, N. (2016). Pd-catalyzed dehydrogenative annulation approach for the efficient synthesis of phenanthridinones. *Chem. Sci.* 7 (8): 5384–5389.
- 78 Wang, J., Qin, C., Lumb, J.P., and Luan, X. (2020). Regioselective synthesis of polyfunctional arenes by a 4-component Catellani reaction. *Chem* 6 (8): 2097–2109.
- 79 Wang, Z., Li, T., Zhao, J. et al. (2018). Expedient synthesis of 6-fluoroalkyl-phenanthridines via palladium-catalyzed norbornene-mediated dehydrogenative annulation. *Org. Lett.* 20 (21): 6640–6645.
- 80 Dong, Z. and Dong, G. (2013). *Ortho* vs *ipso*: site-selective Pd and norbornene-catalyzed arene C–H amination using aryl halides. *J. Am. Chem. Soc.* 135 (49): 18350–18353.
- 81 Gao, Q., Liu, Z.S., Hua, Y. et al. (2019). A palladium/norbornene cooperative catalysis to access N-containing bridged scaffolds. *Chem. Commun.* 55 (60): 8816–8819.
- 82 Li, J., Yang, Y., Liu, Y. et al. (2021). Palladium/norbornene catalyzed ortho amination/cyclization of aryl iodide: process to 3-methyl-indole derivatives and controllable reductive elimination against the second amination. *Org. Lett.* 23 (8): 2988–2993.
- 83 Rago, A.J. and Dong, G. (2021). Synthesis of C3,C4-disubstituted indoles via the palladium/norbornene-catalyzed *ortho*-amination/*ipso*-heck cyclization. *Org. Lett.* 23 (9): 3755–3760.
- 84 Liu, J., Lin, H., Jiang, H., and Huang, L. (2022). Polysubstituted indole synthesis via palladium/norbornene cooperative catalysis of oxime esters. *Org. Lett.* 24 (2): 484–489.
- 85 Chen, S., Liu, Z.S., Yang, T. et al. (2018). The discovery of a palladium(II)-initiated Borono-Catellani reaction. *Angew. Chem. Int. Ed.* 57 (24): 7161–7165.
- 86 Wang, P., Chen, S., Zhou, Z. et al. (2019). Chemoselective Borono-Catellani arylation for unsymmetrical biaryls synthesis. *Org. Lett.* 21 (9): 3323–3327.
- 87 Gao, Y., Li, J., Bai, S. et al. (2020). Direct synthesis of annulated indoles through palladium-catalyzed double alkylations. *Org. Chem. Front.* 7 (9): 1149–1157.
- 88 Jiao, L. and Bach, T. (2011). Palladium-catalyzed direct 2-alkylation of indoles by norbornene-mediated regioselective cascade C–H activation. *J. Am. Chem. Soc.* 133 (33): 12990–12993.
- 89 Martins, A., Marquardt, U., Kasravi, N. et al. (2006). Synthesis of substituted benzoxacycles via a domino ortho-alkylation/heck coupling sequence. *J. Org. Chem.* 71 (13): 4937–4942.

- 90 Zhang, Z., Zhang, B.S., Li, K.L. et al. (2020). Palladium-catalyzed amination/dearomatization reaction of indoles and benzofurans. *J. Org. Chem.* 85 (12): 7817–7839.
- 91 Jiang, W.T., Xu, M.Y., Yang, S. et al. (2020). Alkylation-terminated Catellani reactions using alkyl carbagermatranes. *Angew. Chem. Int. Ed.* 59 (46): 20450–20454.
- 92 Thansandote, P., Chong, E., Feldmann, K.O., and Lautens, M. (2010). Palladium-catalyzed domino C-C/C-N coupling using a norbornene template: synthesis of substituted benzomorpholines, phenoxazines, and dihydrodibenzoxazepines. *J. Org. Chem.* 75 (10): 3495–3498.
- 93 Sui, X., Zhu, R., Li, G. et al. (2013). Pd-catalyzed chemoselective Catellani *ortho*-arylation of iodopyrroles: rapid total synthesis of rhazinal. *J. Am. Chem. Soc.* 135 (25): 9318–9321.
- 94 Zhao, K., Xu, S., Pan, C. et al. (2016). Catalytically asymmetric Pd/norbornene catalysis: enantioselective synthesis of (+)-rhazinal, (+)-rhazinilam, and (+)-kopsiyunnanine C1-3. *Org. Lett.* 18 (15): 3782–3785.
- 95 Ye, J., Li, L., You, Y. et al. (2023). Enantioselective assembly of ferrocenes with axial and planar chiralities via palladium/chiral norbornene cooperative catalysis. *JACS Au.* 3 (2): 384–390.
- 96 An, Y., Zhang, X.Y., Ding, Y.N. et al. (2022). Enantioselective synthesis of both axially and planar chiral ferrocenes via axial-to-planar diastereoiduction. *Org. Lett.* 24: 7294–7299.
- 97 Zuo, Z., Wang, H., Fan, L. et al. (2017). Modular assembly of spirocarbocyclic scaffolds through Pd⁰-catalyzed intermolecular dearomatizing [2+2+1] annulation of bromonaphthols with aryl iodides and alkynes. *Angew. Chem. Int. Ed.* 56 (10): 2767–2771.
- 98 Whyte, A., Olson, M.E., and Lautens, M. (2018). Palladium-catalyzed, norbornene-mediated, *ortho*-amination *ipso*-amidation: sequential C-N bond formation. *Org. Lett.* 20 (2): 345–348.
- 99 An, Y., Zhang, B.S., Zhang, Z. et al. (2020). A carboxylate-assisted amination/unactivated C(sp²)-H arylation reaction via a palladium/norbornene cooperative catalysis. *Chem. Commun.* 56: 5933–5936.
- 100 Abel-Snape, X., Whyte, A., and Lautens, M. (2020). Synthesis of aminated phenanthridinones via palladium/norbornene catalysis. *Org. Lett.* 22 (20): 7920–7925.
- 101 Zhao, S., Han, S., Du, G. et al. (2021). Acid-promoted expeditious syntheses of aminated dibenzosultams via palladium/norbornene cooperatively catalysed C–H amination/arylation. *Adv. Synth. Catal.* 363 (8): 2078–2083.
- 102 Zhang, B.S., Jia, W.Y., Gou, X.Y. et al. (2022). synthesis of c8-aminated pyrrolo-phenanthridines or-indoles via series C(sp² or sp³)-H activation and fluorescence study. *Org. Lett.* 24 (11): 2104–2109.
- 103 Zhang, B.S., Zhao, S.Y., Li, S.X. et al. (2023). Synthesis of C4-aminated carbazoles and their derivatives via Pd/NBE chemistry. *J. Org. Chem.* 88 (3): 1786–1795.

- 104 Zhang, B.S., Li, Y., An, Y. et al. (2018). Carboxylate ligand-exchanged amination/ $C(sp^3)$ -H arylation reaction via Pd/norbornene cooperative catalysis. *ACS Catal.* 8 (12): 11827–11833.
- 105 Pache, S. and Lautens, M. (2003). Palladium-catalyzed sequential alkylation-alkenylation reactions: new three-component coupling leading to oxacycles. *Org. Lett.* 5 (25): 4827–4830.
- 106 Jafarpour, F. and Lautens, M. (2006). Direct, one-step synthesis of condensed heterocycles: a palladium-catalyzed coupling approach. *Org. Lett.* 8 (16): 3601–3604.
- 107 Alberico, D. and Lautens, M. (2006). Palladium-catalyzed alkylation-alkenylation reactions: rapid access to tricyclic mescaline analogues. *Synlett* 18: 2969–2972.
- 108 Rudolph, A., Rackelmann, N., and Lautens, M. (2007). Stereochemical and mechanistic investigations of a palladium-catalyzed annulation of secondary alkyl iodides. *Angew. Chem. Int. Ed.* 46 (9): 1485–1488.
- 109 Alberico, D., Rudolph, A., and Lautens, M. (2007). Synthesis of tricyclic heterocycles via a tandem aryl alkylation/Heck coupling sequence. *J. Org. Chem.* 72 (3): 775–781.
- 110 Thansandote, P., Gouliaras, C., Turcotte-Savard, M.O., and Lautens, M. (2009). A rapid approach to the synthesis of highly functionalized tetrahydroisoquinolines. *J. Org. Chem.* 74 (4): 1791–1793.
- 111 Wang, J., Wang, H., Wang, Z. et al. (2021). Trifunctionalization of aryl iodides with two distinct nitrogen and carbon electrophiles by palladium/norbornene catalysis. *Chin. J. Chem.* 39 (10): 2659–2667.
- 112 Shen, Y., Wu, X.X., Chen, S. et al. (2018). Synthesis of polyfluoroarene-substituted benzofuran derivatives: Via cooperative Pd/Cu catalysis. *Chem. Commun.* 54 (18): 2256–2259.
- 113 Mariampillai, B., Alberico, D., Bidau, V., and Lautens, M. (2006). Synthesis of polycyclic benzonitriles via a one-pot aryl alkylation/cyanation reaction. *J. Am. Chem. Soc.* 128 (45): 14436–14437.
- 114 Mariampillai, B., Alliot, J., Li, M., and Lautens, M. (2007). A convergent synthesis of polysubstituted aromatic nitriles via palladium-catalyzed C-H functionalization. *J. Am. Chem. Soc.* 129 (49): 15372–15379.
- 115 Wu, X.X., Zhou, P.X., Wang, L.J. et al. (2014). Synthesis of polycyclic substituted vinylarenes via a one-pot intramolecular aryl alkylation-N-tosylhydrazone insertion reaction. *Chem. Commun.* 50 (29): 3882–3884.
- 116 Stille, J.K. (1985). Oxidative addition and reductive elimination. In: *The Metal—Carbon Bond (1985)*, 625–787. (PATAI'S Chemistry of Functional Groups).
- 117 Milstein, D. and K. Stille J. (1979). Mechanism of reductive elimination. Reaction of alkylpalladium(II) complexes with tetraorganotin, organolithium, and Grignard reagents. Evidence for palladium(IV) intermediacy. *J. Am. Chem. Soc.* 101 (17): 4981–4991.

- 118 Milstein, D. and K. Stille J. (1979). Palladium-catalyzed coupling of tetraorganotin compounds with aryl and benzyl halides. Synthetic utility and mechanism. *J. Am. Chem. Soc.* 101 (17): 4992–4998.
- 119 Catellani, M., Motti, E., and Baratta, S. (2001). A novel palladium-catalyzed synthesis of phenanthrenes from ortho-substituted aryl iodides and diphenyl- or alkylphenylacetylenes. *Org. Lett.* 3 (23): 3611–3614.
- 120 Zhao, Q., Fu, W.C., and Kwong, F.Y. (2018). Palladium-catalyzed regioselective aromatic extension of internal alkynes through a norbornene-controlled reaction sequence. *Angew. Chem. Int. Ed.* 57 (13): 3381–3385.
- 121 Yang, Y., Zhou, B., Zhu, X. et al. (2018). Palladium-catalyzed synthesis of triphenylenes via sequential C-H activation and decarboxylation. *Org. Lett.* 20 (17): 5402–5405.
- 122 Zhang, B.S., Wang, F., Yang, Y.H. et al. (2020). Synthesis of C4-substituted indoles via a Catellani and C-N bond activation strategy. *Org. Lett.* 22 (21): 8267–8271.
- 123 Candito, D.A. and Lautens, M. (2010). Exploiting the chemistry of strained rings: Synthesis of indoles via domino reaction of aryl iodides with 2H-azirines. *Org. Lett.* 12 (15): 3312–3315.
- 124 Hassner, A., Bunnell, C.A., and Haltiwanger, K. (1978). Chemistry of small rings. 23. Synthesis and structure of stable metal-coordinated 1-azirines. *J. Org. Chem.* 43 (1): 57–61.
- 125 Izumi, T. and Alper, H. (1982). Metal-catalyzed reactions of allylazirines. *Organometallics* 1 (2): 322–325.
- 126 Alper, H., Perera, C.P., and Ahmed, F.R. (1981). A novel synthesis of .beta.-lactams. *J. Am. Chem. Soc.* 103 (5): 1289–1291.
- 127 Alper, H. and Wollowitz, S. (1975). Group 6 metal carbonyl induced dimerization of azirines. *J. Am. Chem. Soc.* 97 (12): 3541–3543.
- 128 Isomura, K., Uto, K., and Taniguchi, H. (1977). Palladium(II)-catalysed formation of indoles from 2,2-diphenyl-2H-azirines. *J. Chem. Soc. Chem. Commun.* 19: 664–665.
- 129 Inada, A. and Hefmgartner, H. (1982). Übergangsmetall-katalysierte addition-sreaktionen von 3-phenyl-2H-azirinen und acetylen-carbonsäureestern. *Helv. Chim. Acta* 65 (5): 1489–1498.
- 130 Liu, C., Liang, Y., Zheng, N. et al. (2018). Synthesis of indolines: via a palladium/norbornene-catalyzed reaction of aziridines with aryl iodides. *Chem. Commun.* 54 (27): 3407–3410.
- 131 Li, R. and Dong, G. (2018). Direct annulation between aryl iodides and epoxides through palladium/norbornene cooperative catalysis. *Angew. Chem. Int. Ed.* 57 (6): 1697–1701.
- 132 Wu, C., Cheng, H.G., Chen, R. et al. (2018). Convergent syntheses of 2,3-dihydrobenzofurans: via a Catellani strategy. *Org. Chem. Front.* 5 (17): 2533–2536.
- 133 Li, R., Liu, F., and Dong, G. (2018). Palladium-catalyzed asymmetric annulation between aryl iodides and racemic epoxides using a chiral norbornene cocatalyst. *Org. Chem. Front.* 5 (21): 3108–3112.

- 134 Larraufie, M.H., Maestri, G., Beaume, A. et al. (2011). Exception to the ortho effect in palladium/norbornene catalysis. *Angew. Chem. Int. Ed.* 50 (51): 12253–12256.
- 135 Della Ca', N., Maestri, G., Malacria, M. et al. (2011). Palladium-catalyzed reaction of aryl iodides with ortho-bromoanilines and norbornene/norbornadiene: unexpected formation of dibenzoazepine derivatives. *Angew. Chem. Int. Ed.* 50 (51): 12257–12261.
- 136 Casnati, A., Fontana, M., Coruzzi, G. et al. (2018). Enhancing reactivity and selectivity of aryl bromides: a complementary approach to dibenzo[b,f]azepine derivatives. *ChemCatChem* 10 (19): 4346–4352.
- 137 Zheng, H., Zhu, Y., and Shi, Y. (2014). Palladium(0)-catalyzed heck reaction/C-H activation/amination sequence with diaziridinone: a facile approach to indolines. *Angew. Chem. Int. Ed.* 53 (42): 11280–11284.
- 138 Jafarpour, F., Jalalimanesh, N., Teimouri, M., and Shamsianpour, M. (2015). Palladium/norbornene chemistry: an unexpected route to methanocarbazole derivatives via three Csp³-Csp²/Csp³-N/Csp²-N bond formations in a single synthetic sequence. *Chem. Commun.* 51 (1): 225–228.
- 139 Hao, T.T., Liang, H.R., Ou-Yang, Y.H. et al. (2018). Palladium-catalyzed domino reaction for stereoselective synthesis of multisubstituted olefins: construction of blue luminogens. *J. Org. Chem.* 83 (8): 4441–4454.
- 140 Wu, F., Wang, H., and Chen, W. (2019). Synthesis and characterization of palladium amido complexes containing pincer CNO ligands through nitrene insertion. *Appl. Organomet. Chem.* 33 (3): e4775.
- 141 Ji, X., Gu, Y., Cheng, C. et al. (2020). Palladium-catalyzed three-component reactions for the synthesis of norbornane-fused indanes. *Adv. Synth. Catal.* 362 (7): 1496–1501.
- 142 Zhu, W.Q., Zhang, Z.W., Han, W.Y. et al. (2021). Aziridine used as a vinylidene unit in palladium-catalyzed [2 + 2 + 1] domino annulation. *Org. Chem. Front.* 8 (13): 3413–3420.
- 143 Zhu, W.Q., Fang, Y.C., Han, W.Y. et al. (2021). Palladium-catalyzed [2 + 2 + 1] annulation: access to chromone fused cyclopentanones with cyclopropenone as the CO source. *Org. Chem. Front.* 8 (12): 3082–3090.
- 144 Li, F., Li, H.M., Xiu, R.F. et al. (2022). Palladium-catalyzed domino reaction for the assembly of norbornane-containing chromones with dimethyl squarate as the solid C1 source. *Org. Lett.* 24: 9392–9397.
- 145 Jiao, L., Herdtweck, E., and Bach, T. (2012). Pd(II)-catalyzed regioselective 2-alkylation of indoles via a norbornene-mediated C-H activation: Mechanism and applications. *J. Am. Chem. Soc.* 134 (35): 14563–14572.
- 146 Yin, L., Guan, T., Cheng, J. et al. (2022). Manipulations of phenylnorbornyl palladium species for multicomponent construction of a bridged polycyclic privileged scaffold. *Commun. Chem.* 5 (1): 140.
- 147 Huang, X.H., Yang, C.Y., Liu, F.L. et al. (2023). Palladium/norbornene cooperative-catalyzed cascade decarboxylative cyclization of 4-iodoisoquinolin-1(2H)-ones with o-bromobenzoic acids: access to diverse

- fused phenanthridinones and hepta[1,2-c]isoquinolinones. *J. Org. Chem.* 88 (13): 8329–8344.
- 148 Vaughan Williams, R., Colvin, M.E., Tran, N. et al. (2000). Exceptionally pyramidalized olefins: a theoretical study of the cyclopropenyl fused tri-cycles tricyclo[3.2.1.0^{2,4}]oct-2(4)-ene, tricyclo[3.2.1.0^{2,4}]octa-2(4),6-diene, tricyclo[3.2.2.0^{2,4}]non-2(4)-ene, and tricyclo[3.2.2.0^{2,4}]nona-2(4),6-diene. *J. Org. Chem.* 65 (2): 562–567.
 - 149 Fan, L., Liu, J., Bai, L. et al. (2017). Rapid assembly of diversely functionalized spiroindenes by a three-component palladium-catalyzed C–H amination/phenol dearomatization domino reaction. *Angew. Chem. Int. Ed.* 56 (45): 14257–14261.
 - 150 Thansandote, P., Hulcoop, D.G., Langer, M., and Lautens, M. (2009). Palladium-catalyzed annulation of haloanilines and halobenzamides using norbornadiene as an acetylene synthon: a route to functionalized indolines, isoquinolinones, and indoles. *J. Org. Chem.* 74 (4): 1673–1678.
 - 151 Zhang, B.S., Li, Y., Zhang, Z. et al. (2019). Synthesis of C4-aminated indoles via a Catellani and retro-Diels-Alder strategy. *J. Am. Chem. Soc.* 141 (24): 9731–9738.
 - 152 An, Y., Zhang, B.S., Ding, Y.N. et al. (2021). Palladium-catalyzed C-H glycosylation and retro Diels-Alder tandem reaction via structurally modified norbornadienes (smNBDs). *Chem. Sci.* 12 (39): 13144–13150.
 - 153 An, Y., Li, Y., Zhang, X.Y. et al. (2021). Palladium-catalyzed C-H amination/[2 + 3] or [2 + 4] cyclization via C(sp³ or sp²)-H activation. *Org. Lett.* 23 (20): 7961–7965.
 - 154 Sun, M., Chen, X., Feng, Z. et al. (2021). A Catellani and: retro-Diels-Alder strategy to access 1-amino phenanthrenes via ortho – and interannular C-H activation of 2-iodobiphenyls. *Org. Chem. Front.* 8 (23): 6535–6540.
 - 155 Fu, W.C., Wang, Z., Chan, W.T.K. et al. (2017). Regioselective synthesis of polycyclic and heptagon-embedded aromatic compounds through a versatile π -extension of aryl halides. *Angew. Chem. Int. Ed.* 56 (25): 7166–7170.
 - 156 Ferraccioli, R., Carenzi, D., and Catellani, M. (2004). Synthesis of 1,2,3,4-tetrahydroisoquinolines and 2,3,4,5-tetrahydro-1H-2- benzazepines combining sequential palladium-catalysed ortho alkylation/ vinylation with aza-Michael addition reactions. *Tetrahedron Lett.* 45 (37): 6903–6907.
 - 157 Ferraccioli, R., Giannini, C., and Molteni, G. (2007). Stereoselective synthesis of 1,3-substituted tetrahydroisoquinolines through palladium-catalyzed three-component reaction. *Tetrahedron: Asymmetry* 18 (12): 1475–1480.
 - 158 Ferraccioli, R. and Forni, A. (2009). Selective synthesis of isoquinolin-3-one derivatives combining Pd-catalysed aromatic alkylation/vinylation with addition reactions: the beneficial effect of water. *Eur. J. Org. Chem.* 19: 3161–3166.
 - 159 Qian, G., Bai, M., Gao, S. et al. (2018). Modular one-step three-component synthesis of tetrahydroisoquinolines using a Catellani strategy. *Angew. Chem. Int. Ed.* 57 (34): 10980–10984.

- 160** Zhang, X., Wei, H., Wei, Q. et al. (2023). Modular one-step three-component assembly of isochromans via palladium/norbornene cooperative catalysis. *Asian J. Org. Chem.* 12: e202200642.
- 161** Faccini, F., Motti, E., and Catellani, M. (2004). A new reaction sequence involving palladium-catalyzed unsymmetrical aryl coupling. *J. Am. Chem. Soc.* 126 (1): 78–79.
- 162** Motti, E., Faccini, F., Ferrari, I. et al. (2006). Sequential unsymmetrical aryl coupling of o-substituted aryl iodides with o-bromophenols and reaction with olefins: palladium-catalyzed synthesis of 6H-dibenzopyran derivatives. *Org. Lett.* 8 (18): 3967–3970.
- 163** Della Ca', N., Motti, E., and Catellani, M. (2008). Palladium-catalyzed synthesis of selectively substituted phenanthridine derivatives. *Adv. Synth. Catal.* 350 (16): 2513–2516.
- 164** Narbonne, V., Retailleau, P., Maestri, G., and Malacria, M. (2014). Diastereoselective synthesis of dibenzazepines through chelation on palladium(IV) intermediates. *Org. Lett.* 16 (2): 628–631.
- 165** Xu, D., Dai, L., Catellani, M. et al. (2015). A novel enantioselective synthesis of 6H-dibenzopyran derivatives by combined palladium/norbornene and cinchona alkaloid catalysis. *Org. Biomol. Chem.* 13 (8): 2260–2263.
- 166** Della Ca, N., Motti, E., Mega, A., and Catellani, M. (2010). One-pot palladium-catalyzed synthesis of selectively substituted phenanthridines by sequential aryl-aryl and heck couplings, aza-Michael and retro-Mannich reactions. *Adv. Synth. Catal.* 352 (9): 1451–1454.
- 167** Yang, S.Y., Han, W.Y., He, C. et al. (2019). 2,2-bifunctionalization of norbornene in palladium-catalyzed domino annulation. *Org. Lett.* 21 (21): 8857–8860.
- 168** Li, M.Z., Tong, Q., Han, W.Y. et al. (2020). Synthesis of chromone-containing polycyclic compounds: via palladium-catalyzed [2 + 2 + 1] annulation. *Org. Biomol. Chem.* 18 (6): 1112–1116.