

Contents

II/20 Molecular Constants

Subvolume C: Non-linear Triatomic Molecules

Part 1: H₂O (HOH)

Part α: H₂¹⁶O (H¹⁶OH), Part I

A	Introduction	IX
I	Energy level designations	IX
I.1	Vibrational assignment	IX
I.1.1	Normal and local modes	IX
I.1.2	Polyads	XI
I.1.3	Vibrational interactions	XV
I.2	Rotational assignment	XV
I.3	Ortho-para transitions	XVI
II	Energy expressions referred to the ground state	XVI
II.1	Vibrational states	XVI
II.2	Simple expressions for the fundamental frequencies	XVI
III	Effective Hamiltonians	XVII
III.1	<i>A</i> -reduced Watson-type rotational Hamiltonian	XVII
III.2	Coudert Hamiltonian with Radau's coordinates	XIX
III.3	Tyuterev Hamiltonian with Generating Function Model	XX
III.4	Rotational Padé Hamiltonian operator	XXII
IV	Perturbation-theory free Hamiltonians	XXIV
IV.1	Jensen Morse Oscillator-Rigid Bender Internal Dynamics Hamiltonian	XXIV
IV.2	Vibrational Hamiltonian expanded in terms of local Morse operators	XXIV
V	Potential energy	XXIV
V.1	Potential Energy Function (PEF) expanded as a power series	XXV
V.2	Spectroscopically determined Potential Energy Surface (PES)	XXVI
V.2.1	Jensen's PES determined by variational calculation of rotation-vibration energies, with MORBID Hamiltonian	XXVI
V.2.2	Effective isotope-independent Born-Oppenheimer (B-O) PES with isotope-dependent adiabatic correction	XXVI
V.2.3	Isotope-dependent PES from high-quality <i>ab-initio</i> analytical potential representation	XXVII
V.2.4	Semitheoretical PES by morphing <i>ab-initio</i> potential	XXVIII
V.2.5	Correction to the <i>ab-initio</i> PES expression from [2000Kai] for the determination of the barrier height	XXVIII
V.2.6	Force constants	XXIX
VI	Dipole moment function (DMF)	XXIX
VI.1	One example of a Taylor series expansion form of the DMF	XXIX
VI.2	DMF expression in [97Cou]	XXIX
VI.3	Analytical expression of the Dipole Moment Surface in [97Par]	XXX

VI.4	Dipole matrix elements in the DMS expansion used in [2005Tot2]	XXX
VI.5	Transition moment for the bending-rotation Coudert Hamiltonian approach	XXXII
VII	Intensities	XXXII
VII.1	Line intensity	XXXII
VII.2	Band intensity	XXXIII
VII.3	Temperature dependence of the absorption	XXXIV
VII.4	Internal partition function	XXXIV
VIII	Line shape	XXXV
VIII.1	Line profiles	XXXV
VIII.1.1	Lorentz profile	XXXV
VIII.1.2	Doppler profile	XXXV
VIII.1.3	Voigt profile	XXXV
VIII.2	Collision-broadening	XXXVI
VIII.2.1	Self-broadening	XXXVI
VIII.2.2	Foreign gas broadening	XXXVI
VIII.2.3	Temperature dependence of the line broadening coefficients	XXXVI
VIII.3	Family of H ₂ O lines	XXXVI
IX	Conversion tables	XXXVII
IX.1	Conversion table for energy-related units and selected fundamental constants	XXXVII
IX.2	Intensity units and conversion table	XXXIX
X	List of symbols	XLIV
XI	Survey	XLVIII
XI.1	Band intensities	XLVIII
XI.2	Band origins	XLVIII
XI.3	Barrier to linearity	XLVIII
XI.4	Difference bands	XLVIII
XI.5	Dipole moments	XLIX
XI.6	Dissociation energy	XLIX
XI.7	Energy levels	XLIX
XI.8	Force constants	XLIX
XI.9	Hamiltonian parameters	L
XI.10	Harmonic frequencies	L
XI.11	Line positions with relative intensities	L
XI.12	Line positions with line intensity unit	LI
XI.13	Line shape related parameters	LII
XI.14	Morphing function	LII
XI.15	Partition function	LII
XI.16	Potential	LIII
XI.17	Spectroscopic parameters	LIII
XI.18	Structure	LIII
XI.19	Temperature-dependent absorption coefficient	LIII
XII	Detailed survey of some tables	LIV
B	Data	1
1	H ₂ O (HOH)	1
1.1	H ₂ ¹⁶ O (H ¹⁶ OH)	1
C	References	411