

Contents

1 Introduction — 1

- 1.1 Basic principle of spectrophotometry — 1
- 1.2 Terms and concepts behind the technique — 2
- 1.2.1 Splitting of d-orbitals — 2
- 1.3 Origin of colors in transition metal complexes — 3
- 1.4 How the nature of ligands influences the splitting — 3
- 1.5 Summary of the concept — 4
- 1.6 Methodology of spectrophotometric determination of metal complexes — 5
- 1.7 A brief outline of procedural details of spectrophotometric methods — 5
- 1.8 Analytical reagents — 6
- 1.9 Applications of spectrophotometry — 7

Section A: Spectrophotometric determination of vanadium: methods and reagents

2 Analytical reagents having oxygen as donor — 21

- 2.1 4-nitrocatechol — 21
- 2.2 2-methyl-8-quinolinol — 21
- 2.3 6-hexyl-4-(2-thiazolyazo)-resorcinol and H_2O_2 — 22
- 2.4 Salicylaldehyde-benzoylhydrazone (Sal-BH) — 22
- 2.5 Dithiophenols and aminophenols — 22
- 2.6 Pyrogallol in the presence of Safranin T — 23
- 2.7 Pyrogallol-2,3,4-trihydroxy-5-sulfoazonaphthalin — 23
- 2.8 1-(2-thiazylazo)-2-naphthol (TAN) — 23
- 2.9 Pyrocatechol violet — 23
- 2.10 N-methylethylxanthocarbamate — 24
- 2.11 Diacetyl monoxime salicyloyl-hydrazone (DMSH) — 24
- 2.12 2,3,4-trihydroxybenzoic acid with bromate — 24
- 2.13 Tannic acid — 25
- 2.14 4-(2-thiazolylazolresorcinol (TAR) and H_2O_2 — 25

3 Analytical reagents having N-S/S-O/N-S-O as donor groups — 27

- 3.1 2-hydroxy-5-bromothiophenol and aromatic amines — 27
- 3.2 2-hydroxy-3-methoxy benzaldehydethiosemicarbazone (HMBATSC) — 27
- 3.3 Acetophenone, 2'-4'-dihydroxy thiosemicarbazone — 27
- 3.4 Dibromo-p-methyl-sulfochlorophenol — 27
- 3.5 2-hydroxy-5-chlorothiophenol (HCTP) — 28
- 3.6 2-hydroxy-3-methoxybenzaldehydethiosemicarbazone (HMBATC) — 28
- 3.7 Gallamine blue (GB^+) and bromate — 28

3.8	Safranine O and Fuchsin — 28
3.9	p-Anisidine and K-bromate — 29
3.10	Xylenol orange: potassium bromate — 29
3.11	Meloxicam — 29
3.12	Chrome Azurol S — 29
3.13	Methyl violet with potassium bromate — 30
3.14	Metanil yellow dye — 30
3.15	Azorubin S and bromate — 30
4	Analytical reagents having N/N-O as donor atom(s) — 32
4.1	Diantipyrine-2-hydroxy-3-methoxy phenyl methane — 32
4.2	Diantipyrine-(2-dihydroxy) phenyl methane (DADHM) — 32
4.3	N-benzoyl-N-phenylhydroxylamine (BPHA) — 32
4.4	N,N-diethyl-p-phenylenediamine (DPD), and bromate — 32
4.5	1-[(4-Antipyril-Azo)-2-naphthol (APANOL) — 33
4.6	Dithiophenol — 33
4.7	N-hydroxy-N (2-methyl) phenyl N' (4-fluoro) phenylbenzamidine hydrochloride and thiocyanate — 33
4.8	N-cinnamohydroxamic acid — 33
4.9	2,6-dithiol-4-methylphenol and hydrophobic amines — 34
4.10	3-hydroxy-2 [3-(4-methoxy-phenyl)-1-phenyl-4-pyrazolyl]-4-oxo-4 H-1-benzopyran (HOMEPB) — 34
4.11	Acetophenone 2',4'-dihydroxy semicarbazone for extractive spectrophotometric determination of vanadium — 34
4.12	2-hydroxy-1-naphthaldehyde-p-hydroxy benzoichydrazone (HNHBH) — 34
4.13	O-aminophenol (OAP) — 35
4.14	5-Br-PADAP-H ₂ O ₂ — 35
4.15	2,4-dihydroxy acetophenone isonicotinoyl hydrazone (RPINH) — 35
4.16	4-(2-pyridylazo) resorcinol (PAR) — 36
4.17	3-hydroxy-1-(p-methoxyphenyl)-2-methylpyridine-4-one — 36
4.18	5-antipyrilazosalicylfluorone (APASF) — 36
4.19	Antipyrilazo compounds — 36
4.20	5-bromo salicylaldehyde isonicotinoyl hydrazone (5-Br SANH) — 36
4.21	3-hydroxy-2-[1'-phenyl-3'-(p-methylphenyl)-H'-pyrazolyl]-4-oxo-4H-1-benzopyran (HPMPB) — 37
4.22	1-(2-pyridylazo)-2-naphthol sensitized reagent (PAN) — 37
4.23	Di-2-pyridylketone benzoylhydrazone (dPKBH) — 37
4.24	1-(2-quinolylazo)-2,4,5-trihydroxybenzene (QATB) — 37
4.25	2,4-dinitrophenylhydrazine with imipramine hydrochloride — 38
4.26	2-methyl-8-quinolinol complex — 38
4.27	2,3-dihydroxynaphthalene — 38
4.28	di-2-pyridylketone benzoylhydrazone (diPKBH) — 38

Section B: **Spectrophotometric determination of chromium: methods and reagents**

- 5 Spectrophotometric determination of chromium using reagents having N, O, S as donor atoms — 43**
- 5.1 Diphenyl hydrazine — **43**
 - 5.2 N,N-diethyl-p-phenylenediamine (DPD) — **43**
 - 5.3 1,5-diphenylcarbazide (DPC) — **43**
 - 5.4 1,5-diphenylcarbazide — **44**
 - 5.5 2-hydroxy-5-bromothiophenol and hydrophobic amines — **44**
 - 5.6 4-(4'-chlorobenzylideneimino)-3-methyl-5-mercapto-1,2,4-triazole (CBIMMT) — **44**
 - 5.7 Diphenyl carbazide — **45**
 - 5.8 2,4-dimethyl-3H-1,5 benzodiazepine (DBA) — **45**
 - 5.9 2-(phenyl)-4,5-di (4-methoxy-phenyl) imidazole — **45**
 - 5.10 2-hydroxy-3-methoxy benzaldehyde thiosemicarbazone (HMBATSC) — **45**
 - 5.11 4-(nitro phenylazo imidazole) (NPAI) — **46**
 - 5.12 2-{[(E)-(1H-benzo[d]imidazole 2-yl) diazenyl]-5-[(E) benzylideneimino) phenol (BIADPi) — **46**
 - 5.13 2,2'-azino-bis (3-ethylbenzo thiazoline-6-sulfonate) (ABTS) — **46**
 - 5.14 2-hydroxynaphthaldehydebenzoyl-hydrazone (HNPBH) and Potassium chlorate — **46**
 - 5.15 N,N'-bis (4-methoxysalicylidene) ethylenediamine (BMSE) — **47**
 - 5.16 Diphenylcarbazide — **47**
 - 5.17 2,2'-ethylenebis (nitrilomethylidene) diphenol (HZEBNMDP) — **47**
 - 5.18 Anipryne methoxy-phenyl methane — **47**
 - 5.19 5,11,17,23-tetra [(2-ethyl acetoethoxyphenyl) (azo) phenyl] calyx [4] arene [TEAC] — **48**
 - 5.20 Diphenyl carbazide — **48**
 - 5.21 1,2,5,8 tetrahydroxyantraquinone (Qumalizarm, QINZ) — **48**
 - 5.22 Diphenylcarbazide — **48**
 - 5.23 3-methoxy-azomethine H — **49**
 - 5.24 1,3-propane diamietetraacetate — **49**
 - 5.25 Diphenylcarbazide — **49**
 - 5.26 Azocalixarene — **49**
 - 5.27 Salicylfluorone — **49**
 - 5.28 1,3-propanediamine-N,N'-diacetate-N,N'-di-3-propionate ion in aqueous acidic media — **50**
 - 5.29 Diphenylcarbazide — **50**
 - 5.30 4-(2-benzothiazolylazo)-2,2'-biphenyldiol — **50**

6 Miscellaneous reagents for spectrophotometric determination of chromium — 53

- 6.1 Bromocresol green — 53
- 6.2 Wright's stain — 53
- 6.3 Fuchsin, Safranin O — 53
- 6.4 Janus Green B — 54
- 6.5 Alizarin red or fast green FCF — 54
- 6.6 O-dianisidine- H_2O_2 — 54
- 6.7 Sodium bismuthate — 54
- 6.8 Rhodamine B (Rh B) — 55
- 6.9 Astrazone brilliant red 4G (ABR) — 55
- 6.10 Rhodamine 6G — 55
- 6.11 Persulfate in the presence of Ag (I) ions — 56
- 6.12 Alizarin red + H_2O_2 — 56
- 6.13 Celestine blue (CB^+) + H_2O_2 — 56
- 6.14 RAWL indicator — 56
- 6.15 Variamine blue — 57
- 6.16 Alizarin green + H_2O_2 — 57
- 6.17 Alizarin red — 57
- 6.18 Crystal violet and Fuchsin — 57
- 6.19 Salen — 58
- 6.20 Thionine — 58
- 6.21 Coloring condition of Cr (VI) — 58
- 6.22 Crystal violet and Rhodamine B — 59
- 6.23 Chromeazurol S — 59
- 6.24 Dimethylnedocarbocyanine dye (DIC) — 59
- 6.25 Astrazone brilliant red-4G (ABR) — 59
- 6.26 Cetyltrimethylammonium bromide — 60

Section C: Spectrophotometric determination of manganese: methods and reagents**7 Spectrophotometric determination of manganese: reagents and methods — 65**

- 7.1 3-aminophenol — 65
- 7.2 Dimercaptophenols and hydrophobic amines — 65
- 7.3 N,N-dimethyl-p-toluidine — 66
- 7.4 2,4-dimethyl-3H-1,5 benzodiazepine — 66
- 7.5 2,6-dimercaptophenol and hydrophobic amines — 66
- 7.6 2,6-dithio-4-methylphenol and amino phenols — 66
- 7.7 2-[(2-hydroxy phenylimino) methyl]-4-nitrophenol [HPIMNP] — 67

- 7.8 2-hydroxy-5-bromothiophenol and amino phenols — **67**
- 7.9 Hexamethylenetetramine — **67**
- 7.10 2,2 [prime] – dipyridyl and 2,4-dinitrobenzolasalicylic acid — **67**
- 7.11 7-(6-bromo-2-benzothiazolyazo-8-hydroxyquinoline [BrBTA8HQ] — **68**
- 7.12 1,10-phenanthroline and O-nitrobenzolasalicylic acid — **68**
- 7.13 2,4-xylidine — **68**
- 7.14 2-aminoacetophenone iso-nicotinoylhydrazone (2-AAINH) — **68**
- 7.15 m-Xylene — **69**
- 7.16 1,2,5,8-tetrahydroxyanthracene 9,10-dione (quinalizarin) — **69**
- 7.17 1,5-di (2-hydroxy-5-chlorophenyl)-3-cyano formazane — **69**
- 7.18 1-(2-pyridylazo)-2-naphthol — **69**
- 7.19 5-Br-PADAP — **70**
- 7.20 3,5-dibromo-2,4-dihydroxy acetophenone oxime — **70**
- 7.21 N,N'-bis (2-aminobenzoyl), triethylenetetramine (BABTD) — **70**
- 7.22 3,7-bis (dimethylamino)-phenothiazin-5-ium-chloride — **70**
- 7.23 2-hydroxy-4-methoxy acetophenoneoxime (HMAO) — **70**
- 7.24 1,3,3-trimethyl-2-[3-(3-methyl-3H-benzothiazol-2-ylidene)-propenyl-3H-indolium (BTIC) and 1-nitroso-2-naphthol — **71**

8 Miscellaneous reagents for spectrophotometric determination of manganese — 73

- 8.1 H₂O₂ in alkaline medium — **73**
- 8.2 Potassium periodate — **73**
- 8.3 Leucomalachite green — **73**
- 8.4 Leucoberbelin blue — **73**
- 8.5 4,7-diphenyl-1-10-phenanthroline and alizarin yellow — **74**
- 8.6 Ammonium persulphate — **74**
- 8.7 Astilbin — **74**
- 8.8 3-[(3-cholamido propyl) dimethyl ammonio]-2-hydroxy-1-propansulfonare (CHAPSO) — **74**
- 8.9 Apo-arginase and 2,3-butanedionemonoxime (DMO) — **75**
- 8.10 SCN–rhodamine B — **75**
- 8.11 Brilliant green — **75**
- 8.12 Periodate — **76**
- 8.13 Congo red and naphthol green B — **76**
- 8.14 Chromotrope2B — **76**
- 8.15 Sudan III with OP — **76**
- 8.16 Aniline blue — **77**
- 8.17 Yellow 2GR — **77**
- 8.18 Ammonium persulfate — **77**
- 8.19 3,3'-dianisole-4,4'-bis (3,5-diphenyltetrazolium chloride) — **77**
- 8.20 Brilliant green and acid chrome blue K — **78**

X — Contents

8.21	Acid fuchsin —	78
8.22	Safranine T and methyl green —	78
8.23	Eriochrome black T —	78
8.24	Tiron and H_2O_2 —	79

Index — 81