

Contents

1	Introduction	1
2	Materials and Methods	7
2.1	Embryos	7
2.2	Light Microscopy	7
2.3	Videomicrography	7
2.4	Transmission Electron Microscopy	8
2.5	Scanning Electron Microscopy	8
2.6	Dopa Reaction	8
2.7	Pterin Fluorescence	9
2.8	Staining with Alcian Blue	9
2.9	Immunohistochemistry	9
2.10	Developmental Protocols	10
2.11	Embryological Experiments	10
2.11.1	Removal of the Epidermis	10
2.11.2	Exchange of the Epidermis	10
2.11.3	Rotation of the Epidermis	10
2.11.4	Chromatophore Number	11
2.11.5	Pronephric Duct	11
2.11.6	Rotation of the Somites	11
2.11.7	Axolotl Embryo as Host	11
2.11.8	<i>Triturus</i> Embryo as Host	11
2.11.9	Younger <i>Triturus</i> Embryo as a Host	12
2.11.10	Microcarrier Experiments	12
2.11.11	Mesoderm	12
2.11.12	Heterotopic Transplantation of the Neural Crest	12
2.12	Tissue Culture	12
2.12.1	Neural Fold Explants	12
2.12.2	Epidermis and Neural Crest	13
3	Results	14
3.1	Fundamental Premises	14
3.2	Development of the Longitudinal Stripe Pattern in <i>Triturus alpestris</i>	16
3.2.1	Dispersion	16

3.2.2	Onset of Longitudinal Stripe Formation	19
3.2.3	Definite Longitudinal Stripe Pattern	19
3.3	Development of the Barred Pigment Pattern in <i>Ambystoma mexicanum</i>	21
3.3.1	Early Spreading	21
3.3.2	Dispersion	21
3.3.3	Definite Barred Pattern	25
3.4	Analysis of Factors and Mechanisms Governing Pigment Pattern Formation in <i>Triturus alpestris</i>	29
3.4.1	Pigment Cells	29
3.4.1.1	Segregation	29
3.4.1.2	Shape, Number and Behaviour	29
3.4.1.3	Cell Number and Pigment Pattern	37
3.4.2	Epidermis	39
3.4.2.1	Epidermis-Deprived Embryos and Epidermis-Crest Cultures	39
3.4.2.2	Heterotopic Exchange	41
3.4.2.3	Rotation	41
3.4.2.4	Subepidermal Extracellular Matrix	43
3.4.2.4.1	Electron Microscopic Investigations	43
3.4.2.4.2	Histochemical and Immunohistochemical Studies	45
3.4.3	Other Tissues	47
3.4.3.1	Pronephric Duct	47
3.4.3.2	Blood Vessels	50
3.4.3.3	Neural Tube, Somites	50
3.4.3.4	Mesoderm and Endoderm	52
3.4.3.5	Axolotl Embryo as Host	54
3.4.3.6	Younger <i>Triturus</i> Embryo as Host	56
3.5	Analysis of Factors and Mechanisms Governing Pigment Pattern Formation in <i>Ambystoma</i> <i>mexicanum</i>	58
3.5.1	Pigment Cells	58
3.5.1.1	Segregation	58
3.5.1.2	Cell Shape	58
3.5.1.3	Cell Number	58
3.5.2	Epidermis	59
3.5.2.1	Epidermis-Deprived Embryos	59
3.5.2.2	White Axolotl Embryos	59
3.5.2.3	Epidermal Implants	60
3.5.2.4	Microcarrier Experiments	62
3.5.3	Other Tissues	63
3.5.3.1	Blood Vessels	63
3.5.3.2	Neural Tube, Somites, Gut	63
3.5.3.3	Neural Crest at Heterotopic Sites	63
3.5.3.4	<i>Triturus</i> Embryo as Host	64

4	Discussion	65
4.1	Dopa Reaction and Pterin Fluorescence: Criticism of the Methods	65
4.2	Organization of the Neural Crest and Early Dispersion of Pigment Cells	69
4.3	Arrangement of Pigment Cells in the Trunk	71
4.3.1	Dispersion of Pigment Cells	72
4.3.2	Formation of Melanophore and Xanthophore Zones in <i>Triturus alpestris</i>	72
4.3.2.1	Cell Shape and Rearrangement of Chromatophores	72
4.3.2.2	Dorsal Stripe	73
4.3.2.3	Lateral Stripe	75
4.3.2.4	Arrangement of Melanophores and Xanthophores in Relation to Axial Cues	76
4.3.3	Bar Formation in <i>Ambystoma mexicanum</i>	80
4.4	Comparison of the Development of the Larval Pigment Pattern in <i>Triturus alpestris</i> and <i>Ambystoma mexicanum</i>	81
4.5	Conclusion and Perspectives	85
5	Summary	87
	Acknowledgements	91
	References	92
	Subject Index	100