

Contents

Figures.....	iv
Tables.....	vii
Abbreviations.....	ix
1 Introduction and objective.....	1
2 Literature review.....	3
2.1 Insect Classification – Order Diptera.....	3
2.1.1 Suborder Nematocera.....	3
2.1.1.1 Family Culicidae.....	3
2.1.1.2 Family Simuliidae.....	4
2.1.1.3 Family Ceratopogonidae.....	5
2.1.1.4 Family Psychodidae.....	6
2.1.2 Suborder Brachycera.....	6
2.1.2.1 Family Tabanidae.....	6
2.1.2.2 Family Glossinidae.....	7
2.1.2.3 Family Muscidae.....	7
2.2 Vector control.....	8
2.2.1 Chemical control – destruction of adult vectors.....	10
2.2.1.1 Insecticide-treated cattle (ITC).....	10
2.2.1.2 Insecticide-treated targets (ITT).....	11
2.2.1.3 Insecticide-treated nets (ITN).....	11
2.2.1.4 Indoor residual spraying (IRS).....	12
2.2.2 Chemical control – destruction of vectors in early developing stages.....	12
2.2.2.1 Insect growth regulators (IGR) and larvicides.....	12
2.2.2.2 Insect chemosterilants.....	13
2.2.3 Non-chemical control.....	13
2.2.3.1 Biological control.....	13
2.2.3.2 Nuclear techniques (SIT).....	14
2.2.3.3 Sanitation.....	15
2.2.3.4 Reduced host-vector interface.....	15
2.3 Entomological techniques.....	16

2.3.1 Sampling methods – mosquitoes.....	16
2.3.1.1 Hand catch.....	16
2.3.1.2 Spray sheet collection.....	16
2.3.1.3 Exit traps.....	17
2.3.1.4 Human landing catch (HLC).....	17
2.3.1.5 Baited mosquito traps.....	18
2.3.1.6 Light traps.....	19
2.3.2 Sampling methods – nuisance flies and tsetse flies.....	19
2.3.3 Dissection techniques (Culicidae).....	20
2.3.3.1 Determination of species, age and infection incrimination..	20
2.3.4 Entomological parameters in malaria vector control.....	22
3 Materials and Methods.....	25
3.1 Study site.....	25
3.2 Experimental stables.....	26
3.3 Experimental animals.....	27
3.4 Experimental nets.....	27
3.5 Experimental design.....	27
3.6 Sampling and recording methods.....	28
3.6.1 Sampling methods.....	28
3.6.1.1 Mono-conical traps (Vavoua traps).....	28
3.6.1.2 Human landing catch (HLC).....	29
3.6.1.3 Odour-baited trap (BG-Sentinel™ trap).....	30
3.6.2 Recording methods.....	31
3.6.2.1 Digital photos of selected animal body regions.....	31
3.6.2.2 Video recordings of animal defensive movements.....	31
3.7 Handling of collected insects.....	32
3.7.1 Flies.....	32
3.7.2 Mosquitoes.....	32
3.7.3 Other haematophagous insects.....	33
3.8 Entomological data analysis.....	33
3.9 Statistical analysis.....	34
4 Results.....	35
4.1 Nuisance and biting flies.....	35
4.1.1 Mono-conical traps.....	35

4.1.2 Digital photo observations.....	37
4.1.3 Video recording observations.....	39
4.2 Mosquitoes.....	42
4.2.1 Human Landing Catch (HLC).....	42
4.2.1.1 Culicines.....	43
4.2.1.2 Anophelines.....	46
4.2.1.3 Entomological data analysis.....	53
4.3 Catches with the BG-Sentinel™ trap.....	58
4.4 Evaluation of the insecticide activity persistence in the treated net.....	61
5 Discussion.....	63
5.1 Entomological monitoring.....	63
5.1.1 Impact of the nets on nuisance and biting flies	63
5.1.1.1 Assessment with mono-conical traps, photos and videos..	63
5.1.2 Impact of the nets on mosquitoes	65
5.1.2.1 Assessment with HLC (human landing catch).....	65
5.1.2.2 Assessment with BG-Sentinel™ traps.....	70
5.1.3 Comparison between HLC and BG-Sentinel™ trap.....	71
5.2 Persistence of insecticide activity in the treated net.....	72
6 Conclusions and recommendations.....	73
7 Summary.....	75
8 Zusammenfassung.....	77
9 References.....	79
Wissenschaftliche Veröffentlichungen.....	117
Acknowledgements	118
Selbständigkeitserklärung	119