

Table of contents

	Preface	3
	Abstract	5
I.	Introduction	7
II.	Material and Methods	11
III.	Characterization of style types	13
IV.	Style morphology and anatomy and some remarks on other characters used for classification in the past and present – An overview	27
	A. Basal families of Asteraceae (Barnadesioideae, Famatinanthoideae, Mutisioideae, Stifftioideae, Wunderlichioideae, Gochnatioideae, Hecastocleidoideae)	27
	I. Subfamily Barnadesioideae	29
	I.1. Single tribe Barnadesieae	29
	II. Subfamily Famatinanthoideae	29
	II.2. Single tribe Famatinantheae	29
	III. Subfamily Stifftioideae	34
	III.3. Tribe Stifftieae	34
	III.3a. Stifftieae – <i>Stifftia</i>	34
	III.3b. Stifftieae – <i>Hyaloseris</i> clade	36
	III.3c. Stifftieae – <i>Gongylolepis</i> clade	36
	III.4. Tribe Hyalideae	36
	IV. Subfamily Mutisioideae	38
	IV.5. Tribe Onoserideae	39
	IV.6. Tribe Nassauvieae	42
	IV.7. Tribe Mutisieae	45
	V. Subfamily Wunderlichioideae	49
	V.8. Single tribe Wunderlichieae	49
	VI. Subfamily Gochnatioideae	52
	VI.9. Single tribe Gochnatieae	52
	VII. Subfamily Hecastocleidoideae	52
	VII.10. Single tribe Hecastocleideae	52
	B. Pertyoideae, Carduoideae, and Gymnarrhenioideae	54
	VIII. Subfamily Pertyoideae	55
	VIII.11. Single tribe Pertyeae	55
	IX. Subfamily Carduoideae	57
	IX.12. Tribe Oldenburgieae	57
	IX.13. Tribe Tarchonantheae	59
	IX.14. Tribe Dicomeae	60
	IX.15. Tribe Cardueae	61
	X. Subfamily Gymnarrhenioideae	64
	X.16. Tribe Gymnarrheneae	64
	C. Vernonioideae, Cichorioideae, and Corymbioideae	69
	XI. Subfamily Vernonioideae	69
	XI.17. Tribe Arctotideae	69
	XI.17a. Subtribe Arctotidinae	70
	XI.17b. Subtribe Gorteriinae	70
	XI.17c. Tribe Arctotideae, unplaced genus: <i>Heterolepis</i>	73
	XI.18. Tribe Platycarpheae	74
	XI.19. Tribe Eremothamneae	74
	XI.20. Tribe Liabeae	77
	XI.21. Tribe Moquinieae	77
	XI.22. Tribe Vernonieae	78
	XII. Subfamily Cichorioideae	81
	XII.23. Single tribe Cichorieae	81
	XIII. Subfamily Corymbioideae	84

	XIII.24. Single tribe Corymbieae	84
D.	Subfamily Asteroideae	86
	D1. Supertribe Senecionodae	87
	XIV.25. Tribe Senecioneae	87
	D2. Supertribe Asterodae	92
	XIV.26. Tribe Calenduleae	92
	XIV.27. Tribe Gnaphalieae	95
	XIV.28. Tribe Astereae	100
	XIV.29. Tribe Anthemideae	104
	D3. Supertribe Helianthodae	107
	XIV.30. Tribe Inuleae	107
	XIV.30a. Subtribe Inulinae	109
	XIV.30b. Subtribe Plucheinae	112
	XIV.31. Tribe Athroismeae	114
	XIV.32. Tribe Feddeae	116
	XIV.33. Tribe Helenieae	116
	XIV.34. Tribe Coreopsidae	117
	XIV.35. Tribe Neurolaeneae	119
	XIV.36. Tribe Tageteae	120
	XIV.37. Tribe Chaenactideae	122
	XIV.38. Tribe Bahieae	123
	XIV.39. Tribe Polymnieae	124
	XIV.40. Tribe Heliantheae	124
	XIV.41. Tribe Millerieae	129
	XIV.42. Tribe Madieae	130
	XIV.43. Tribe Perityleae	133
	XIV.44. Tribe Eupatorieae	133
V.	The value of style characters for more detailed phylogenetic considerations	136
	1. Style characteristics in basal subfamilies of Asteraceae	136
	2. Style characteristics in Pertyoideae, Carduoideae (Oldenburgieae, Dicomeae, Cardueae), and Gymnarrhenoideae	139
	3. Style characteristics in Vernonioideae, Cichorioideae, and Corymbioideae	143
	4. Style characteristics in Asteroideae	149
	4.1. Senecionodae / Senecioneae	149
	4.2. Asterodae	150
	4.3. Helianthodae	154
	4.3.1. Basal Helianthodae: Inuleae, Athroismeae, and Feddeae	154
	4.3.2. "Heliantheae alliance"	157
	5. Stigma characters	164
	5.1. 'Non-Asteroideae' group	164
	5.2. Asteroideae clade	167
VI.	Secondary pollen presentation	169
	1. Pollen portioning and optimization of pollination	169
	2. The mechanisms	169
	3. Assignment of the mechanisms based on style characters and the position of the style tip just before pollen presentation	173
	4. Secondary pollen presentation – comparative comments on morphology and function	195
VII.	Final remarks	202
VIII.	Acknowledgements	206
IX.	References	207
X.	Appendix (Table 1)	227
	Index	254