

Contents

1	Introduction	1
1.1	Background	1
1.2	Finite-dimensional approximation	2
1.3	Four-manifolds	5
1.4	Further directions	7
1.5	Organization	7
2	Finite-dimensional approximation on 3-manifolds	9
2.1	Spectral sections	9
2.2	Connections on Hilbert bundles	12
2.3	Notation and main statements	15
2.4	Construction of spectral sections	21
2.5	Derivative of projections	34
2.6	Weighted Sobolev space	43
2.7	Proof of Theorem 2.3.3	49
3	Well-definedness	57
3.1	Variation of approximations	57
3.2	Spin^c structure for family of manifolds	68
3.3	Independence of metric	70
3.4	Variation of Sobolev norms	72
3.5	The Seiberg–Witten invariant	73
3.6	Elementary properties of $\mathcal{SWF}(Y, \mathfrak{s})$	81
4	Computation	87
4.1	Seiberg–Witten Floer homotopy type in reducible case	87
4.2	Examples	91
5	Finite-dimensional approximation on 4-manifolds	95
5.1	Construction of the relative Bauer–Furuta invariant	95
5.2	Well-definedness of the relative Bauer–Furuta invariant	103
6	Frøyshov-type invariants	111
6.1	Equivariant cohomology	111
6.2	Frøyshov-type invariant	111
6.3	K -theoretic Frøyshov invariant	120

A The Conley index and parameterized stable homotopy 129

 A.1 The unstable parameterized homotopy category 129

 A.2 The parameterized Conley index 133

 A.3 Spectra 136

Afterword: Finite-dimensional approximation in other settings 139

References 141