

Contents

1	A Variational Model on Labelled Graphs with Cusps and Crossings	1
1.1	The Reconstruction Problem	1
1.2	The Mumford–Shah Model	4
1.3	The Nitzberg–Mumford Model	6
1.4	Other Curvature-Depending Functionals	11
1.5	The Variational Model on Labelled Graphs	13
	References	21
2	Stable Maps and Morse Descriptions of an Apparent Contour	25
2.1	Stability of Maps	25
2.2	Stable Maps from a Two-Manifold to the Plane	31
2.3	Ambient Isotopies	36
2.4	Ambient Isotopic and Diffeomorphically Equivalent Apparent Contours	41
2.5	Morse Descriptions of an Apparent Contour	42
2.5.1	Genericity of Morse Lines in Case of No Cusps	44
2.5.2	Morse Lines in Case of Cusps: Markers	45
2.5.3	The Morse Description	48
2.5.4	Recovering the Shape from a Morse Description	50
	References	51
3	Apparent Contours of Embedded Surfaces	53
3.1	Three-Dimensional Scenes	53
3.1.1	Splitting of $\mathbb{R}^3$	54
3.2	Apparent Contours of Embedded Surfaces	55
3.3	The Function f_Σ	59
3.4	Labelling an Apparent Contour: The Function d_Σ	64

3.5	Ambient Isotopic and Diffeomorphically Equivalent Labelled Apparent Contours	69
3.6	Visible Contours	70
	References	72
4	Solving the Completion Problem	73
4.1	Some Concepts from Graph Theory	73
4.1.1	Contour Graphs and Visible Contour Graphs	75
4.2	Complete Contour Graphs and Labelling	77
4.3	Statement of the Completion Theorem	80
4.4	Morse Descriptions of a Visible Contour Graph	82
4.4.1	Localization	84
4.5	Proof of the Completion Theorem	84
4.5.1	Analysis at the Global Maximum and at Local Maxima	86
4.5.2	Analysis at Terminal Points	87
4.5.3	Analysis at T-Junctions	88
4.5.4	Analysis at Local Minima and at the Global Minimum	92
4.6	Examples	96
	References	100
5	Topological Reconstruction of a Three-Dimensional Scene	101
5.1	Statement of the Reconstruction Theorem	101
5.1.1	Depth-Equivalent Scenes	102
5.2	Proof of Existence	103
5.2.1	Glueing	104
5.2.2	Smooth Local Embedding of $\mathcal{T}$ in $\mathbb{R}^3$	108
5.2.3	Smooth Global Embedding of M in $\mathbb{R}^3$	112
5.2.4	Definition of the 3D-Shape	117
5.3	Proof of Uniqueness	118
5.A	Appendix	125
	References	128
6	Completeness of Reidemeister-Type Moves on Labelled Apparent Contours	131
6.1	Moves on a Labelled Apparent Contour	133
6.1.1	List of All Simple Rules	134
6.2	Stratifications and Stratified Morse Functions	136
6.2.1	Stratifications Induced by a Stable Map	137
6.3	Informal Statement	140
6.4	Rigorous Statement	142
6.5	Proof of the Completeness Theorem	147
6.6	Completeness of Moves	153
	References	155

7 Invariants of an Apparent Contour 157

7.1 Definition of $\mathfrak{B}(\text{appcon}(\varphi))$ 158

7.2 Definition of $BL(\text{appcon}(\varphi))$ 160

7.3 Coincidence Between $\mathfrak{B}(\text{appcon}(\varphi))$ and $BL(\text{appcon}(\varphi))$ 162

7.3.1 Proof of Coincidence Up to a Constant 163

7.3.2 Proof of Coincidence 167

7.4 Euler–Poincaré Characteristic of ∂E 171

7.5 Cell Complexes and Fundamental Groups 176

7.5.1 Cell Complexes 178

7.5.2 Fundamental Groups 179

7.6 Alexander Polynomials and Invariants
of Fundamental Groups 181

7.7 Free Differential Calculus 183

7.8 Links with Two Components: Deficiency One 189

7.9 Surfaces with Genus 2: Deficiency Two 191

References 192

8 Elimination of Cusps 195

8.1 Embedding Sign of a Cusp 196

8.2 Connectable Cusps in an Open Set 198

8.3 Statement of the Elimination Theorem 201

8.4 Proof of the Elimination Theorem 202

8.5 Application to Closed Embedded Surfaces 206

References 207

9 The Program “Visible” 209

9.1 An Example 209

9.2 Encoding a Morse Description of the Visible Contour 212

9.2.1 Encoding the Morse Events 213

9.2.2 Implicit Orientation 213

9.2.3 The “e” Region Marking 214

9.3 Using the Program 214

9.4 Encoding a Morse Description of the Constructed
Apparent Contour 215

9.5 Some Examples 216

Reference 223

10 The Program “Appcontour”: User’s Guide 225

10.1 An Overview of the Software 226

10.2 Region Description 230

10.2.1 Extended Arcs 231

10.2.2 Describing a Region 232

10.2.3 Completeness of the Region Description 234

10.3 Encoding an Apparent Contour with Labelling 235

10.3.1 Region Description as a Stream of Characters 235

10.3.2 Morse Description 236

10.3.3 Knot Description 238

10.4	The Rules (Reidemeister-Type Moves)	240
10.4.1	Simple Rules	240
10.4.2	A Nonlocal Effect of the B Rule	244
10.4.3	Composite Rules	244
10.4.4	Inverse Rules	247
10.5	Surgeries on Apparent Contours	248
10.5.1	Vertical Surgery	248
10.5.2	Horizontal Surgery	249
10.6	Canonical Description and Comparison	249
10.6.1	On the Isomorphism Problem for Graphs	250
10.6.2	The “Regions” Graph: $\mathcal{R}$ -Graph	250
10.6.3	The Depth-First Search of an $\mathcal{R}$ -Graph	252
10.6.4	The Canonization Procedure	253
10.6.5	Comparison of Apparent Contours	258
10.7	Fundamental Groups and Cell Complexes	258
10.7.1	Computing the Euler–Poincaré Characteristic and the Number of Connected Components	259
10.7.2	Fundamental Groups	260
10.7.3	Invariants of Finitely Presented Groups and the Alexander Polynomial	262
10.7.4	Alexander Polynomials and Alexander Ideals in Two Indeterminates	267
10.8	The Mendes Graph	274
10.9	Invariants	275
10.9.1	Euler–Poincaré Characteristic	275
10.9.2	Bennequin Invariant	275
10.9.3	Examples of Invariants Computation	276
10.10	contour Reference Guide	280
10.10.1	Informational Commands	280
10.10.2	Operating Commands	284
10.10.3	Conversion and Standardization Commands	287
10.10.4	Cell Complex and Fundamental Group Commands	288
10.10.5	Options Specific to Fundamental Group Computations	289
10.10.6	Common Options	290
10.10.7	Direct Input of a Finitely Presented Group or an Alexander Ideal	291
10.11	showcontour Reference Guide	293
10.11.1	Producing a Proper Morse Description	293
10.11.2	From the Morse description to a polygonal drawing	294
10.11.3	Discrete Optimization of the Polygonal Drawing	294
10.11.4	Dynamic Smoothing of the Polygonal	295

10.12	Using contour in Scripts	296
10.12.1	contour_interact.sh	296
10.12.2	contour_describe.sh	297
10.12.3	contour_transform.sh	299
10.13	Example: knotted Surface of Genus 2	300
10.14	Example: Knots in a Solid Torus	301
10.15	Example: Klein Bottle and the “House with Two Rooms”	304
10.16	Example: Mixed Internal/External Knot	309
10.17	Using appcontour on Apparent Contours Without Labelling	312
10.17.1	Haefliger Sphere	312
10.17.2	Boy Surface	313
10.17.3	Milnor Curve	313
10.17.4	Millett curve	314
10.17.5	Klein bottle	315
10.A	Appendix: Practical Canonization of Laurent Polynomials	317
10.A.1	One-Dimensional Support	318
10.A.2	Two-Dimensional Support	318
	References	321
11	Variational Analysis of the Model on Labelled Graphs	323
11.1	The Action Functional	324
11.1.1	Graphs with Cusps and Curvature in L^p	324
11.1.2	The Functional	325
11.1.3	A Notion of Convergence	326
11.2	Lower Semicontinuity	327
11.3	On the Lower Semicontinuous Envelope of the Action	332
11.3.1	Limits of Labellings	335
11.3.2	Sufficient Conditions: An Example	337
11.A	Appendix A: Systems of Curves	342
11.A.1	Curves of Class pwr_c^p	342
11.A.2	Systems of Curves	344
11.A.3	Parametrizations of Complete Contour Graphs	345
11.B	Appendix B: Convergence and Compactness of Systems of Curves	346
11.B.1	Convergence	347
	References	353
	Nomenclature	355
	Index	361