

Contents

1	Abstract	7
2	The Cahn-Hilliard equation with noise	9
2.1	The deterministic Cahn-Hilliard equation as a model for spinodal decomposition	9
2.2	The Cahn-Hilliard equation as a gradient flow	11
2.3	An overview over existing literature	14
2.4	The Cahn-Hilliard-Cook equation	16
2.4.1	Linear Dynamics	16
2.4.2	Pattern Analysis	19
3	An intuitive approach to the L^∞ norm of random fourier sums	23
3.1	Boundary Behavior and Sign Forcing	28
3.1.1	Random Fourier Cosine Sums with Forced Signs	28
3.1.2	The Role of the Left Endpoint of the Domain	38
3.1.3	The Behavior at the Remaining Points	41
3.2	Modeling Extreme Values	43
3.2.1	The Typical Oscillation Magnitude	43
3.2.2	A Model for the Magnitude of Extreme Values	49
3.2.3	Growth Rate of the Maximum Norm	52
3.2.4	Generalization to Higher Dimensions	60
4	Pattern size analysis	65
4.1	Ergodic Theory	65
4.2	Some results about averages	70
4.2.1	Weighted averages in one dimension	71

Contents

4.2.2	ε -dependent coefficients in one dimension	74
4.2.3	ε -dependent non-multiplicative coefficients in 2d . .	77
4.3	Weighted averages on arbitrary domains	86
4.4	Counting zeros of random sums	95
4.5	Random cosine sums model for Cahn-Hilliard	98
4.6	Specific Fourier domains	106
4.6.1	Cahn-Hilliard-like domain	106
4.6.2	Other domains	107
4.6.3	A more realistic model for Cahn-Hilliard-like spin- odal decomposition	113
4.7	Numerical Simulations	116
4.7.1	The Edelman density	116
4.7.2	The connection of number of zeros and nodal do- main size	119
4.8	Sloped lines	120
4.9	Extensions and generalizations	126
5	Code	131
5.1	Spectral Galerkin methods	131
5.2	Application of the Galerkin method on the CHC equation .	133
5.3	Documentation of the Matlab code	135
	Bibliography	139
	Acknowledgments	145