

Contents

Introduction	1
Part I Cu	11
Magnetic properties of copper(II) complex with 4-(pyridin-4-yloxy)-2,2,6,6-tetramethyl-1-piperidinoxyl ligand	13
Magnetic properties of heterobimetallic iron(II)-copper(II) compound containing ferrocenyl Schiff-base	15
Magnetic properties of bis[(1-ethoxycarbonyl)-1-cyanoeththylene-2,2-dithiolato]cuprate(II)	17
Magnetic properties of copper(II) nitrate complex with 4,5-dicyanoimidazole	18
Magnetic properties of copper(II) complex with 4,5-dicyanoimidazole	20
Magnetic properties of copper(II) bis-complex with 4,5-dicyanoimidazole	22
Magnetic properties of azide bridged copper(II) complex with 4,4'-dimethyl-2,2'-bipyridine . . .	24
Magnetic properties of copper(II) complex with 2-methyl-1-vinyl-imidazole	26
Magnetic properties of copper(II) bis-complex with 2-methyl-1-vinyl-imidazole	27
Magnetic properties of bis(hexafluoroacetylacetonato)copper(II) complex with 4-(N-<i>tert</i>-butylaminoxyl)-2,2'-bipyridine	28
Magnetic properties of copper(II) complex with oxalodihydroxamic acid	30
Magnetic properties of copper(II) bis(pyrazolide)	32
Magnetic properties of copper(II) bis(4-methylpyrazolide)	33
Magnetic properties of copper(II) bis(4-chloropyrazolide)	34
Magnetic properties of polymeric hydrogen 2,3-pyrazinedicarboxylate of copper(II)	35
Magnetic properties of 2,3-pyrazinedicarboxylate of copper(II)	37
Magnetic properties of copper(II) 1,2,4-triazolate complex	39
Magnetic properties of copper(II) complex with bis(2-benzothiazolyl)methanate	41
Magnetic properties of copper(II) complex with bis(2-benzothiazolyl)methanate	42
Magnetic properties of copper(II) complex with glycylalanine	43
Magnetic properties of copper(II) complex with glycylalanine and imidazole	44
Magnetic properties of imidazole bridged hetero bimetallic copper(II)-Zinc(II), Cu-Zn complex with glycylalanine	45
Magnetic properties of copper(II) complex with dihydrobis(tetrazolyl)borate	46
Magnetic properties of singly <i>anti-anti</i> carboxylate-bridged zig-zag chain complex of Cu(II) containing Schiff-base and hexafluoroacetylacetonate ligands	47

Magnetic properties of mixed ligand copper(II) complex with substituted pyrrolizidine and 2,4- <i>tert</i> -butylacetylacetonide	49
Magnetic properties of copper(II) complex with macrocyclic ligand	50
Magnetic properties of copper(II) complex with macrocyclic ligand	51
Magnetic properties of copper(II) complex with macrocyclic ligand	52
Magnetic properties of copper(II) complex with macrocyclic ligand	54
Magnetic properties of copper(II) complex with macrocyclic ligand	56
Magnetic properties of copper(II) complex with macrocyclic ligand	58
Magnetic properties of copper(II) complex with Schiff-base obtained from acetoacetic-2-pyridylamide and 2-aminoethanol	60
Magnetic properties of copper(II) complex with Schiff-base obtained from acetoacetic-2-pyridylamide and 3-aminopropanol	61
Magnetic properties of copper(II) complex with Schiff-base obtained from acetoacetic-2-pyridylamide and 2-aminoethanol	62
Magnetic properties of copper(II) complex with Schiff-base obtained from acetoacetic-2-pyridylamide and 2-aminoethanol	63
Magnetic properties of copper(II) complex with 19-membered pentaazamacrocyclic ligand bearing tetramide groups	64
Magnetic properties of copper(II) complex with 19-membered pentaazamacrocyclic ligand bearing tetramide groups	65
Magnetic properties of copper(II) complex with 2,3-pyruvaldehyde <i>bis</i> {N(3)-methylthiosemicarbazone}	66
Magnetic properties of copper(II) complex with 2,3-pyruvaldehyde <i>bis</i> {N(3)-ethylthiosemicarbazone}	67
Magnetic properties of copper(II) complex with 2,3-pyruvaldehyde <i>bis</i> {N(3)-dimethylthiosemicarbazone}	68
Magnetic properties of copper(II) complex with 2,3-pyruvaldehyde <i>bis</i> {piperidylthiosemicarbazone}	69
Magnetic properties of copper(II) chloro complex with 4-formylantipyrine N(4)-methylthiosemicarbazone	70
Magnetic properties of copper(II) bromo complex with 4-formylantipyrine N(4)-methylthiosemicarbazone	71
Magnetic properties of copper(II) acetato complex with 4-formylantipyrine N(4)-methylthiosemicarbazone	72
Magnetic properties of copper(II) chloro complex with 4-formylantipyrine N(4)-dimethylthiosemicarbazone	73
Magnetic properties of copper(II) bromo complex with 4-formylantipyrine N(4)-dimethylthiosemicarbazone	74
Magnetic properties of copper(II) acetato complex with 4-formylantipyrine N(4)-dimethylthiosemicarbazone	75

Magnetic properties of copper(II) complex with 4-formylantipyrine <i>N</i>(4)-dimethylthiosemicarbazone	76
Magnetic properties of copper(II) complex with 4-formylantipyrine-3-piperidylthiosemicarbazone	77
Magnetic properties of copper(II) bromo complex with 4-formylantipyrine-3-piperidylthiosemicarbazone	78
Magnetic properties of copper(II) complex with 4-formylantipyrine-3-piperidylthiosemicarbazone	79
Magnetic properties of copper(II) complex with 4-formylantipyrine-3-piperidylthiosemicarbazone	80
Magnetic properties of copper(II) chloro complex with formylpyrazine-<i>N</i>(4)-methylthiosemicarbazone	81
Magnetic properties of copper(II) bromo complex with formylpyrazine-<i>N</i>(4)-methylthiosemicarbazone	82
Magnetic properties of copper(II) chloro complex with formylpyrazine-<i>N</i>(4)-methylthiosemicarbazone	83
Magnetic properties of copper(II) bromo complex with formylpyrazine-<i>N</i>(4)-methylthiosemicarbazone	84
Magnetic properties of copper(II) chloro complex with formylpyrazine-<i>N</i>(4)-ethylthiosemicarbazone	85
Magnetic properties of copper(II) bromo complex with formylpyrazine-<i>N</i>(4)-ethylthiosemicarbazone	86
Magnetic properties of copper(II) chloro complex with formylpyrazine-<i>N</i>(4)-dimethylthiosemicarbazone	87
Magnetic properties of copper(II) bromo complex with formylpyrazine-<i>N</i>(4)-dimethylthiosemicarbazone	88
Magnetic properties of copper(II) chloro complex with formylpyrazine-<i>N</i>(4)-dimethylthiosemicarbazone	89
Magnetic properties of copper(II) bromo complex with formylpyrazine-<i>N</i>(4)-dimethylthiosemicarbazone	90
Magnetic properties of copper(II) chloro complex with formylpyrazine-3-azapanyl-thiosemicarbazone	91
Magnetic properties of copper(II) bromo complex with formylpyrazine-3-azapanyl-thiosemicarbazone	92
Magnetic properties of copper(II) chloro complex with formylpyrazine-3-azapanyl-thiosemicarbazone	93
Magnetic properties of copper(II) bromo complex with formylpyrazine-3-azapanyl-thiosemicarbazone	94
Magnetic properties of copper(II) chloride complex with formylpyrazine-{3-azabicyclo[3,2,2],nonyl}-thiosemicarbazone	95
Magnetic properties of copper(II) bromo complex with formylpyrazine-{3-azabicyclo[3,2,2],nonyl}-thiosemicarbazone	96

Magnetic properties of copper(II) complex with formylpyrazine-{3-azabicyclo[3,2,2]nonyl}-thiosemicarbazone	97
Magnetic properties of copper(II) complex with hexadentate imidazole functionalised triazamacrocycle	98
Magnetic properties of copper(II) complex with hexadentate pyrazole functionalised triazamacrocycle	99
Magnetic properties of monomeric ferrocenecarboxylatocopper(II) complex with dimethyl-imidazole	100
Magnetic properties of monomeric ferrocenecarboxylatocopper(II) complex with N-methylimidazole	101
Magnetic properties of copper(II) chloro complex with 1-hydroxypyridine-2-thione	102
Magnetic properties of mixed ligand copper(II) chloro complex with 1-hydroxypyridine-2-thione and triphenylphosphine	103
Magnetic properties of mixed ligand copper(II) chloro complex with 1-hydroxypyridine-2-thione and bis(diphenylphosphino)methane	104
Magnetic properties of mixed ligand copper(II) chloro complex with 1-hydroxypyridine-2-thione and 1,2-bis(diphenylphosphino)ethane	105
Magnetic properties of mixed ligand copper(II) chloro complex with 1-hydroxypyridine-2-thione and 1,2-bis(diphenylphosphino)propane	106
Magnetic properties of mixed ligand copper(II) chloro complex with 1-hydroxypyridine-2-thione and 1,2-bis(diphenylphosphino)butane	107
Magnetic properties of mixed ligand copper(II) bromo complex with 1-hydroxypyridine-2-thione	108
Magnetic properties of mixed ligand copper(II) bromo complex with 1-hydroxypyridine-2-thione and triphenylphosphine	109
Magnetic properties of mixed ligand copper(II) bromo complex with 1-hydroxypyridine-2-thione and bis(diphenylphosphino) methane	110
Magnetic properties of mixed ligand copper(II) bromo complex with 1-hydroxypyridine-2-thione and 1,2-bis(diphenylphosphino)ethane	111
Magnetic properties of mixed ligand copper(II) bromo complex with 1-hydroxypyridine-2-thione and 1,2-bis(diphenylphosphino)butane	112
Magnetic properties of copper(II) complex with N,N'-bis{2-phenylmethyl-thiophenyl}-methylene}ethane-1,2-diamine	113
Magnetic properties of di-μ-hydroxo-bis[diaqua(2,2'-bipyrimidine)copper(II) perchlorate dihydrate	114
Magnetic properties of copper(II) complex with pyridylmethylphosphonate ligand having nitrate counterion	116
Magnetic properties of copper(II) complex with pyridylmethylphosphonate ligand having perchlorate counterion	118
Magnetic properties of (L-prolinato)copper(II) prepared from aqueous solvent	120
Magnetic properties of (L-prolinato)copper(II) prepared from methanol solvent	121

Magnetic properties of arylazothiouracil complex of copper(II)	122
Magnetic properties 1, 3-phenylenediacetatocopper(II)	123
Magnetic properties of copper(II) complex with 12-membered Schiff-base macrocycle ligand containing pendant group	124
Magnetic properties of copper(II) complex with 12-membered Schiff-base macrocycle ligand containing pendant group	125
Magnetic properties of copper(II) complex with 12-membered Schiff-base macrocycle ligand containing pendant group	126
Magnetic properties of copper(II) chloro complex with bis(n-butyl-benzimidazolyl)ether	127
Magnetic properties of copper(II)nitrato complex with bis(n-butyl-benzimidazolyl)ether	128
Magnetic properties of copper(II) chloro complex with bis(n-butyl-benzimidazolyl)thioether	129
Magnetic properties of copper(II) nitrato complex with bis(n-butyl-benzimidazolyl)thioether	130
Magnetic properties of copper(II) complex with ethylenediamine and chlorosulphate ligands	131
Magnetic properties of copper(II) complex with N-methyl-ethylenediamine and chlorosulphate ligands	132
Magnetic properties of copper(II) complex with N-ethyl-ethylenediamine and chlorosulphate ligands	133
Magnetic properties of copper(II) complex with N, N'-dimethyl-ethylenediamine and chlorosulphate ligands	134
Magnetic properties of copper(II) complex with N, N'-diethylethylene-diamine and chlorosulphate ligands	135
Magnetic properties of copper(II) complex with malonyldihydrazide based macrocyclic ligand	136
Magnetic properties of copper(II) complex with succinyldihydrazide based macrocyclic ligand	137
Magnetic properties of copper(II) complex with phenylhydrazoacetyl-acetone isonicotinoyl hydrazone	138
Magnetic properties of copper(II) complex with hydrogen-phenylhydrazo-acetylacetone-isonicotinoylhydrazone	139
Magnetic properties of $[\text{ZnCl}_4]^{2-}$ moiety stabilized by bis(ethylenediamine)copper(II) cation	140
Magnetic properties of $[\text{CdCl}_4]^{2-}$ moiety stabilized by bis(ethylenediamine)copper(II) cation	141
Magnetic properties of $[\text{HgCl}_4]^{2-}$ moiety stabilized by bis(ethylenediamine)copper(II) cation	142
Magnetic properties of copper(II) complex with diphenyl diketone mono-thiosemicarbazone	143
Magnetic properties of adduct of dichlorosuphato-bis(1,3-propylenediamine)copper(II) with dimethyltin substrate	144

Magnetic properties of adduct of dichlorosuphato-bis(1,3-propylenediamine)copper(II) with dibutyltin substrate	145
Magnetic properties of copper(II) complex with pyridin-2-carbaldehyde pyridin-2'-ylhydrazine	146
Magnetic properties of copper(II) complex with 6-methylpyridin-2-carbaldehyde-pyridin-2'-ylhydrazine	147
Magnetic properties of copper(II) complex with 5,5-dimethylcyclohexane-1,2,3-trione-2-(<i>p</i>-chlorophenylhydrazine)	148
Magnetic properties of copper(II) complex with 5,5-dimethylcyclohexane-1,2,3-trione-2-(<i>p</i>-methoxyphenylhydrazine)	150
Magnetic properties of copper(II) complex with 5,5-dimethylcyclohexane-1,2,3-trione-2-(<i>o</i>-chlorophenylhydrazine)	152
Magnetic properties of copper(II) complex with 5,5-dimethylcyclohexane-1,2,3-trione-2-(<i>o</i>-methylphenylhydrazine)	153
Magnetic properties of copper(II) complex with 5,5-dimethylcyclohexane-1,2,3-trione-2-(<i>m</i>-methylphenylhydrazine)	154
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-N(4)-methylthiosemicarbazone	155
Magnetic properties of copper(II) complex with 6-methyl-2-acetyl-pyridine-N(4)-ethylthiosemicarbazone	156
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-N(4)-dimethylthiosemicarbazone	157
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-N(4)-diethylthiosemicarbazone	158
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-N(4)-dipropylthiosemicarbazone	159
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-3-piperidinylthiosemicarbazone	160
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-3-hexamethyleneiminylthiosemicarbazone	161
Magnetic properties of copper(II) complex with 6-methyl-2-acetylpyridine-3-hexamethyleneiminylthiosemicarbazone	162
Magnetic properties of copper(II) complex with 2-hydroxy-ζ-4-cinnamoyl acetophenone . . .	163
Magnetic properties of copper(II) complex with 2-hydroxy-ζ-4-chloro-cinnamoyl acetophenone	164
Magnetic properties of copper(II) complex with 2-hydroxy-ζ-4-methyl-cinnamoyl acetophenone	165
Magnetic properties of copper(II) complex with 2-hydroxy-ζ-4-methoxy-cinnamoyl acetophenone	166
Magnetic properties of pyridine adduct of copper(II) complex with 2-hydroxy-ζ-4-cinnamoyl acetophenone	167

Magnetic properties of pyridine adduct of copper(II) complex with 2-hydroxy- G -4-chloro-cinnamoyl acetophenone	168
Magnetic properties of pyridine adduct of copper(II) complex with 2-hydroxy- G -4-methyl-cinnamoyl acetophenone	169
Magnetic properties of pyridine adduct of copper(II) complex with 2-hydroxy- G -4-methoxy-cinnamoyl acetophenone	170
Magnetic properties of copper(II) complex with 2-mercaptobenzothiazole	171
Magnetic properties of copper(II) complex with sugar- α -amino acid as ligand	172
Magnetic properties of copper(II) complex with sugar- α -amino acid as ligand	174
Magnetic properties of copper(II) chloro complex with (4,5-dimethyl-3-pyrazolyl)aldazine	175
Magnetic properties of copper(II) bromo complex with (4,5-dimethyl-3-pyrazolyl)aldazine	176
Magnetic properties of copper(II) aquated complex with (4,5-dimethyl-3-pyrazolyl)aldazine as perchlorate salt	177
Magnetic properties of copper(II) chloro complex with (4,5-dimethyl-3-pyrazolyl)aldazine as tetrafluoroborate salt	178
Magnetic properties of copper(II) bromo complex with (4,5-dimethyl-3-pyrazolyl)aldazine as tetrafluoroborate salt	180
Magnetic properties of copper(II) complex with 1-salicyl-4-phenylthiosemicarbazide	182
Magnetic properties of copper(II) complex with cyclohexylacetohydroxamic acid	183
Magnetic properties of tetraamide macrocyclic complex of copper(II)	184
Magnetic properties of tetraamide macrocyclic complex of copper(II)	185
Magnetic properties of copper(II) complex with hydrogen N-picolinamido-N'-benzoylthiocarbamide	186
Magnetic properties of copper(II) complex with hydrogen N-picolinamido-N'-benzoylthiocarbamide	187
Magnetic properties of copper(II) chelate with 8-arylazo-6-formyl-7-hydroxy-5-methoxy-2-methylchromone	188
Magnetic properties of copper(II) complex with dithiofuroate	189
Magnetic properties of copper(II) complex with 1-methylisatin-N(4)-methylthiosemicarbazone	190
Magnetic properties of copper(II) chloro complex with 1-methylisatin-N(4)-methylthiosemicarbazone	191
Magnetic properties of copper(II) complex with 1-methylisatin-N(4)-dimethylthiosemicarbazone	192
Magnetic properties of copper(II)chloro complex with 1-methylisatin-N(4)-dimethylthiosemicarbazone	193
Magnetic properties of copper(II) complex with Schiff-base ligand containing nitrogen and sulfur donors	194

Magnetic properties of copper(II) complex with Schiff-base ligand containing nitrogen and sulfur donors	195
Magnetic properties of copper(II) complex derived from 1,2-di(phenyltellurato)-<i>o</i>-xylene	196
Magnetic properties of copper(II) complex derived from 1,2-di(<i>p</i>-tolyltellurato)-<i>o</i>-xylene	197
Magnetic properties of copper(II) complex derived from 1,2-di(4-methoxyphenyl-tellurato)-<i>o</i>-xylene	198
Magnetic properties of copper(II) complex derived from 1,2-di(4-ethoxyphenyl-tellurato)-<i>o</i>-xylene	199
Magnetic properties of copper(II) complex derived from 1,2-diphenylselenato-<i>o</i>-xylene	200
Magnetic properties of copper(II) complex derived from 1,2-di(<i>p</i>-tolylselenato)-<i>o</i>-xylene	201
Magnetic properties of copper(II) chloro complex with α-formyl pyridyl hydrazone	202
Magnetic properties of copper(II) chloro complex with α-benzoyl pyridyl hydrazone	203
Magnetic properties of copper(II) complex with fluoro and imidazole ligands	204
Magnetic properties of copper(II) complex with fluoro and imidazole ligands	205
Magnetic properties of copper(II) complex with fluoro and imidazole ligands	206
Magnetic properties of copper(II) complex with fluoro and imidazole ligands	207
Magnetic properties of copper(II) chloro complex with 4-formylantipyrine-<i>N</i>(4)-antipyrinylthiosemicarbazone	208
Magnetic properties of copper(II) bromo complex with 4-formylantipyrine-<i>N</i>(4)-antipyrinylthiosemicarbazone	209
Magnetic properties of copper(II) acetato complex with 4-formylantipyrine-<i>N</i>(4)-antipyrinylthiosemicarbazone	210
Magnetic properties of copper(II) complex with furfurylidene-nicotinamide Schiff-base	211
Magnetic properties of copper(II) complex with 1,3,4-oxadiazole	212
Magnetic properties of copper(II) complex with 1,3,4-oxadiazole	213
Magnetic properties of copper(II) chloro complex with benzoin phenylhydrazone	214
Magnetic properties of copper(II) nitrate complex with benzoin phenylhydrazone	215
Magnetic properties of copper(II) thiocyanato complex with benzoin phenylhydrazone	216
Magnetic properties of copper(II) complex with benzoin phenylhydrazone	217
Magnetic properties of copper(II) chloro complex with benzoin thiosemicarbazone	218
Magnetic properties of copper(II) complex with benzoin thiosemicarbazone	219
Magnetic properties of copper(II) complex with benzaldehyde morpholine-<i>N</i>-thiohydrazone	220
Magnetic properties of copper(II) complex with cinnamaldehyde morpholine-<i>N</i>-thiohydrazone	221
Magnetic properties of copper(II) complex with cyclohexanone-morpholine-<i>N</i>-thiohydrazone	222

Magnetic properties of copper(II) chloro complex with cyclohexanone-morpholine-N-thiohydrazone	223
Magnetic properties of copper(II) complex with isonitrosoacetophenone thiosemicarbazone	224
Magnetic properties of copper(II) nitrate complex with isonitrosoacetophenone thiosemicarbazone	225
Magnetic properties of copper(II) acetate complex with isonitrosoacetophenone thiosemicarbazone	226
Magnetic properties of copper(II) chloro complex with 4-chloroisoinitroso acetophenone thiosemicarbazone	227
Magnetic properties of copper(II) nitrate complex with 4-chloroisoinitroso acetophenone thiosemicarbazone	228
Magnetic properties of copper(II) acetate complex with 4-chloroisoinitroso acetophenone thiosemicarbazone	229
Magnetic properties of copper(II) complex with bis-heterocyclic ligand	230
Magnetic properties of copper(II) complex with bis-heterocyclic ligand	231
Magnetic properties of mixed ligand copper(II) complex with bis-(4-hydroxybenzalsemicarbazone) and pyridine	232
Magnetic properties of mixed ligand copper(II) complex with bis-(4-hydroxybenzalsemicarbazone) and 4-methylpyridine	233
Magnetic properties of mixed ligand copper(II) complex with bis-(4-hydroxybenzalsemicarbazone) and quinoline	234
Magnetic properties of mixed ligand copper(II) complex with bis-(4-hydroxybenzalsemicarbazone) and isoquinoline	235
Magnetic properties of copper(II) complex with 2-furfurylidene-<i>p</i>-nitroaniline	236
Magnetic properties of copper(II) complex with 2-furfurylidene-<i>m</i>-nitroaniline	237
Magnetic properties of coordination polymer of copper(II) with poly-Schiff-base	238
Magnetic properties of copper(II) complex with 7-(α-phenyl-α-<i>m</i>-nitroanilinomethyl)-8-quinolinol	239
Magnetic properties of copper(II) complex with 7-(α-phenyl-α-<i>p</i>-nitroanilinomethyl)-8-quinolinol	240
Magnetic properties of copper(II) complex with 2-(2-thiazolylazo)-1,3-acetylacetone	241
Magnetic properties of copper(II) complex with 2-(2-thiazolylazo)-1,3-trifluoroacetylacetone	242
Magnetic properties of copper(II) complex with 2-(2-thiazolylazo)-1,3-benzoylacetone	243
Magnetic properties of copper(II) complex with 2-(2-thiazolylazo)-1,3-thenoyltrifluoroacetylacetone	244
Magnetic properties of copper(II) complex with 2-(2-thiazolylazo)-1,3-methylacetoacetate	245
Magnetic properties of copper(II) complex with thiadiazole	246
Magnetic properties of copper(II) chelate of 3-(2-hydroxy-5-chloro-benzylideneamino)-5-methylisoxazole	247

Magnetic properties of copper(II) chelate of 3-(2-hydroxy-5-bromo-benzylideneamino)-5-methylisoxazole	248
Magnetic properties of copper(II) complex with 8-quinolyldrazonoacetylacetone	249
Magnetic properties of copper(II) complex with 8-quinolyldrazonocyclohexandione	250
Magnetic properties of copper(II) complex with chlordiazepoxide	251
Magnetic properties of copper(II) complex with chlordiazepoxide	252
Magnetic properties of copper(II) complex with chlordiazepoxide	253
Magnetic properties of copper(II) chloro complex with 2-acetylpyridine-2-aminobenzophenone-2-furoylhydrazone	254
Magnetic properties of copper(II) complex with 2-acetylpyridine-2-aminobenzophenone-2-furoylhydrazone	255
Magnetic properties of copper(II) complex with 7-(α-phenyl-α-<i>o</i>-chloroanilinomethyl)-8-quinolinol	256
Magnetic properties of copper(II) complex with 7-(α-phenyl-α-<i>m</i>-chloroanilinomethyl)-8-quinolinol	257
Magnetic properties of copper(II) polychelate with Schiff-base (4,4'-dihydroxy-3,3'-diacetylbi-phenyl-2,3-butanedione dihydrazone)	258
Magnetic properties of copper(II) complex with a ketoanil	259
Magnetic properties of copper(II) complex with a ketoanil	260
Magnetic properties of copper(II) complex with N-salicylideneglycine	261
Magnetic properties of copper(II) complex with N-salicylidene-L-alanine	262
Magnetic properties of copper(II) complex with N-salicylidene-L-phenylalanine	263
Magnetic properties of copper(II) chloro complex with 2,6-diacetylpyridine-bis(benzalhydrazone)	264
Magnetic properties of copper(II) bromo complex with 2,6-diacetylpyridine-bis(benzalhydrazone)	265
Magnetic properties of copper(II) nitrate complex with 2,6-diacetylpyridine-bis(benzalhydrazone)	266
Magnetic properties of copper(II) thiocyanato complex with 2,6-diacetylpyridine-bis(benzalhydrazone)	267
Magnetic properties of copper(II) azino complex	268
Magnetic properties of copper(II) azino complex	269
Magnetic properties of copper(II) complex with Schiff-base derived from thiophene-2-aldehyde and 4-chloroaniline	270
Magnetic properties of copper(II) complex with Schiff-base derived from thiophene-2-aldehyde and 4-bromoaniline	271
Magnetic properties of copper(II) complex with heterocyclic benzoylhydrazone	272
Magnetic properties of copper(II) complex with heterocyclic benzoylhydrazone	273

Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and chloride	274
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and chloride	275
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and thiocyanate	276
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and sulphate	277
Magnetic properties of copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline as sulphate salt	278
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and nitrate	279
Magnetic properties of copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline as nitrate salt	280
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and bromide	281
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and acetate	282
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and chloride	283
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and chloride	284
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and ammonia	285
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and ammonia	286
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and thiourea	287
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyridine and thiourea	288
Magnetic properties of mixed ligand copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyrimidine and chloride	289
Magnetic properties of mixed ligand copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyrimidine and chloride	290
Magnetic properties of mixed ligand copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyrimidine and ammonia	291
Magnetic properties of mixed ligand copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyrimidine and thiourea	292
Magnetic properties of mixed ligand copper(II) complex with Schiff-base derived from thiophene-2-glyoxal and 2-aminopyrimidine and thiourea	293

Magnetic properties of copper(II) complex with Morin	294
Magnetic properties of copper(II) complex with acetyl salicylhydrazone	295
Magnetic properties of mixed ligand complex of copper(II) with [(methylthiophen-2-yl) methylene]hydrazono-carbothioamide and thiocyanate	296
Magnetic properties of mixed ligand complex of copper(II) with [(methylthiophen-2-yl) methylene]hydrazono-carbothioamide and cyanate	297
Magnetic properties of mixed ligand complex of copper(II) with hydrazinecarbothioamide and thiocyanate	298
Magnetic properties of mixed ligand complex of copper(II) with hydrazinecarbothioamide and cyanate	299
Magnetic properties of copper(II) complex with 2-(benzothiazolyl-2'-aminoimino)-1, 2-diphenylethanone	300
Magnetic properties of copper(II) complex with isovanillin isonicotinoylhydrazone	301
Magnetic properties of copper(II) complex with 2-hydroxyacetophenoneoxime-thiourea-trioxane resin	302
Magnetic properties of copper(II) chloro complex with Schiff-base derived from condensation of 5-[4'-(nitrophenyl)azo]salicylaldehyde with thiosemicarbazone	303
Magnetic properties of copper(II) acetate complex with Schiff-base derived from condensation of 5-[4'-(nitrophenyl)azo]salicylaldehyde with thiosemicarbazone	304
Magnetic properties of copper(II) complex with Schiff-base derived from 8-acetyl-7-hydroxy-4-methylcoumarin- 4-chloroaniline	305
Magnetic properties of copper(II) complex with Schiff-base derived from 8-acetyl-7-hydroxy-4-methylcoumarin-4-nitroaniline	306
Magnetic properties of copper(II) complex with benzothiazole-2-carboxyhydrazide	307
Magnetic properties of copper(II) complex with benzothiazole-2-(1'-N-phenyl)carboxyhydrazide	308
Magnetic properties of copper(II) complex with benzothiazole-2-(1'-N-methyl)carboxyhydrazide	309
Magnetic properties of copper(II) complex with morpholine-N-thio-hydrazone of cyclohexanone	310
Magnetic properties of copper(II) complex with morpholine-N-thio-hydrazone of benzaldehyde	311
Magnetic properties of copper(II) complex with morpholine-N-thio-hydrazone of cinnamaldehyde	312
Magnetic properties of copper(II) acetate complex with Schiff-base derived from thiophene-2-aldehyde and <i>o</i> -phenylenediamine	313
Magnetic properties of copper(II) nitrate complex with Schiff-base derived from thiophene-2-aldehyde and <i>o</i> -phenylenediamine	314
Magnetic properties of copper(II) acetate complex with Schiff-base derived from thiophene-2-aldehyde and <i>m</i> -phenylenediamine	315

Magnetic properties of copper(II) nitrate complex with Schiff-base derived from thiophene-2-aldehyde and <i>m</i>-phenylenediamine	316
Magnetic properties of copper(II) chloro complex with Schiff-base derived from 2-thiophenecarboxaldehyde and 2-monopropanolamine	317
Magnetic properties of copper(II) complex with Schiff-base derived from 3-aminodibenzofuran and salicylaldehyde	318
Magnetic properties of copper(II) complex with thiophene-2-aldehyde thiosemicarbazone	319
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and salicylaldehyde	320
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and salicylaloxime	321
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and salicylaldehyde phenylhydrazone	322
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and salicylaldehyde semicarbazone	323
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and salicylaldehyde thio-semicarbazone	324
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and 8-hydroxy quinoline	325
Magnetic properties of copper(II) mixed ligand complex with thiophene-2-aldehyde thiosemicarbazone and 2-hydroxypyridine	326
Magnetic properties of copper(II) complex with 2-furfurylidene-2-aminopyridine	327
Magnetic properties of copper(II) complex with ethyl methyl ketone-2-aminopyridine	328
Magnetic properties of copper(II) complex with salicylidene-2-aminopyridine	329
Magnetic properties of copper(II) complex with salicylaldehyde-4,4-dimethyl-3-thiosemicarbazone and pyridine	330
Magnetic properties of copper(II) complex with salicylaldehyde-4,4-dimethyl-3-thiosemicarbazone	331
Magnetic properties of dinuclear copper(II) complex with 6-amino-1,3-dimethyl-5-[(2-carboxyphenyl)azo]uracil and pyridine	332
Magnetic properties of mixed ligand complex of copper(II) with <i>o</i>-hydroxybenzoic acid and hydrazine	333
Magnetic properties of copper(II) chelate(1:1) with 3-methoxy-salicylidene-2-aminothiophenol Schiff-base	334
Magnetic properties of copper(II) chelate(1:2) with 3-methoxy-salicylidene-2-aminothiophenol Schiff-base	335
Magnetic properties of copper(II) complex with bis(2-benzothiazolyl)methanate	336
Magnetic properties of copper(II) complex with bis(2-benzoxazolyl)methanate	337
Magnetic properties of copper(II) acetate complex with thymine	338

Magnetic properties of copper(II) complex with 6-(phenyl)thymine	339
Magnetic properties of dimetallic copper(II) complex with thiobarbituric acid	340
Magnetic properties of copper(II) aqua complex with glycylalanine	341
Magnetic properties of copper(II) complex with glycylalanine and imidazole	342
Magnetic properties of copper(II) chloro complex with semicarbazone of methyl-n-pentylketone	343
Magnetic properties of copper(II) bromo complex with semicarbazone of methyl-n-pentylketone	344
Magnetic properties of copper(II) nitrate complex with semicarbazone of methyl-n-pentylketone	345
Magnetic properties of copper(II) acetate complex with semicarbazone of methyl-n-pentylketone	346
Magnetic properties of copper(II) chloro complex with semicarbazone of methyl-n-hexyl ketone	347
Magnetic properties of copper(II) bromo complex with semicarbazone of methyl-n-hexyl ketone	348
Magnetic properties of copper(II) nitrate complex with semicarbazone of methyl-n-hexyl ketone	349
Magnetic properties of copper(II) acetate complex with semicarbazone of methyl-n-hexyl ketone	350
Magnetic properties of copper(II) complex with substituted diiminoquinone	351
Magnetic properties of copper(II) complex with substituted diiminoquinone	352
Magnetic properties of copper(II) complex with substituted diiminoquinone	354
Magnetic properties of copper(II) complex with substituted phenylenediamine	355
Magnetic properties of layered copper-oxyhalide perovskite	356
Magnetic properties of layered copper-oxyhalide perovskite	358
Magnetic properties of copper(II) complex with macrocyclic Schiff-base ligand	359
Magnetic properties of one-dimensional assembly of copper(II) polyhedra <i>via</i> dual use of H-bonding and π - π interaction	360
Magnetic properties of copper(II) complex with 1-ethoxycarbonyl-1-cyanoethylene-2, 2-dithiolate	362
Magnetic properties of copper(II) chloride complex with tridentate Schiff-base (6-amino-5-formyl-1, 3-dimethyluracil-benzoylhydrazine)	363
Magnetic properties of copper(II) perchlorate complex with tridentate Schiff-base (6-amino-5-formyl-1, 3-dimethyluracil-benzoylhydrazine)	364
Magnetic properties of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(phenyl)pent-4-ene-1,3-dione	365
Magnetic properties of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -chloro-phenyl)pent-4-ene-1,3-dione	366

Magnetic properties of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -methylphenyl)pent-4-ene-1,3-dione	367
Magnetic properties of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -methoxyphenyl)pent-4-ene-1,3-dione	368
Pyridine adduct of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(phenyl)pent-4-ene-1,3-dione	369
Pyridine adduct of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -chlorophenyl)pent-4-ene-1,3-dione	370
Pyridine adduct of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -methylphenyl)pent-4-ene-1,3-dione	371
Pyridine adduct of copper(II) complex with 1-(3-hydroxy-2-naphthyl)-5-(<i>p</i> -methoxyphenyl)pent-4-ene-1,3-dione	372
Magnetic properties of copper(II) chloro complex with 2-(benzylthio)ethylamine	373
Magnetic properties of copper(II) nitrate complex with 2-(benzylthio)ethylamine	374
Magnetic properties of copper(II) thiocyanato complex with 2-(benzylthio)ethylamine	375
Magnetic properties of copper(II) complex with quinquedentate macrocycle incorporating oxygen and nitrogen donor atoms	376
Magnetic properties of copper(II) complex with quinquedentate macrocycle incorporating oxygen and nitrogen donor atoms	377
Magnetic properties of copper(II) complex with quinquedentate macrocycle incorporating sulphur and nitrogen donor atoms	378
Magnetic properties of copper(II) complex with macrocycle incorporating sulphur and nitrogen donor atoms	379
Magnetic properties of copper(II) complex with macrocycle incorporating sulphur and oxygen donor atoms	380
Magnetic properties of copper(II) chloride complex with pyrrole-2-carboxaldehyde Schiff-base of <i>S</i> -methylthiocarbamate	381
Magnetic properties of copper(II) chloride complex with pyrrole-2-aldehyde Schiff-base of <i>S</i> -benzylthiocarbamate	382
Magnetic properties of copper(II) bromide complex with pyrrole-2-aldehyde Schiff-base of <i>S</i> -methylthiocarbamate	383
Magnetic properties of copper(II) bromide complex with pyrrole-2-aldehyde Schiff-base of <i>S</i> -benzylthiocarbamate	384
Magnetic properties of copper(II) complex with pyrrole-2-carboxaldehyde Schiff-base of <i>S</i> -methylthiocarbamate	385
Magnetic properties of copper(II) complex with pyrrole-2-aldehyde Schiff-base of <i>S</i> -benzylthiocarbamate	386
Magnetic properties of calcium copper formate	387
Magnetic properties of calcium copper formate	389
Magnetic properties of polymeric copper(II) complex with macrocyclic ligand	391

Magnetic properties of polymeric copper(II) complex with macrocyclic ligand	392
Magnetic properties of mononuclear copper(II) acylperoxo complex	393
Magnetic properties of mononuclear copper(II) acylperoxo complex	394
Magnetic properties of mononuclear copper(II) carboxylate complex	395
Magnetic properties of mononuclear copper(II) complex with substituted 2-propanol	396
Magnetic properties of copper chloro complex as 3-chloroanilinium salt	397
Magnetic properties of binuclear copper(II)–manganese(II) complex with <i>N,N</i> -bis(<i>N</i> -hydroxyethylaminoethyl)oxamio ligand	399
Magnetic properties of cyanide-bridged, Fe ^{III} -Cu ^{II} assembly from ferricyanide and CuN ₄ ²⁺ building blocks	401
Magnetic properties of cyanide-bridged, Fe ^{II} -Cu ^{II} assembly from ferricyanide and CuN ₄ ²⁺ building blocks	403
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	404
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	405
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	406
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	407
Magnetic properties of Zn(II)-Cu(II) chloro complex with [di(4-bromo-2-iminomethyl-6- methylaminomethylphenolate)]	408
Magnetic properties of Zn(II)-Cu(II) complex with [di(4-bromo-2-iminomethyl-6- methylaminomethylphenolate)]	409
Magnetic properties of Cu(II)-Zn(II) perchlorato complex with [di(4-bromo-2- iminomethyl-6-methylaminomethylphenolate)]	410
Magnetic properties of hetero-bimetallic, Cu-Zn, imidazolate-bridged complex with glycylalanine and imidazole	411
Magnetic properties of heterodinuclear Cu ^{II} -Zn ^{II} complex with dione dioxime	412
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	414
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	415
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	417
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	419
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	421
Magnetic properties of bimetallic, Cu(II)-Pb(II) complex with binucleating macrocyclic ligand having dissimilar coordination sites	423
Magnetic properties of copper diethylenetriamine complex containing vanadium oxide	425
Magnetic properties of layered copper-oxychloride perovskite	427

Magnetic properties of layered copper-oxybromide perovskite	429
Magnetic properties of mixed valence copper(II)–copper(I) complex with acetophenone thiosemicarbazone	430
Magnetic properties of mixed valence copper(II)–copper(I) complex with propiophenone thiosemicarbazone	431
Magnetic properties of mixed valence copper(II)–copper(I) complex with benzyl methyl thiosemicarbazone	432
Magnetic properties of mixed valence copper(II)–copper(I) complex with cyclohexanone thiosemicarbazone	433
Magnetic properties of linear heterotrinnuclear $\text{Fe}^{\text{II}}\text{Cu}^{\text{II}}\text{Cu}^{\text{II}}$ complex with 1,4,7-trimethyl-1, 4,7-triazacyclononane and asymmetric dicompartmented imine-oxime ligand	434
Magnetic properties of bis(heterodinuclear) $[\text{Cu}^{\text{II}}\text{Ni}^{\text{II}}]_2$ complex with unsymmetrical discompartmental imine-oxime ligand	436
Magnetic properties of copper(II) Schiff-base complex having binuclear units linked by isonicotinate ligand	438
Magnetic properties of dicopper(II) complex with 4-methyl-2,6-bis(pyrazol-1-ylmethyl)phenol	440
Magnetic properties of dicopper(II) complex with 2,6-bis(3,5-dimethylpyrazol-1-ylmethyl)-4-methylphenol	442
Magnetic properties of dinuclear copper(II) complex with 1,2-bis(2,2'-bipyridyl-6-yl)ethane	444
Magnetic properties of dinuclear copper(II) complex with bis-1,3-(<i>cis</i> , <i>cis</i> -1,3,5-triaminocyclohexane)xylyliene	446
Magnetic properties of binuclear copper(II) complex of 2,6-diformyl-4-methylphenol dioxime	448
Magnetic properties of binuclear copper(II) complex with 7-(2-hydroxy-2-methylpropyl)-7-aza-1,4-dithiacyclononane	450
Magnetic properties of trichloroacetato derivative of dicopper(II)	452
Magnetic properties of tribromoacetato derivative of dicopper(II)	454
Magnetic properties of dicopper complex with unsymmetrical dinucleating Schiff-base ligand	455
Magnetic properties of dicopper complex with unsymmetrical dinucleating Schiff-base ligand	457
Magnetic properties of dinuclear copper(II) chloro complex with bridging diazine ligand	459
Magnetic properties of dinuclear copper(II) bromo complex with bridging diazine ligand	461
Magnetic properties of dinuclear copper(II) chloro complex with bridging diazine ligand	463
Magnetic properties of dinuclear copper(II) oxalato complex with pyrazino[2,3- <i>f</i>][4,7]phenanthroline	465
Magnetic properties of dinuclear copper(II) sulphato complex with pyrazino-[2,3- <i>f</i>][4,7]phenanthroline	467

Magnetic properties of dinuclear copper(II) nitrate complex with pyrazino[2,3-<i>f</i>]-[4,7]phenanthroline	469
Magnetic properties of dinuclear Cu₂ chloro complex with [di(4-bromo-2-iminomethyl-6-methylaminomethylphenolate)]	471
Magnetic properties of Cu₂ complex with [di(4-bromo-2-iminomethyl-6-methylaminomethylphenolate)]	473
Magnetic properties of Cu₂ perchlorato complex with [di(4-bromo-2-iminomethyl-6-methylaminomethylphenolate)]	475
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous thiocyanate bridging ligand	477
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous cyanate bridging ligand	478
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and exogenous hydroxy bridging ligand	479
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous pivalic acid bridging ligand	480
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous <i>o</i>-toluic acid bridging ligand	481
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-ethyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous imidazole ligand	483
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-methyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous thiocyanate ligand	485
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-methyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous hydroxy ligand	486
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-methyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino-propane and exogenous acetate ligand	487
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-methyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diamino propane and exogenous pyrazolato bridging ligand	488
Magnetic properties of dinuclear copper(II) complex with N,N,N',N'-tetrakis-[(1-methyl-2-benzimidazolyl)methyl]-2-hydroxy-1,3-diaminopropane and exogenous imidazolato bridging ligand	490
Magnetic properties of dinuclear copper(II) bromide complex with N,N,N',N'-tetramethylethylenediamine bridged by oxalato ligand	492
Magnetic properties of dinuclear copper(II) perchlorate complex with bipyridine bridged by oxalato ligand	493

Magnetic properties of dinuclear copper(II) bromide complex with bipyridine bridged by oxalato ligand	494
Magnetic properties of dinuclear copper(II) perchlorate complex with N,N,N',N'-tetramethylethylenediamine bridged by oxamato ligands	495
Magnetic properties of dinuclear copper(II) bromide complex with N,N,N',N'-tetramethylethylenediamine bridged by oxamato ligands	496
Magnetic properties of dicopper(II) perchlorate complex with bipyridine bridged by oxamato ligands	497
Magnetic properties of dicopper(II) bromide complex with bipyridine bridged by oxamato ligands	498
Magnetic properties of dicopper(II) perchlorate complex with N,N,N',N'-tetramethylethylenediamine bridged by oxamidato ligand	499
Magnetic properties of dicopper(II) bromide complex with N,N,N',N'-tetramethylethylenediamine bridged by oxamidato ligand	500
Magnetic properties of dicopper(II) perchlorate complex with bipyridine bridged by oxamidato ligand	501
Magnetic properties of dicopper(II) bromide complex with bipyridine bridged by oxamidato ligand	502
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous hydroxo bridging ligand	503
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and an exogenous acetato bridging ligand	504
Magnetic properties of dinuclear copper(II) Schiff-base bridged by an alkoxo and an exogenous acetato bridging ligand	505
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous pyrazolato bridging ligand	506
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous hydroxo bridging ligand	507
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and an exogenous acetato bridging ligand	508
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous pyrazolato bridging ligand	509
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by an alkoxo and an exogenous azido ligand	510
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous hydroxo bridging ligand	512
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and an exogenous acetato bridging ligand	513
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous pyrazolato bridging ligand	514
Magnetic properties of dinuclear copper(II) Schiff-base complex bridged by alkoxo and exogenous hydroxo bridging ligand	515

Magnetic properties of dicopper(II) hydroxide tris(pyrazolide)	516
Magnetic properties of dicopper(II) hydroxide tris(4-methylpyrazolide)	517
Magnetic properties of dicopper(II) hydroxide tris(4-chloropyrazolide)	519
Magnetic properties of dicopper(II) hydroxide tris(4-bromopyrazolide)	521
Magnetic properties of dicopper(II) hydroxide tris(4-chloropyrazolide) bis(4-chloropyrazole)	523
Magnetic properties of dicopper(II) hydroxide tris(4-bromopyrazolide) bis(4-bromopyrazole)	525
Magnetic properties of copper(II) complex with 2-thiobarbituric acid	527
Magnetic properties of imidazole bridged bimetallic copper(II), Cu ₂ complex with glycylalanine	528
Magnetic properties of copper polymer with infinite chloride ion channels	529
Magnetic properties of dinuclear copper(II) complex with diethyl 2-quinolylmethyl- phosphonate ligand	531
Magnetic properties of octakis(dimethylamino)porphyrinato-dicopper(II).[60]fullerene complex	533
Magnetic properties of isophthlato bridged dinuclear copper(II) complex with 2, 2-bipyridine	535
Magnetic properties of isophthlato bridged dinuclear copper(II) complex with 1,10- phenanthroline	537
Magnetic properties of isophthlato bridged dicopper(II) complex with 4,4'-dimethyl- 2,2'-bipyridine	539
Magnetic properties of isophthlato bridged dicopper(II) complex with 5-nitro- 1,10-phenanthroline	540
Magnetic properties of dimeric copper(II) chloro complex with 4-formylantipyrine N(4)-methylthiosemicarbazone	541
Magnetic properties of dimeric copper(II) acetato complex with 4-formylantipyrine N(4)-methylthiosemicarbazone	542
Magnetic properties of dimeric copper(II) chloro complex with 4-formylantipyrine N(4)-dimethylthiosemicarbazone	543
Magnetic properties of dimeric copper(II) chloro complex with 4-formylantipyrine-3- piperidylthiosemicarbazone	544
Magnetic properties of copper(II) complex with Schiff-base cryptand	545
Magnetic properties of di- μ -hydroxo-bis[(2,2'-bipyridine)(trifluoromethanesulfonato-O) copper(II)]	547
Magnetic properties of dinuclear copper(II) complex with octaazamacrocyclic ligand	549
Magnetic properties of copper(II) complex with 6-(2-hydroxyphenyl)-2,2'-bipyridine	550
Magnetic properties of dinuclear copper(II), Cu ₂ hydroxide complex with octaamino cryptand (as perchlorate salt)	551

Magnetic properties of dicopper(II), Cu ₂ , complex with octaamino cryptand (as triflate salt)	552
Magnetic properties of dinuclear copper(II), Cu ₂ imidazolate complex with octaamino cryptand (as triflate salt)	553
Magnetic properties of dinuclear copper(II), Cu ₂ azide complex with octaamino cryptand (as triflate salt)	554
Magnetic properties of dinuclear copper(II), Cu ₂ azide complex with octaamino cryptand (as triflate salt, hydrated)	555
Magnetic properties of dinuclear copper(II) complex with phenanthroline-phenol ligand	556
Magnetic properties of dinuclear copper(II) complex with pyridine-phenol ligand	558
Magnetic properties of copper(II) ion pair complex with phenylhydrazoacetylacetone isonicotinoyl hydrazone	560
Magnetic properties of [CuCl ₄] ²⁻ moiety stabilized by bis(ethylenediamine)copper(II) cation	561
Magnetic properties of binuclear copper(II) complex with a 30-membered hexaaza-macrocyclic ligand	562
Magnetic properties of 2-hydroxy-1,3-propanediylbis(oxamato) bridged trinuclear copper(II) complex with 1, 10-phenanthroline	563
Magnetic properties of dinuclear copper(II) chloro complex with Schiff-base derived from 1,2,3,5,6,7,8,8a-octahydro-3-oxo-N,1-diphenyl-5(phenylmethylene)-2-naphthalenecarboxamide and <i>o</i> -aminophenol	565
Magnetic properties of dinuclear copper(II) chloro complex with Schiff-base derived from 1,2,3,5,6,7,8,8a-octahydro-3-oxo-N,1-diphenyl-5(phenylmethylene)-2-naphthalenecarboxamide and <i>o</i> -aminothiophenol	567
Magnetic properties of dinuclear copper(II) complex with Schiff-base derived from 1,2,3,5,6,7,8,8a-octahydro-3-oxo-N,1-diphenyl-5(phenyl-methylene)-2-naphthalenecarboxamide and <i>o</i> -aminobenzoic acid	569
Magnetic properties of dinuclear copper(II) complex with macrocyclic Schiff-base derived from highly conjugated β-diketone	571
Magnetic properties of dinuclear copper(II) complex with macrocyclic Schiff-base derived from highly conjugated β-diketone	572
Magnetic properties of dinuclear copper(II) complex with macrocyclic Schiff-base derived from highly conjugated β-diketone	573
Magnetic properties of dinuclear copper(II) complex with macrocyclic Schiff-base derived from highly conjugated β-diketone	574
Magnetic properties of dinuclear copper(II) complex with N,N'-diacetyloximo-1,3-phenylenediamine	575
Magnetic properties of dinuclear copper(II) complex with N,N'-diacetyloximo-1,3-phenylenediamine	576
Magnetic properties of dinuclear copper(II) complex with N,N'-diacetyloximo-1,3-phenylenediamine	577

Magnetic properties of dinuclear copper(II) complex with 8-arylazo-6-formyl-7-hydroxy-5-methoxy-2-methylchromone	578
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	579
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	580
Magnetic properties of copper(II) complex with a ferrocene Schiff-base derivative	581
Magnetic properties of dinuclear copper(II) bromo complex with 1-methylisatin-<i>N</i>(4)-methylthiosemicarbazone	582
Magnetic properties of dinuclear copper(II) bromo complex with 1-methylisatin-<i>N</i>(4)-dimethylthiosemicarbazone	583
Magnetic properties of chloro bridged dinuclear copper(II) complex with α-acetylpyridyl hydrazone	584
Magnetic properties of dinuclear copper(II) complex with 4-formylantipyrine-<i>N</i>(4)-antipyrinylthiosemicarbazone as tetrafluoroborate salt	585
Magnetic properties of dinuclear copper(II) complex with Schiff-base derived from 4,4'-diaminodiphenylsulphone with salicylaldehyde	586
Magnetic properties of dinuclear copper(II) complex with Schiff-base derived from 4,4'-diaminodiphenylsulphone with 2-hydroxy-1-naphthaldehyde	587
Magnetic properties of dinuclear copper(II) complex with ONNO donor azo dye	588
Magnetic properties of dinuclear copper(II) complex with chelating azo dye ligand	589
Magnetic properties of dinuclear copper(II) complex with chelating azo dye ligand	590
Magnetic properties of bimetallic copper(II) chelate with biphenyldihydrazone of 1,3-diketone	591
Magnetic properties of copper(II) complex with bis-heterocyclic ligand	592
Magnetic properties of hetero-tetrametallic, Fe₂(III)-Cu₂(II) complex with a ketoanil	593
Magnetic properties of hetero-trimetallic, Fe(III)-Cu₂(II) complex with a ketoanil	594
Magnetic properties of mixed ligand complex of copper(II) with thiophene-2-glyoxal and 2-mercaptoaniline and thiocyanate	595
Magnetic properties of mixed ligand complex of copper(II) with Schiff-base derived from thiophene-2-glyoxal and 2-mercaptoaniline and phosphate	596
Magnetic properties of binuclear copper(II) chelate with 1-(2-hydroxy-phenyl)-3-phenyl-1,3-propanedione	597
Magnetic properties of copper(II) complex with Schiff-base ligand derived from 3-formylsalicylic acid and 2-aminothiophenol	598
Magnetic properties of copper(II) complex with Schiff-base ligand derived from 3-formylsalicylic acid and 2-aminothiophenol	599
Magnetic properties of dinuclear copper(II) complex with 6-amino-1,3-dimethyl-5-[(2-carboxyphenyl)azo]uracil	600
Magnetic properties of dimeric copper(II) complex with citric acid	601
Magnetic properties of homo-bimetallic, Cu-Cu, imidazolate-bridged complex with glycylalanine and imidazole	602

Magnetic properties of dinuclear copper(II) complex with macrocyclic Schiff-base ligand	603
Magnetic properties of dinuclear copper(II) complex with <i>meso</i>-bis(phenylsulphinyl)ethane	604
Magnetic properties of dinuclear copper(II) complex with <i>racemic</i>-bis-(phenylsulphinyl)ethane	605
Magnetic properties of chain complex of dicopper(II) trimethylacetate linked by pyridyl nitronyl nitroxide	607
Magnetic properties of croconato bridged dinuclear copper(II) complex with substituted amide	609
Magnetic properties of dinuclear copper(II) complex with macrocyclic ligand	610
Magnetic properties of dinuclear copper(II) complex with macrocyclic ligand	612
Magnetic properties of dinuclear copper(II) complex with macrocyclic ligand	613
Magnetic properties of polymeric copper(II) complex with macrocyclic ligand	614
Magnetic properties of heterodinuclear Cu^{II}-Cu^{II} complex with dione dioxime	616
Magnetic properties of non-symmetric dinuclear copper(II) complex with non-symmetric dinucleating ligand	618
Magnetic properties of μ-1,1-azido-bridged dicopper(II) complex	620
Magnetic properties of μ-1,1-azido-bridged dicopper(II) complex	621
Magnetic properties of μ-1,1-azido and μ_2-bromo bridged dicopper(II) complex	622
Magnetic properties of μ-1,1-azido and μ_2-aqua bridged dicopper(II) complex	623
Magnetic properties of μ-1,1-azido and μ_2-nitrate bridged dicopper(II) complex	624
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	625
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	626
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	627
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	628
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	629
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	630
Magnetic properties of μ_2-1,1-azido bridged dicopper(II) complex	631
Magnetic properties of an oxalate-linked copper(II) coordination polymer	633
Magnetic properties of bromine-bridged copper(II) dimer with 5,7-dimethyl[1,2,4-triazolo(1,5-α)]pyrimidine	635
Magnetic properties of μ_2-1,1-bridged dinuclear copper(II) complex with diazine ligand	637
Magnetic properties of μ_2-1,1-bridged dinuclear copper(II) complex with diazine ligand	639
Magnetic properties of μ_2-1,1-bridged dinuclear copper(II) complex with diazine ligand	640
Magnetic properties of μ_2-1,1-N₃ bridged dinuclear copper(II) complex with diazine ligand	641
Magnetic properties of μ_2-1,1-N₃ bridged dinuclear copper(II) complex with diazine ligand	642

Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	643
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	644
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	646
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	648
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	650
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	652
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	653
Magnetic properties of μ_2 -1,1- N_3 bridged dinuclear copper(II) complex with diazine ligand	654
Magnetic properties of dinuclear copper(II) complex with carboxylate-rich coordination environment	656
Magnetic properties of hydroxo bridged dinuclear copper(II) complex with carboxylate-rich coordination environment	657
Magnetic properties of μ - N^1 , N^2 1,2,4-triazole and μ -sulphato bridged dinuclear copper(II) complex	658
Magnetic properties of (μ -oxalato)-bis(tetramine)dicopper(II) complex	660
Magnetic properties of (μ -oxalato)-bis(tetramine)dicopper(II) complex	661
Magnetic properties of dinuclear copper(II) complex with substituted 2-propanol	662
Magnetic properties of dimeric copper(II) complex with 3,5-di- <i>tert</i> -butylsemiquinonate	663
Magnetic properties of copper(II) dimer with hydrogen-bonded bridges	664
Magnetic properties of <i>cis</i> - μ -1,3-azide-bridged copper(II) dimer	666
Magnetic properties of di(μ -chloro) bridged dicopper(II) complex with tetradentate binucleating ligand 2,5-bis[(2-pyridylmethyl)thio]thiadiazole	667
Magnetic properties of di(μ -chloro) bridged dicopper(II) complex with tetradentate binucleating ligand 2,5-bis(2-pyridylthio)thiadiazole	669
Magnetic properties of di(μ -bromo) bridged dicopper(II) complex with tetradentate binucleating ligand 2,5-bis(2-pyridylthio)thiadiazole	670
Magnetic properties of (μ -hydroxo) bridged dicopper(II) complex with tetradentate binucleating ligand 2,5-bis(2-pyridylthio)thiadiazole	671
Magnetic properties of trinuclear copper(II) complex with formamidine ligand	672
Magnetic properties of trinuclear copper(II) complex with formamidine ligand	674
Magnetic properties of trinuclear copper(II) complex with formamidine ligand	676
Magnetic properties of three dimensional copper(II) complex with 1,2-bis(1,2,4-triazol-4-yl)ethane	678
Magnetic properties of a compound having two different (oxalato)(bipyridine) copper(II) complexes in one single crystal	680
Magnetic properties of a compound having two different (oxalato)(bipyridine) copper(II) complexes in one single crystal	682

Magnetic properties of a compound having two different (oxalato)(bipyridine) copper(II) complexes in one single crystal	683
Magnetic properties of double complex salt of cyclic diamine containing cationic copper(II) complex and a hexacyanoferrate anion	684
Magnetic properties of double complex salt of cyclic diamine containing cationic copper(II) complex and a hexacyanoferrate anion	685
Magnetic properties of double complex salt of cyclic diamine containing cationic copper(II) complex and a hexacyanoferrate anion	686
Magnetic properties of trinuclear copper(II) complex with thiazolo-2,4-pentanedione	687
Magnetic properties of trinuclear copper(II) complex with azolo-2,4-pentanedione	688
Magnetic properties of trimetallic copper(II) complex with bis-bidentate azo dye	689
Magnetic properties of trimetallic copper(II) complex with bis-bidentate azo dye	690
Magnetic properties of trimeric copper(II) chloride complex with N,N-dimethylformamide	691
Magnetic properties of copper(II) complex with nitronyl nitroxide biradical	692
Magnetic properties of angular trinuclear copper(II) complex with acetate and bipyridine ligands	694
Magnetic properties of square-planar tetranuclear copper(II)	696
Magnetic properties of tetranuclear copper(II) complex with phenyl-N,N'-bis(salicylidene) methanediamine	698
Magnetic properties of tetranuclear copper(II) complex with N,N'-bis(salicylidene) methanediamine	700
Magnetic properties of tetranuclear copper(II) complex with Schiff-base and dipyrzolate ligands	702
Magnetic properties of tetranuclear copper(II) complex with Schiff-base and dipyrzolate ligands	704
Magnetic properties of copper(II) arsenate	705
Magnetic properties of phenoxo-bridged tetranuclear Cu^{II} complex	707
Magnetic properties of tetranuclear copper(II) complex with 36-membered macrocycle	709
Magnetic properties of tetranuclear copper(II) exchange-coupled cluster encapsulated in hetropolytungastate complex	710
Magnetic properties of tetranuclear copper(II) exchange-coupled cluster encapsulated in hetropolytungastate complex	712
Magnetic properties of heteronuclear complex with Hg^{II} metal and cyclic Cu₄^{II} unit	714
Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	716
Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	718
Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	720

Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	722
Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	723
Magnetic properties of tetranuclear copper(II) complex with bis(diazine) involving a rectangular array of Cu(II) centers	725
Magnetic properties of copper(II) complex with nitronyl nitroxide biradical	727
Magnetic properties of polymeric copper(II) interconnected by 2-5-(2-pyridyl)pyrazine and acetate groups	729
Magnetic properties of tetranuclear copper(II) complex with μ-1,2,3,4-squarato coordination mode	730
Magnetic properties of tetranuclear copper(II) complex with novel dendritic poly(phthalazine) ligand	732
Magnetic properties of tetranuclear copper(II) complex with novel dendritic poly(phthalazine) ligand	734
Magnetic properties of tetranuclear copper(II) complex with novel dendritic poly(phthalazine) ligand	736
Magnetic properties of azido bridged tetranuclear copper(II) complex with substituted 2-propanol	738
Magnetic properties of copper–zirconium oxide–alkoxide species	739
Magnetic properties of (μ_4-oxo) bridged copper tetramer with 2,6-bis(morpholinomethyl)-4-methylphenol	740
Magnetic properties of (μ_4-oxo) bridged copper tetramer with 2,6-bis(morpholinomethyl)-4-methylphenol	742
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 3-hydroxy-2-naphthohydroxamic acid as supporting ligand	744
Magnetic properties of copper(II)12-metallacrown-4 (12-MC-4) having 3-hydroxy-2-salicylhydroxamic acid as supporting ligand	746
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 3,5-dideuteriosalicylhydroxamic acid as supporting ligand	747
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 2,4-dihydroxybenzohydroxamic acid as supporting ligand	748
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 4-methylsalicylhydroxamic acid as supporting ligand	749
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 2,5-dihydroxybenzohydroxamic acid as supporting ligand	750
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 5-methoxysalicylhydroxamic acid as supporting ligand	751
Magnetic properties of copper(II) 12-metallacrown-4 (12-MC-4) having 5-butylsalicylhydroxamic acid as supporting ligand	752
Magnetic properties of hexanuclear copper(II) chloro complex with glycerol phthalazine	753

Magnetic properties of hexanuclear copper(II) bromo complex with glycerol phthalazine . . .	755
Magnetic properties of hexanuclear copper(II) nitrate complex with glycerol phthalazine . . .	757
Magnetic properties of mixed metallic copper(II)–barium(II) arsenate	758
Magnetic properties of (μ-oxo)-heterometallic Y-Cu complex with 2-hydroxypyridine	759
Magnetic properties of octanuclear copper(II) chloro complex with pentaerythritol phthalazine	760
Magnetic properties of octanuclear copper(II) bromo complex with pentaerythritol phthalazine	762
Magnetic properties of polymeric copper(II) complex with tetradentate binucleating ligand 2,5-bis(2-pyridylmethylthio)thiadiazole	763
Part II Ag	765
Magnetic properties of μ-hydroxy-μ-nitritodicopper(II)–disilver(II) complex with triazacyclononane	767
Part III Au	769
Magnetic properties of gold(I) complex containing aminoxyl radical-substituted phosphine ligand	771
Magnetic properties of gold(I) complex containing aminoxyl radical-substituted phosphine ligand	773
Magnetic properties of gold(II) chloro complex with N-(2-pyridylmethyl)-2-mercaptoaniline	775
Magnetic properties of gold(II) bromo complex with N-(2-pyridylmethyl)-2-mercaptoaniline	776
Magnetic properties of gold(II) chloro complex with N-(2-pyridylmethyl)-2-mercaptoaniline	777
Magnetic properties of gold(II) bromo complex with N-(2-pyridylmethyl)-2-mercaptoaniline	778
Magnetic properties of gold(II) complex with <i>o</i>-(methylthio)aniline	779
Magnetic properties of dimeric gold(II) chloro complex with N-(2-pyridyl-methyl)-2-mercaptoaniline	781
Part IV Zn	783
Magnetic properties of zinc(II) complex with substituted diiminoquinone	785
Part V Cd	787
Magnetic properties of chloro-bridged tetranuclear cadmium(II) cluster with pyridyl-substituted nitronyl nitroxide	789
Part VI Ce	791
Magnetic properties of γ-picoline-N-oxide adduct of cerium(III) glycolate	793
Magnetic properties of cerium(III) complex with 2-(3-coumarinylthio-acetyl) benzimidazole	794

Magnetic properties of cerium(III) complex with 3-amino-2-hydroxy-1,4-naphthoquinone	795
Magnetic properties of cerium(III) phthalate as hydrazinium salt	796
Magnetic properties of anionic Ce^{3+} selenophosphate coordination complex	797
Part VII Pr	799
Magnetic properties of γ-picoline-N-oxide adduct of praseodymium(III) glycolate	801
Magnetic properties of praseodymium(III) complex with <i>p</i>-methoxy-phenylglyoxalthiosemicarbazone	802
Magnetic properties of praseodymium(III) with 8-acetyl-7-hydroxy-4-methylcoumarin	803
Magnetic properties of praseodymium(III) complex with 1-naphthylacetic acid	804
Magnetic properties of praseodymium(III) complex with 2-(3-coumarinyl-thioacetyl) benzimidazole	805
Magnetic properties of praseodymium(III) perchlorate complex with phenylalanine	806
Magnetic properties of praseodymium(III) nitrate complex with phenylalanine	807
Magnetic properties of praseodymium(III) complex with 3-amino-2-hydroxy-1,4-naphthoquinone	808
Magnetic properties of praseodymium(III) phthalate as hydrazinium salt	809
Magnetic properties of a mononuclear ten-coordinate praseodymium(III) complex with di-Schiff-base	810
Magnetic properties of copper praseodymium (CuPr_2) trichloroacetate	811
Magnetic properties of reduced praseodymium rhenate	813
Magnetic properties of chains of centered praseodymium cluster with some distortion	814
Magnetic properties of reduced praseodymium rhenate	816
Part VIII Nd	817
Magnetic properties of γ-picoline-N-oxide adduct of neodymium(III) glycolate	819
Magnetic properties of neodymium(III) complex with <i>p</i>-chlorophenyl-glyoxalthiosemicarbazone	820
Magnetic properties of neodymium(III) with 8-acetyl-7-hydroxy-4-methylcoumarin	821
Magnetic properties of neodymium(III) complex with 1-naphthylacetic acid	822
Magnetic properties of neodymium(III) complex with 2-(3-coumarinyl-thioacetyl) benzimidazole	823
Magnetic properties of neodymium(III) perchlorate complex with phenylalanine	824
Magnetic properties of neodymium(III) nitrate complex with phenylalanine	825
Magnetic properties of neodymium(III) complex with 22-membered tetraaza macrocycle	826
Magnetic properties of neodymium(III) complex with 18-membered tetraaza macrocycle	827
Magnetic properties of neodymium(III) complex with 14-membered tetraaza macrocycle	828

Magnetic properties of neodymium(III) complex with 3-amino-2-hydroxy-1, 4-naphthoquinone	829
Magnetic properties of neodymium(III) phthalate as hydrazinium salt	830
Magnetic properties of neodymium(III) complex with 2-formyl-4-methyl-6-{N-(2-pyridylethyl)formimidoyl}phenol ligand	831
Magnetic properties of polymeric mixed metal neodymium-copper complex with diglycolic acid	833
Magnetic properties of (μ-oxo)-heterometallic Nd-Cu complex with 2-hydroxypyridine	834
Magnetic properties of reduced neodymium rhenate	835
Part IX Sm	837
Magnetic properties of samarium(II) polymeric complex containing cyclopentadienylvanadiumnaphthalene anion	839
Magnetic properties of samarium(II) silyl substituted tellurolate as TMEDA derivative	840
Magnetic properties of samarium(II) silyl substituted selenolate as TMEDA derivative	841
Magnetic properties of γ-picoline-N-oxide adduct of samarium(III) glycolate	842
Magnetic properties of samarium(III) complex with <i>p</i>-methoxyphenyl-glyoxalthiosemicarbazone	843
Magnetic properties of samarium(III) with 8-acetyl-7-hydroxy-4-methylcoumarin	844
Magnetic properties of samarium(III) complex with 1-naphthylacetic acid	845
Magnetic properties of samarium(III) complex with 2-(3-coumarinylthioacetyl) benzimidazole	846
Magnetic properties of samarium(III) perchlorate complex with phenylalanine	847
Magnetic properties of samarium(III) nitrate complex with phenylalanine	848
Magnetic properties of samarium(III) complex with 3-iodolawsone-monoxime	849
Magnetic properties of samarium(III) complex with 3-bromolawsone-monoxime	850
Magnetic properties of samarium(III) complex with 3-chlorolawsone-monoxime	851
Magnetic properties of samarium(III) complex with 3-amino-2-hydroxy-1, 4-naphthoquinone	852
Magnetic properties of samarium(III) phthalate as hydrazinium salt	853
Magnetic properties of sandwich samarium(III) carborane cluster	854
Magnetic properties of samarium(III) complex with Schiff-base	856
Magnetic properties of dimeric samarium(II) complex with Schiff-base	857
Magnetic properties of dimeric oxo bridged samarium(II) complex with Schiff-base	858
Magnetic properties of (μ-chloro)bridged dinuclear samarium(III) supported by the Calix-tetrapyrrole ligand	859
Magnetic properties of (μ-methyl)bridged dinuclear samarium(III) supported by the Calix-tetrapyrrole ligand	860

Magnetic properties of (μ-chloro)bridged mixed valence samarium(II, III) complex supported by the Calix-tetrapyrrole ligand	861
Magnetic properties of dinuclear samarium(II) complex with Calix-tetrapyrrole ligand	862
Magnetic properties of mononuclear samarium(III) hydride with Calix-tetrapyrrole ligand	863
Magnetic properties of cuboid cluster of samarium(II) having Schiff-base	864
Magnetic properties of reduced samarium rhenate	865
Part X Eu	867
Magnetic properties of europium(II) complex containing cyclopentadienylvanadiumnaphthalene anion	869
Magnetic properties of europium(II) polymeric complex containing cyclopentadienylvanadiumnaphthalene anion	870
Magnetic properties of europium(II) silyl substituted selenolate as TMEDA derivative	871
Magnetic properties of europium(II) silyl substituted selenolate as DMPE derivative	872
Magnetic properties of europium(II) silyl substituted tellurolate as TMEDA derivative	873
Magnetic properties of europium(III) complex with <i>p</i>-methoxyphenylglyoxalthiosemicarbazone	874
Magnetic properties of europium(III) phthalate as hydrazinium salt	875
Magnetic properties of europium(III) aqua complex with phthalic acid as hydrazinium salt	876
Magnetic properties of europium(II) silyl substituted tellurolate as DMPE derivative	877
Magnetic properties of europium bismuthide	878
Magnetic properties of europium germanide	880
Part XI Gd	881
Magnetic properties of mixed oxide of strontium(II), gadolinium(III) and rhodium(III)	883
Magnetic properties of γ-picoline-N-oxide adduct of gadolinium(III) glycolate	885
Magnetic properties of gadolinium(III) with 8-acetyl-7-hydroxy-4-methylcoumarin	886
Magnetic properties of gadolinium(III) complex with 1-naphthylacetic acid	887
Magnetic properties of gadolinium(III) complex with 2-naphthylacetic acid	888
Magnetic properties of gadolinium(III) perchlorate complex with phenylalanine	889
Magnetic properties of gadolinium(III) nitrate complex with phenylalanine	890
Magnetic properties of gadolinium(III) complex with 3-amino-2-hydroxy-1,4-naphthoquinone	891
Magnetic properties of gadolinium(III) aqua complex with phthalic acid as hydrazinium salt	892
Magnetic properties of gadolinium(III) complex with <i>p</i>-chlorophenyl-glyoxalthiosemicarbazone	893
Magnetic properties of a mononuclear ten-coordinate gadolinium(III) complex with di-Schiff-base	894

Magnetic properties of gadolinium(III) complex with 2-(3-coumarinylthioacetyl) benzimidazole	895
Magnetic properties of gadolinium(III) complex with 2-formyl-4-methyl-6-{N-(2-pyridylethyl)formimidoyl}phenol ligand	896
Magnetic properties of gadolinium(III) complex with pyridine-substituted nitronyl nitroxide radical	897
Magnetic properties of gadolinium(III) complex with pyridine-substituted nitronyl nitroxide radical	899
Magnetic properties of mixed metallic gadolinium-copper telluride	901
Magnetic properties of sandwich gadolinium(III) carborane cluster	903
Magnetic properties of bimetallic Gd^{III}Cu^{II} molecular material having oxalate and 1, 3-propylene-bis(oxamate) ligands	904
Magnetic properties of diphenylbutadiene-bridged gadolinium complex	906
Magnetic properties of Mn-Gd pivalate	907
Part XII Tb	909
Magnetic properties of terbium(III) complex with <i>p</i>-chlorophenylglyoxalthiosemicarbazone	911
Magnetic properties of terbium(III) complex with 1-naphthylacetic acid	912
Magnetic properties of terbium(III) complex with 2-naphthylacetic acid	913
Magnetic properties of terbium(III) aqua complex with phthalic acid as hydrazinium salt	914
Magnetic properties of terbium rhenate	915
Part XIII Dy	917
Magnetic properties of dysprosium(III) complex with 1-naphthylacetic acid	919
Magnetic properties of dysprosium(III) aqua complex with phthalic acid as hydrazinium salt	920
Magnetic properties of dysprosium(III) complex with 2-naphthylacetic acid	921
Magnetic properties of dysprosium(III) perchlorate complex with phenylalanine	922
Magnetic properties of dysprosium(III) nitrate complex with phenylalanine	923
Magnetic properties of dysprosium(III) with 8-acetyl-7-hydroxy-4-methylcoumarin	924
Magnetic properties of dysprosium(III) complex with 3-iodolawsone-monoxime	925
Magnetic properties of dysprosium(III) complex with 3-bromolawsone-monoxime	926
Magnetic properties of dysprosium(III) complex with 3-chlorolawsone-monoxime	927
Magnetic properties of dysprosium(III) complex with 3-amino-2-hydroxy-1, 4-naphthoquinone	928
Magnetic properties of dysprosium(III) complex with 2-(3-coumarinyl-thioacetyl) benzimidazole	929
Magnetic properties of dysprosium(III) complex with <i>p</i>-methoxyphenyl-glyoxalthiosemicarbazone	930

Magnetic properties of dysprosium(III) complex with 22-membered tetraaza macrocycle	931
Magnetic properties of dysprosium(III) complex with 18-membered tetraaza macrocycle	932
Magnetic properties of dysprosium(III) complex with 14-membered tetraaza macrocycle	933
Magnetic properties of sandwich dysprosium(III) carborane cluster	934
Part XIV Ho	935
Magnetic properties of sandwich holmium(III) carborane cluster	937
Magnetic properties of holmium rhenate	938
Part XV Er	939
Magnetic properties of polymeric mixed metal erbium-copper complex with diglycolic acid	941
Part XVI Yb	943
Magnetic properties of ytterbium(III) complex with <i>p</i>-chlorophenyl-glyoxalthiosemicarbazone	945
Magnetic properties of ytterbium nickel carbide	946