

Contents

1	Introduction	1
	References	5
 Part I Synthesis of Indole Derivatives		
2	Construction of 2-(Aminomethyl)indoles through Copper-Catalyzed Domino Three-Component Coupling and Cyclization	9
2.1	Introduction	9
2.1.1	Synthesis of 2-(Aminomethyl)indoles Using Several Amines and Aldehydes	10
2.1.2	Synthesis of Substituted 2-(Aminomethyl)indoles Using Various Ethynylanilines and Secondary Amines	13
2.1.3	Construction of Polycyclic Indoles by Palladium-Catalyzed C–H Functionalization	14
2.1.4	Synthetic Application to Calindol, Benzo[e][1,2]thiazines, and Indene	14
2.2	Experimental Section	19
2.2.1	General Methods	19
2.2.2	General Procedure for Synthesis of 2-(Aminomethyl)indole	24
2.2.3	General Procedure for Synthesis of Tetrahydropyridine-Fused Indole	33
2.2.4	General Procedure for Synthesis of Benzo[e][1,2]thiazine-1,1-dioxide	41
	References	44
3	Facile Synthesis of 1,2,3,4-Tetrahydro-β-Carbolines by One-Pot Domino Three-Component Indole Formation and Nucleophilic Cyclization	47
3.1	Experimental Section	52
3.1.1	General Methods	52

3.1.2	General Procedure for Synthesis of <i>N</i> -Arylsulfonyl-2-ethynylaniline: Synthesis of 2-Ethynyl- <i>N</i> -mesitylenesulfonylaniline (1c)	53
3.1.3	<i>N</i> -(<i>p</i> -Bromobenzenesulfonyl)-2-ethynylaniline (1e)	53
3.1.4	<i>N</i> -(<i>p</i> -Chlorobenzenesulfonyl)-2-ethynylaniline (1f)	54
3.1.5	2-Ethynyl- <i>N</i> -(<i>p</i> -fluorosulfonyl)aniline (1g)	54
3.1.6	General Procedure for Synthesis of 1,2,3,4-Tetrahydro- β -carboline by Domino Copper-Catalyzed Three-Component Indole Formation and Cyclization with <i>t</i> -BuOK: Synthesis of 2-Methyl-1-propyl-2,3,4,9-tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]- indole (6a) and 2-Methyl-1-propyl-1,2,3,4- tetrahydropyrazino[1,2- <i>a</i>]indole (8a) (Table 1, Entry 11)	54
3.1.7	2-Methyl-1-[2-(trimethylsilyl)ethenyl]-2,3,4,9- tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indole (6b) and 2-Methyl-1-[2(trimethylsilyl)ethenyl]-1,2,3,4- tetrahydropyrazino[1,2- <i>a</i>]indole (8b)	55
3.1.8	1-(Benzyloxymethyl)-2-methyl-2,3,4,9-tetrahydro- 1 <i>H</i> -pyrido[3,4- <i>b</i>]indole (6c) and 1-(Benzyloxymethyl)- 2-methyl-1,2,3,4-tetrahydropyrazino[1,2- <i>a</i>]indole (8c) . .	56
3.1.9	2-Methyl-2,3,4,9-tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]- indole (6d)	56
3.1.10	General Procedure for Synthesis of 1,2,3,4-Tetrahydro- β -carboline by Domino Copper-Catalyzed Three-Component Indole Formation and Cyclization with MsOH: Synthesis of 2-Methyl-2,3-dihydropyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)-one (7a) (Conditions A)	57
3.1.11	2-Allyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)- one (7b)	57
3.1.12	2-Butyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)- one (7c)	58
3.1.13	General Procedure for Synthesis of 1,2,3,4-Tetrahydro- β -carboline by Domino Copper-Catalyzed Three-Component Indole Formation and Cyclization by MsOH: Synthesis of 2-Benzyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)-one (7d) (Conditions B)	58
3.1.14	(<i>R</i>)-2,3-Dimethyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indol- 4(9 <i>H</i>)-one (7e) (Conditions C)	59
3.1.15	(<i>R</i>)-3-Isobutyl-2-methyl-2,3-dihydro- 1 <i>H</i> -pyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)-one (7f)	59

3.1.16	(<i>R</i>)-3-Benzyl-2-methyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i>]indol-4(9 <i>H</i>)-one (7g)	59
3.1.17	5,6,8,9,10,11,11 <i>a</i> ,12-Octahydroindolo[3,2- <i>b</i>]quinolizine (7h)	60
References		60
4	Concise Synthesis of Indole-Fused 1,4-Diazepines through Copper(I)-Catalyzed Domino Three-Component Coupling–Cyclization–<i>N</i>-Arylation under Microwave Irradiation	63
4.1	Experimental Section	67
4.1.1	General Methods	67
4.1.2	General Procedure for Synthesis of 2-Ethynyl- <i>N</i> -methanesulfonylaniline: Synthesis of 2-Ethynyl- <i>N</i> -methanesulfonyl-4-methoxycarbonylaniline (1b)	68
4.1.3	2-Ethynyl- <i>N</i> -methanesulfonyl-4-trifluoromethyl-carbonylaniline (1c)	68
4.1.4	2-Ethynyl- <i>N</i> -methanesulfonyl-4-methylaniline (1d)	69
4.1.5	2-Ethynyl- <i>N</i> -methanesulfonyl-5-trifluoromethyl-carbonylaniline (1e)	69
4.1.6	General Procedure for Synthesis of Indole-Fused 1,4-Diazepine through Three-Component Indole Formation- <i>N</i> -Arylation: Synthesis of 7- <i>n</i> -Butyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>]-[1,4]diazepine (3a)	69
4.1.7	7-Methyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>]-[1,4]diazepine (3b)	70
4.1.8	7-Benzyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>]-[1,4]diazepine (3c)	70
4.1.9	7-Allyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>]-[1,4]diazepine (3d)	71
4.1.10	7-Allyl-3-methoxycarbonyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>][1,4]diazepine (3e)	71
4.1.11	7-Allyl-3-trifluoromethyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>][1,4]di-azepine (3f)	71
4.1.12	7-Allyl-3-methyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>][1,4]diazepine (3g)	72
4.1.13	7-Allyl-2-trifluoromethyl-7,8-dihydro-6 <i>H</i> -benzo[<i>f</i>]indolo[1,2- <i>a</i>][1,4]diazepine (3h)	72
4.1.14	Synthesis of <i>N</i> -[(2-bromothiophen-3-yl)methyl]-butan-1-amine (4)	73
4.1.15	7-Allyl-7,8-dihydro-6 <i>H</i> -pyrido[3,2- <i>f</i>]indolo[1,2- <i>a</i>][1,4]diazepine (5)	73
4.1.16	Synthesis of <i>N</i> -((2-bromopyridin-3-yl)methyl)prop-2-en-1-amine (6)	73

4.1.17	7-Allyl-7,8-dihydro-6 <i>H</i> -indolo[1,2- <i>a</i>]thieno-[2,3- <i>f</i>][1,4]diazepine (7)	74
References		74

Part II Synthesis of Isoquinoline Derivatives

5	Facile Synthesis of 3-(Aminomethyl)isoquinolines by Copper-Catalyzed Domino Four-Component Coupling and Cyclization	79
5.1	Introduction	79
5.2	Experimental Section	83
5.2.1	General Methods	83
5.2.2	General Procedure for Four-Component Isoquinoline Formation	85
References		88
6	Rapid Access to 3-(Aminomethyl)isoquinoline-Fused Polycyclic Compounds by Copper-Catalyzed Four-Component Coupling, Cascade Cyclization, and Oxidation	89
6.1	Experimental Section	94
6.1.1	General Procedure for Synthesis of (Aminomethyl)-isoquinoline-Fused Polycyclic Compounds by Domino Mannich-Type Reaction and Cascade Cyclization: Synthesis of 6-[(<i>N,N</i> -Diisopropylamino)methyl]-3,4-dihydro-2 <i>H</i> -pyrimido[2,1- <i>a</i>]isoquinoline (12a) (Table 1, Entry 10)	94
6.1.2	6-[(<i>N,N</i> -Diisopropylamino)methyl]-9-Fluoro-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12b)	95
6.1.3	6-[(<i>N,N</i> -Diisopropylamino)methyl]-10-Fluoro-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12c)	95
6.1.4	6-[(<i>N,N</i> -Diisopropylamino)methyl]-9-Methyl-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12d)	96
6.1.5	6-[(<i>N,N</i> -Diisopropylamino)methyl]-10-Methoxy-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12e)	96
6.1.6	6-(Piperidin-1-ylmethyl)-3,4-Dihydro-2 <i>H</i> -Pyrimido-[2,1- <i>a</i>]isoquinoline (12f)	96
6.1.7	6-[(<i>N,N</i> -Diallylamino)methyl]-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12g)	97
6.1.8	6-[[<i>N,N</i> -Bis((<i>R</i>)-1-phenylethyl)amino]methyl]-3,4-Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i>]isoquinoline (12h)	97
6.1.9	5-[(<i>N,N</i> -Diisopropylamino)methyl]-2,3-Dihydroimidazo-[2,1- <i>a</i>]isoquinoline (13)	98

6.1.10	7-[(<i>N,N</i> -Diisopropylamino)methyl]-2,3,4,5-Tetrahydro [1,3]diazepino[2,1- <i>a</i>]isoquinoline (14)	98
6.1.11	8-[(<i>N,N</i> -Diisopropylamino)methyl]-3,4,5,6- Tetrahydro-2 <i>H</i> -[1,3]diazocino[2,1- <i>a</i>]isoquinoline (15). . .	98
6.1.12	6-[(<i>N,N</i> -Diisopropylamino)methyl]benzimidazo- [2,1- <i>a</i>]isoquinoline (16)	99
References		99
7	Conclusions	101