## **Contents**

## Preface — VII

| A | C | kn | 0 | W | led | lg | m | en | ts |  | I) | ( |
|---|---|----|---|---|-----|----|---|----|----|--|----|---|
|---|---|----|---|---|-----|----|---|----|----|--|----|---|

| 1     | Topics of this book —— 1   |
|-------|--|
| 1.1   | Topic 1: Derivation of Carleman estimates and stability estimates for some ill-posed Cauchy problems, Chapter 2 —— 1 |
| 1.2   | Topic 2: Global uniqueness theorems for multidimensional CIPs on the   |
|       | basis of the BK method, Chapter 3 —— 1   |
| 1.3   | Topic 3: Carleman estimates for numerical methods for ill-posed Cauch  |
|       | problems for PDEs, Chapters 4, 5, and 12 —— 3  |
| 1.4   | Topic 4: The convexification globally convergent numerical concept for   |
|       | CIPs: A far reaching consequence of the BK method, Chapters 6–11—  |
| 2     | Carleman estimates and Hölder stability for ill-posed Cauchy   |
|       | problems 7   |
| 2.1   | What is the Carleman estimate —— 8   |
| 2.2   | Hölder stability for ill-posed Cauchy problems —— 9  |
| 2.3   | Carleman estimate for the parabolic operator —— 13   |
| 2.4   | Carleman estimate for the elliptic operator —— 20  |
| 2.5   | Carleman estimate for a hyperbolic operator —— 22  |
| 2.6   | Specifying Hölder stability estimates for ill-posed Cauchy   |
|       | problems —— 28   |
| 2.6.1 | The parabolic operator —— 28   |
| 2.6.2 | The elliptic operator —— 31  |
| 2.6.3 | A hyperbolic operator —— 32  |
| 2.7   | Lipschitz stability estimate for an ill-posed problem for a hyperbolic   |
|       | equation —— 34   |
| 2.7.1 | The pointwise case —— 35   |
| 2.7.2 | The integral inequality —— <b>40</b>   |
| 3     | Global uniqueness for coefficient inverse problems and Lipschitz stability   |
|       | for a hyperbolic CIP —— 41   |
| 3.1   | Estimating an integral —— 41   |
| 3.2   | Hyperbolic equation —— 42  |
| 3.3   | Hyperbolic equation when one of initial conditions equals zero —— 51   |
| 3.4   | Parabolic equations —— 55  |
| 3.4.1 | Case 1: Parabolic and hyperbolic equations —— 56   |



| 3.4.2 | Case 2: The data at $\{t = t_0 \in (0, T)\}$ as well as the lateral Cauchy data —— <b>59</b> |
|-------|--|
| 3.4.3 | Case 3: Final overdetermination —— 64  |
| 3.5   | A coefficient inverse problem for an elliptic equation —— 67                                 |
| 3.6   | Lipschitz stability estimate of a CIP for a hyperbolic equation —— 67                        |
| 4     | The quasi-reversibility numerical method for ill-posed Cauchy problems for                   |
|       | linear PDEs —— 77  |
| 4.1   | Introduction — 77  |
| 4.2   | The Quasi-Reversibility Method (QRM) —— 79   |
| 4.2.1 | The Carleman estimate —— 79  |
| 4.2.2 | The Quasi-Reversibility Method (QRM) —— 80   |
| 4.3   | Elliptic equation —— 84  |
| 4.4   | Parabolic equation with the lateral Cauchy data —— 87  |
| 4.5   | Hyperbolic equation with lateral Cauchy data —— 90   |
| 5     | Convexification for ill-posed Cauchy problems for quasi-linear PDEs —— 93                    |
| 5.1   | Introduction —— 93   |
| 5.2   | Some facts of the convex analysis —— 94  |
| 5.3   | The general scheme of the method —— 97   |
| 5.3.1 | The Cauchy problem —— 97   |
| 5.3.2 | The pointwise Carleman estimate —— 99  |
| 5.3.3 | Theorems —— 100  |
| 5.4   | Proof of Theorem 5.3.1 —— 103  |
| 5.5   | Proof of Theorem 5.3.2 —— 106  |
| 5.6   | Proof of Theorem 5.3.4 —— 108  |
| 5.7   | Proof of Theorem 5.3.5 —— 108  |
| 5.8   | Specifying equations —— 109  |
| 5.8.1 | Quasilinear elliptic equation —— 109   |
| 5.8.2 | Quasilinear parabolic equation —— 110  |
| 5.8.3 | Quasilinear hyperbolic equation —— 111   |
| 5.9   | Numerical study —— 112   |
| 5.9.1 | The forward problem —— 112   |
| 5.9.2 | The ill-posed Cauchy problem and noisy data —— 114   |
| 5.9.3 | Specifying the functional $J_{\lambda,\beta}$ —— 114   |
| 5.9.4 | Minimization of $J_{\lambda,\beta}(v,F)$ — 115   |
| 5.9.5 | Results —— 115   |
| 5.10  | Summary —— <b>119</b>  |
| 6     | A special orthonormal basis in $L_2(a,b)$ for the convexification for CIPs                   |
|       | without the initial conditions—restricted Dirichlet-to-Neumann map —— 121                    |
| 6.1   | Introduction —— 121  |

| 6.2   | A CIP with the restricted DN data —— 122                                    |
|-------|---|
| 6.2.1 | The Carleman estimate —— 122  |
| 6.2.2 | Statement of the problem — 123  |
| 6.2.3 | A special orthonormal basis in $L_2(0,1)$ —— 125                            |
| 6.3   | An ill-posed problem for a coupled system of quasilinear PDEs —— 126        |
| 6.4   | Convexification — 129   |
| 6.4.1 | Weighted Tikhonov-like functional —— 129                                    |
| 6.4.2 | Numerical scheme —— 134   |
| 6.5   | Two specific examples —— 134  |
| 6.5.1 | Parabolic equation —— 134   |
| 6.5.2 | Hyperbolic equation —— 136  |
| 7     | Convexification of electrical impedance tomography with restricted          |
|       | Dirichlet-to-Neumann map data —— 139  |
| 7.1   | Introduction —— 139   |
| 7.2   | EIT with restricted Dirichlet-to-Neumann (DtN) data —— 140                  |
| 7.2.1 | The mathematical model —— 140   |
| 7.2.2 | An equivalent problem —— 142  |
| 7.3   | Cauchy problem for a system of coupled quasilinear elliptic                 |
|       | equations —— 143  |
| 7.3.1 | The orthonormal basis of Section 6.2.3 —— 143                               |
| 7.3.2 | Cauchy problem for a system of coupled quasilinear elliptic                 |
|       | equations —— 143  |
| 7.3.3 | Two new Carleman estimates —— 145   |
| 7.3.4 | Hölder stability and uniqueness of the Cauchy problem (7.21),               |
|       | (7.23) —— <b>150</b>  |
| 7.4   | Convexification —— 151  |
| 7.5   | Theorems —— 152   |
| 7.5.1 | Formulations of theorems —— 152   |
| 7.5.2 | Proof of Theorem 7.5.1 —— 155   |
| 7.5.3 | Proof of Theorem 7.5.4 —— 158   |
| 7.5.4 | Proof of Theorem 7.5.5 —— 160   |
| 7.6   | Numerical studies —— 160  |
| 7.6.1 | Some details of the numerical implementation —— 160                         |
| 7.6.2 | A multilevel method of the minimization of functional (7.52) —— 162         |
| 7.6.3 | Numerical testing —— 162  |
| 8     | Convexification for a coefficient inverse problem for a hyperbolic equation |
|       | with a single location of the point source —— 171                           |
| 8.1   | Introduction —— 171   |
| 8.2   | Statement of the inverse problem —— 172                                     |
| 8.3   | A system of coupled quasi-linear elliptic equations —— 174                  |

| 8.3.1  | The function $w(\mathbf{x},t)$ —— 174                                      |
|--------|--|
| 8.3.2  | The system of coupled quasi-linear elliptic PDEs —— 176                    |
| 8.4    | Globally strictly convex Tikhonov-like functional —— 178                   |
| 8.4.1  | The functional —— 178  |
| 8.4.2  | Theorems —— 180  |
| 8.5    | Proofs —— <b>182</b>   |
| 8.5.1  | Proof of Theorem 8.4.1 —— 182  |
| 8.5.2  | Proof of Theorem 8.4.2 —— 185  |
| 8.5.3  | Proof of Theorem 8.4.6 —— 187  |
| 8.6    | Numerical studies —— 188   |
| 8.6.1  | Some details of the numerical implementation —— 189                        |
| 8.6.2  | A multilevel minimization method of the functional $J_{\lambda}(W)$ —— 191 |
| 8.6.3  | Numerical testing —— 191   |
| 9      | Convexification for an inverse parabolic problem —— 199                    |
| 9.1    | Introduction —— 199  |
| 9.2    | Statement of the coefficient inverse problem —— 199                        |
| 9.3    | Weighted globally strictly convex Tikhonov-like functional —— 201          |
| 9.3.1  | Nonlinear integral differential equation —— 201                            |
| 9.3.2  | The functional —— 202  |
| 9.4    | Theorems —— 204  |
| 9.5    | Proofs of Theorems 9.4.2 and 9.4.4 —— <b>210</b>                           |
| 9.5.1  | Proof of Theorem 9.4.2 —— <b>210</b>                                       |
| 9.5.2  | Proof of Theorem 9.4.4 213   |
| 9.6    | Proof of Theorem 9.4.5 —— <b>214</b>                                       |
| 9.7    | Proof of Theorem 9.4.7 —— <b>217</b>                                       |
| 9.8    | Numerical testing —— 218   |
| 9.9    | Proof of Theorem 9.4.1 —— 221  |
| 10     | Experimental data and convexification for the recovery of the dielectric   |
|        | constants of buried targets using the Helmholtz equation —— 229            |
| 10.1   | Introduction —— 229  |
| 10.2   | Statement of the coefficient inverse problem —— 231                        |
| 10.3   | An auxiliary system of coupled quasilinear elliptic equations —— 233       |
| 10.3.1 | An equation without the unknown coefficient —— 233                         |
| 10.3.2 | Truncated Fourier series —— 235  |
| 10.3.3 | Boundary data (10.24), (10.25) —— <b>237</b>                               |
| 10.3.4 | Lipschitz stability of the boundary value problem                          |
|        | (10.23)–(10.25) <b>—— 238</b>  |
| 10.4   | Weighted Tikhonov-like functional —— 240                                   |
| 10.5   | Analysis of the functional $J_{\lambda \nu}(V)$ — 241                      |

| 10.5.1 | Strict convexity on $\overline{B(M)}$ — 241                               |
|--------|---|
| 10.5.2 | The minimizer of $J_{\lambda,\gamma}(V)$ on $\overline{B(M)}$ —— 245      |
| 10.5.3 | The distance between the minimizer and the "ideal" solution —— 245        |
| 10.6   | The globally convergent gradient projection method —— 248                 |
| 10.7   | Work with experimental data —— 250  |
| 10.7.1 | Experimental setup —— <b>250</b>  |
| 10.7.2 | Buried targets to be imaged —— 251  |
| 10.7.3 | The necessity of data propagation —— 252                                  |
| 10.7.4 | Data propagation revisited —— 254   |
| 10.7.5 | Computational setup —— 257  |
| 10.7.6 | Reconstruction results —— <b>261</b>                                      |
| 11     | Travel time tomography with formally determined incomplete data in        |
|        | 3D — 263  |
| 11.1   | Introduction —— 263   |
| 11.2   | Statement of the problem —— 265   |
| 11.3   | A special orthonormal basis — 267   |
| 11.4   | Estimate of $\tau_z^2(\mathbf{x}, \alpha)$ from the below —— 267          |
| 11.5   | A boundary value problem for a system of nonlinear coupled                |
|        | integro-differential equations —— 269                                     |
| 11.5.1 | A nonlinear integro-differential equation —— 269                          |
| 11.5.2 | Boundary value problem for a system of coupled integro- differential      |
|        | equations —— 270  |
| 11.5.3 | The positivity of the function $(u_0 + w + g)(\mathbf{x}, \alpha)$ —— 272 |
| 11.5.4 | Applying the multidimensional analog of Taylor formula —— 273             |
| 11.6   | Problem (11.53), (11.54) in the semidiscrete form —— <b>276</b>           |
| 11.7   | Lipschitz stability and uniqueness —— 281                                 |
| 11.8   | Weighted globally strictly convex Tikhonov-like functional —— 282         |
| 11.8.1 | Estimating an integral —— 282   |
| 11.8.2 | The functional —— 283   |
| 11.9   | Proofs of Theorems 11.8.1 and 11.8.4 —— <b>286</b>                        |
| 11.9.1 | Proof of Theorem 11.8.1 —— 286  |
| 11.9.2 | Proof of Theorem 11.8.4 —— 288  |
| 12     | Numerical solution of the linearized travel time tomography problem with  |
|        | incomplete data —— 291  |
| 12.1   | Introduction —— 291   |
| 12.2   | The linearization —— 293  |
| 12.3   | A boundary value problem for a system of coupled PDEs of the first        |
|        | order —— <b>296</b>   |
| 12.4   | The QRM in partial finite differences —— 300                              |
| 12.5   | Numerical implementation —— 303   |

XVI — Contents

12.5.1 Numerical tests — 304

Bibliography --- 311

Index --- 323