

Contents

Introduction	1
Magnetic properties of di- μ -hydroxo-bis(bipyridyl)dicopper(II) sulphate pentahydrate	11
Magnetic properties of bis{copper(II)} complex of ascidiacyclamide	12
Magnetic properties of dafone chlorocuprate complex	14
Magnetic properties of a mixed valence copper (I, II) phosphate	16
Magnetic properties of copper(II) porphyrin π -cation radical	18
Magnetic properties of di- μ -hydroxo-bis(1, 10-phenanthroline)dicopper(II) sulphate pentahydrate	20
Magnetic properties of di- μ -hydroxo-bis(N, N, N', N'-tetramethylethylenediamine) dicopper(II) nitrate trihydrate	21
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and 1-aminoethane	22
Magnetic properties of polymeric, two dimensional μ , 2, 2'-bipyrimidine, μ -oxalato bridged and μ -chloro-bridged copper(II) complex	23
Magnetic properties of copper(II) complex of N, N'-bis(2-cyanoethyl)-5, 7-dioxo-1, 4, 8, 11-tetraazacyclotetradecane	25
Magnetic properties of π -cation radical of Cu(II) porphyrin	26
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and 1, 2-diaminoethane	28
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and 1, 3-diaminopropane	29
Magnetic properties of copper(II) methylphosphonate	30
Magnetic properties of copper(II) ethylphosphonate	32
Magnetic properties of copper(II) phenylphosphonate	34
Magnetic properties of α -copper(II) methylphosphonate	37
Magnetic properties of α -copper(II) ethylphosphonate	39
Magnetic properties of α -copper(II) phenylphosphonate	41
Magnetic properties of β -copper(II) methylphosphonate	43
Magnetic properties of trimetallic(Cu ₃) oxomethylphosphonate dihydrate	45
Magnetic properties of Bis[(μ -nitrate-O)bis(nitrato)bis(μ -pyridazine)(pyridazine) copper(II)]copper(II)	47
Magnetic properties of (nitrato)(nitrato-O, O')tris(pyridazine)copper(II)	49
Magnetic properties of bis(nitrato)tetrakis(pyridazine)copper(II)	50
Magnetic properties of (nitrato)tetrakis(pyridazine)copper(II)nitrate	51
Magnetic properties of copper oxide arsenate	52

Magnetic properties of anhydrous β -copper(II) formate	54
Magnetic properties of anhydrous α -copper(II) formate	56
Magnetic properties of linear trinuclear Cu(II) complex derived from acyclic hexadentate Schiff base ligand	57
Magnetic properties of dinuclear imidazole bridged copper(II) complex of singly condensed Schiff base	59
Magnetic properties of dinuclear imidazole bridged copper(II) complex of non-symmetrical Schiff base	61
Magnetic properties of dinuclear copper(II) acetate complex of macrocyclic ligand	62
Magnetic properties of tetranuclear copper(II) acetate complex of macrocyclic ligand	64
Magnetic properties of mononuclear copper(II) complex of macrocyclic ligand	66
Magnetic properties of mononuclear copper(II) complex of methylated macrocyclic ligand	68
Magnetic properties of oximate-bridged asymmetric dinuclear copper(II) complex with a Cu(II)-I bond	69
Magnetic properties of copper(II) hydrogenselenite	71
Magnetic properties of copper(II) hydrogenselenite aqua complex	74
Magnetic properties of copper(II) hydrogenselenite nitrate complex	77
Magnetic properties of copper(II) manganese(II) heterobimetallic hydrogenselenite	80
Magnetic properties of copper(II) cobalt(II) heterobimetallic hydrogenselenite	82
Magnetic properties of copper(II) nickel(II) heterobimetallic hydrogenselenite	84
Magnetic properties of copper(II) copper(II) homobimetallic hydrogenselenite	86
Magnetic properties of copper(II) zinc(II) heterobimetallic hydrogenselenite	88
Magnetic properties of binuclear copper(II) complex of a macrocyclic ligand derived from pyridine-N-oxide	90
Magnetic properties of aqua(L-aspartato)(2,2'-bipyridine)copper(II) trihydrate	92
Magnetic properties of diethylenetriamine-bis(tetracyano-quinodimethanido) copper(II) complex	93
Magnetic properties of binuclear copper(II) complex of binucleating ligand 2, 6-diformyl-4-methylphenoldi(benzoylhydrazone)	95
Magnetic properties of copper hexafluoroacetylacetonate complex with nitroxide-4-amido-2, 2-dimethyl-5, 5-dimethoxy-3-imidazoline-1-oxyl	97
Magnetic properties of copper(II) chloride complex with 1, 2, 4-triazole	99
Magnetic properties of copper(II) bromide complex with 1, 2, 4-triazole	101
Magnetic properties of copper(II) chloride complex with 1, 2, 4-triazole	103
Magnetic properties of copper(II) bromide complex with 1, 2, 4-triazole	105
Magnetic properties of copper(II) complex of 2-aminopyrimidine	107
Magnetic properties of binuclear copper(II) complex of macrocyclic aza-amido ligand	109

Magnetic properties of copper(II)-praseodymium(III) complex of 2, 6-bis[(dimethylamino) methyl]-4-methylphenolato ligand	111
Magnetic properties of tetranuclear copper(II) complex of 2, 6-bis[(dimethylamino) methyl]-4-methylphenol ligand	113
Magnetic properties of trinuclear pyridazine-bridged Cu ₃ (II) complex	115
Magnetic properties of trinuclear pyridazine-bridged Cu ₂ Ni complex	116
Magnetic properties of Cu(II) complex with pyridazinecarboxamide ligand	117
Magnetic properties of heterodinuclear iron(III)-Cu(II) complex with Schiff base and acetylacetone ligands	118
Magnetic properties of copper(II) squarate tetrahydrate	120
Magnetic properties of bis[μ-chloro-dichlorobis{2, 6-bis(4-amino-2-thiabutyl)} 4-methylphenolato-(N, N ¹ , S, S ¹ -μ-O)dicopper(II)-μ-Cl]	122
Magnetic properties of bis[2, 6-diacetylpyridine-bis(1'-phthalazinylylhydrazonato)copper(II)	124
Magnetic properties of copper(II) chloride adduct of undeprotonated 2, 6-diacetylpyridine-bis(1'-phthalazinylylhydrazone)	125
Magnetic properties of copper(II) chloride adduct of monodeprotonated 2, 6-diacetylpyridine-bis(1'-phthalazinylylhydrazone)	127
Magnetic properties of copper(II) chloride adduct of undeprotonated 2, 6-diacetylpyridine-bis(1'-phthalazinylylhydrazone) water solvate	129
Magnetic properties of copper(II) chlorate adduct of monoprotated 2, 6-diacetylpyridine-bis(1'-phthalazinylylhydrazone) water solvate	131
Magnetic properties of tetrakis{[3, 5-bis(pyridine-2-yl)-pyrazolate] (aqua) copper(II) nitrate monohydrate} complex	133
Magnetic properties of {[3, 5-bis(pyridine-2-yl)-pyrazolate](aqua) copper(II)] chloride}	136
Magnetic properties of {[3, 5-bis(pyridine-2-yl)-pyrazolate](aqua) copper(II)] bromide}	138
Magnetic properties of copper(II) complex of Schiff-base derived from amino acid and ethyl-α-ketocyclopentylcarboxylate	140
Magnetic properties of copper(II) complex of Schiff-base derived from amino acid and ethyl-α-ketocyclopentylcarboxylate	141
Magnetic properties of copper(II) complex of Schiff-base derived from amino acid and ethyl-α-ketocyclopentylcarboxylate	142
Magnetic properties of copper(II) dipropyleneetriamine thiocyanate nitrate complex	144
Magnetic properties of binuclear copper(II) complex of pyridoxal benzoyl hydrazone	146
Magnetic properties of binuclear copper(II) complexes of pyridoxal <i>o</i> -hydroxy hydrazones	149
Magnetic properties of binuclear copper(II) complexes of pyridoxal picolinoyl hydrazone	151
Magnetic properties of binuclear copper(II) complexes of pyridoxal nicotinoyl hydrazone	153
Magnetic properties of binuclear copper(II) complexes of pyridoxal isonicotinoyl hydrazone	155
Magnetic properties of dimeric copper(II) acetate complex of 2-formyl pyridinethiosemicarbazone	157

Magnetic properties of tetranuclear ternary pyrophosphato-copper(II)-2-formyl pyridine thiosemicarbazone complex	159
Magnetic properties of rare earth-copper molecular materials	161
Magnetic properties of rare earth-copper molecular materials	163
Magnetic properties of copper complex with di-pyridylmethane	165
Magnetic properties of diaqua-bis(N-acetyl- <i>D, L</i> -phenylglycinato)-bis(imidazole)copper(II)	167
Magnetic properties of <i>cis</i> -(9, 10-dihydro-9-oxo-10-acridineacetato- μ -O) copper(II)	168
Magnetic properties of copper(II) complex of Schiff base derived from 2-[(2-aminoethyl)thiomethyl]-benzimidazole and 2, 6-diacetylpyridine	170
Magnetic properties of imidazolate bridged dinuclear Cu(II)-Fe(III) octaethylporphyrin	172
Magnetic properties of imidazolate bridged dinuclear Cu(II)-Fe(III) porphyrin	174
Magnetic properties of copper(II) chelate containing symmetric N, N'-dimethyl-1, 2-diaminoethane and acetylacetone, with chlorate as anion	176
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and acetylacetone, with chlorate as counter anion	177
Magnetic properties of copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and 3-cyano-2, 4-pentanedione with hexafluorophosphate as anion	178
Magnetic properties of copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and water with tetrafluoroborate anion	179
Magnetic properties of copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and water with perchlorate anion	180
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and 3-cyano-2, 4-pentanedione with azide anion	181
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and o-nitrite	182
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and thiocyanate	183
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and isocyanide	184
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and fluoride	185
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and chloride	186
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane, 3-cyano-2, 4-pentanedione and bromide	187
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and 3-cyano-2, 4-pentanedione	188
Magnetic properties of mixed ligand copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and 3-cyano-2, 4-pentanedione with perchlorate anion	189
Magnetic properties of copper(II) chelate containing N, N-dimethyl-1, 2-diaminoethane and 3-cyano-2, 4-pentanedione with nitrate counterion	190

Magnetic properties of mixed ligand copper(II) chelate containing bis(N, N-dimethyl-1, 2-diaminoethane) and bis(3-cyano-2, 4-pentanedione)	191
Magnetic properties of mixed ligand copper(II) chelate containing N, N, N', N'-tetramethyl-1, 2-diaminoethane and bis(3-cyano-2, 4-pentanedione)	193
Magnetic properties of mixed bridged, dinuclear copper(II) chloride complex with dinucleating, 3, 5-bis(pyridin-2-yl)pyrazole	195
Magnetic properties of mixed bridged, dinuclear copper(II) bromide complex with dinucleating 3, 5-bis(pyridin-2-yl)pyrazole	197
Magnetic properties of mixed bridged, dinuclear copper(II) chloride complex with 3-(6-methylpyridin-2-yl)-5-(pyridin-2-yl) pyrazole	199
Magnetic properties of mixed bridged, dinuclear copper(II) bromide complex with 3-(6-methylpyridin-2-yl)-5-(pyridin-2-yl)pyrazole	201
Magnetic properties of mixed bridged, dinuclear copper(II) chloride complex with dinucleating 3, 5-bis(6-methylpyridine-2-yl)pyrazole	203
Magnetic properties of dinuclear bis(μ -3-pyridin-2-yl-1, 2, 4-triazolato) copper(II) compound containing N-methylimidazole as CO-ligand	205
Magnetic properties of dinuclear bis(μ -3-pyridin-2yl-1, 2, 4-triazolato) copper(II) compound containing pyrazole as CO-ligand	207
Magnetic properties of polynuclear bis(μ -3-pyridin-2-yl-1, 2, 4-triazolato) copper(II) compound containing 4, 4'-bipyridine as CO-ligands	208
Magnetic properties of copper(II) croconate trihydrate	209
Magnetic properties of dinuclear (Cu/Cu) complex of a novel oxamide-type ligand coordinating in the <i>trans</i> conformation	211
Magnetic properties of phenolato bridged, heterodinuclear Cu ^{II} Mn ^{II} dimer	213
Magnetic properties of oximato bridged, heterodinuclear Cu ^{II} Mn ^{II} dimer	215
Magnetic properties of oxamidato bridged, heterodinuclear Cu ^{II} Mn ^{II} dimer	217
Magnetic properties of dinuclear hydroxo bridged copper(II) chloride complex of tetradentate (N4) thio-diazine ligand	219
Magnetic properties of dinuclear hydroxo bridged copper(II) bromide complex of tetradentate (N4) thio-diazine ligand	221
Magnetic properties of dinuclear hydroxo-bridged copper(II) nitrate complex of tetradentate (N4) thio-diazine ligand	223
Magnetic properties of binuclear fumarato-bridged bis[(N-(2-diethylamino)ethyl)-salicylideneaminato]copper(II) complex	225
Magnetic properties of bis(dimethyl sulfoxide) adduct of tetra- μ -formato-dicopper(II); tetracarboxylato-bridged copper(II) dimer	227
Magnetic properties of a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II)	229
Magnetic properties of a pyrazine-bridged dinuclear copper(II) complex	231
Magnetic properties of a hydrogen bonded polynuclear copper(II) complex	233
Magnetic properties of <i>syn-anti</i> carboxylate-bridged dinuclear copper(II) complex	235

Magnetic properties of μ -phthalato-bis(piperidine)Cu(II) chain complex	237
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and 1-aminopropane	238
Magnetic properties of oximate bridged Cu ^{II} Cu ^{II} dimeric complex containing bipyridine and aqua ligands	239
Magnetic properties of oximate-bridged Cu ^{II} Cu ^{II} dimeric complex containing terpyridine	241
Magnetic properties of oximate-bridged Cu ^{II} Ni ^{II} dimeric complex containing cyclam	243
Magnetic properties of copper(II) complex of 2-nitroso-4-methylaniline	245
Magnetic properties of copper(II) complex of 2-nitroso-4, 6-dimethylaniline	246
Magnetic properties of a 2D layered copper(II) complex with bridging <i>trans</i> -oxamate and pyrimidine ligands	247
Magnetic properties of dinuclear copper(II) complex of thioether-pyridazine macrocycle	249
Magnetic properties of dinuclear copper(II) complex of thioether-pyridazine macrocycle	251
Magnetic properties of binuclear copper(II) complex of substituted methylpyrazole	253
Magnetic properties of polymeric 3, 5-dimethylpyrazolatecopper(II)	255
Magnetic properties of polymeric 4-chloro-3, 5-dimethylpyrazolate copper(II)	257
Magnetic properties of polymeric 4-bromo-3, 5-dimethylpyrazolate copper(II)	259
Magnetic properties of polymeric 3, 4, 5-trimethylpyrazolate copper(II)	261
Magnetic properties of dinuclear copper(II, II) complex of thioether-pyridazine (hexathiapyridazinophane) macrocycle ligand	263
Magnetic properties of dinuclear copper(II, I) complex of thioether-pyridazine macrocycle ligand	265
Magnetic properties of dinuclear copper(II) complex of thioether-pyridazine macrocyclic ligand	266
Magnetic properties of copper(II) complex of thioether-pyridazine macrocyclic ligand	268
Magnetic properties of copper(II) complex of Schiff-base derived from S-methyldithiocarbazate and glyoxal	270
Magnetic properties of copper(II) complex of Schiff-base derived from S-benzylthiocarbazate and glyoxal	271
Magnetic properties of copper(II) dichloride complex of 2-acetylpyridine-3-(4-methyl) piperazinyl-thiosemicarbazone	272
Magnetic properties of copper(II) dibromide complex of 2-acetylpyridine-3-(4-methyl) piperazinyl-thiosemicarbazone	273
Magnetic properties of copper(II) chloride complex of 2-acetylpyridine-3-(4-methyl) piperazinyl-thiosemicarbazone	274
Magnetic properties of copper(II) bromide complex of 2-acetylpyridine-3-(4-methyl) piperazinyl-thiosemicarbazone	275
Magnetic properties of copper(II) complex of indol-3- β -acrylic acid	276
Magnetic properties of copper(II) acetate complex with piperazine	277

Magnetic properties of copper(II) acetate complex with 1-methyl piperazine	279
Magnetic properties of copper(II) acetate complex with bis(piperazinemonohydrochloride)	281
Magnetic properties of copper(II) acetate complex with half mole(0.5) piperazine	283
Magnetic properties of copper(II) acetate complex with piperazinemonohydrochloride	285
Magnetic properties of copper(II) acetate complex of 1-methylpiperazine	287
Magnetic properties of copper(II) acetate complex with half-mole(0.5)-1-methylpiperazine	289
Magnetic properties of copper(II) acetate complex with 1-methylpiperazinemonohydrochloride . . .	291
Magnetic properties of copper(II) acetate complex with half-mole(0.5)-1, 4-dimethylpiperazine . . .	293
Magnetic properties of copper(II) acetate complex with 1, 4-dimethylpiperazinemonohydrochloride	295
Magnetic properties of copper(II) chloride complex of potassium dihydrobis(5-nitroindolyl)borate	297
Magnetic properties of copper(II) chloride complex of potassium hydrotris(5-nitroindolyl)borate	298
Magnetic properties of copper(II) chloride complex of tetrakis(5-nitroindolyl)borate	299
Magnetic properties of benzoate copper(II) dioxygen complex	300
Magnetic properties of 2-furylcarboxylatocopper(II) dioxygen complex	301
Magnetic properties of 2-pyridylcarboxylatocopper(II) dioxygen complex	302
Magnetic properties of heterobinuclear copper(II)-Nickel(II) complex with N, N'-bis(2-aminopropyl)oxamide and 2, 2'-bipyridine	303
Magnetic properties of heterobinuclear copper(II)-Nickel(II) complex with N, N'-bis(2-aminopropyl)oxamide and 1, 10-phenanthroline	305
Magnetic properties of heterobinuclear copper(II)-nickel(II) complex with N, N'-bis(2-aminopropyl)oxamide and 5-nitro-1, 10-phenanthroline	307
Magnetic properties of imidazolate-bridged copper(II) with 6-amino-1, 3-dimethyl- 5(2-carboxyphenyl)azouracil	308
Magnetic properties of imidazole-bridged dinuclear copper(II) with 1, 3-dimethyl- 5(2-carboxyphenyl)azobarbituric acid	310
Magnetic properties of dimeric copper(II) complex of fluorenone thiosemicarbazone	312
Magnetic properties of copper(II) complex with fluorenone thiosemicarbazone	313
Magnetic properties of copper(II) <i>O</i> -nitrate complex with fluorenone thiosemicarbazone	314
Magnetic properties of copper(II) sulphate complex with fluorenone thiosemicarbazone	315
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and 3-amino-1-propanol	316
Magnetic properties of copper(II) chelate of 1-phenyl-4-methylpent-4-en-1, 3-dione	317
Magnetic properties of copper(II) complex of 5-(<i>p</i> -tolyl)-2-phenyl-6H-pyrazolo- [1, 5- <i>c</i>]pyrimidine-7-thione	318
Magnetic properties of copper(II) complex of 5-(<i>p</i> -methoxyphenyl)-2-phenyl-6H- pyrazolo-[1, 5- <i>c</i>]pyrimidine-7-thiones	319

Magnetic properties of copper(II) complex of 5, 2-diphenyl-6H-pyrazolo-[1, 5- <i>c</i>]pyrimidine-7-thiones	320
Magnetic properties of copper(II) complex of 5- <i>p</i> -chlorophenyl-2-phenyl-6H-pyrazolo-[1, 5- <i>c</i>]pyrimidine-7-thiones	321
Magnetic properties of copper(II) complex of 5- <i>p</i> -bromophenyl-2-phenyl-6H-pyrazolo-[1, 5- <i>c</i>]pyrimidine-7-thiones	322
Magnetic properties of copper(II) complex of 2, 9-di(<i>o</i> -tolyl)-1, 10-phenanthroline	323
Magnetic properties of copper(II) complex of 2, 9-di(2-methoxyphenyl)-1, 10-phenanthroline	324
Magnetic properties of copper(II) complex of 2, 9-di(2-ethoxyphenyl)-1, 10-phenanthroline	325
Magnetic properties of copper(II) complex of 2-(<i>o</i> -tolyl)-1, 10-phenanthroline	326
Magnetic properties of copper(II) complex of 2-(2-ethoxyphenyl)-1, 10-phenanthroline	327
Magnetic properties of copper(II) chelate of N-(2-pyridyl) methacrylamide	328
Magnetic properties of copper(II) chloride complex of 2-(1-indazolyl)-pyridine	329
Magnetic properties of copper(II) chloride complex of 2-(1-indazolyl)-pyrimidine	330
Magnetic properties of dichlorotetrakis(3, 4-dicyano-5-aminopyrazole)-copper(II)	331
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 2, 6-diformyl-4-methylphenol and aniline	332
Magnetic properties of 2-(3, 5-dimethyl-pyrazole-1-yl)benzothiazole complex of copper(II) chloride	333
Magnetic properties of 2-(3, 5-dimethyl-pyrazole-1-yl)benzothiazole complex of copper(II) bromide	334
Magnetic properties of 2-(3, 5-dimethyl-pyrazole-1-yl)benzothiazole complex of copper(II) nitrate	335
Magnetic properties of 2-(3, 5-dimethyl-pyrazole-1-yl)benzothiazole complex of copper(II) sulphate	336
Magnetic properties of 2-(3, 5-dimethyl-pyrazole-1-yl)benzothiazole complex of copper(II) perchlorate	337
Magnetic properties of acetylacetonato derivative of mono(dizirconium-ennea-isopropoxy)copper(II)	338
Magnetic properties of hexafluoroacetylacetonato derivative of mono(dizirconium-ennea-isopropoxy)copper(II)	339
Magnetic properties of trifluoroacetylacetonato derivative of mono(dizirconium-ennea-isopropoxy)copper(II)	340
Magnetic properties of dimeric μ -(2, 6-dimethylphenolato)-mono(dizirconium-ennea-isopropoxy)copper(II)	341
Magnetic properties of dimeric μ -(3, 5-dimethylphenolato)-mono(dizirconium-ennea-isopropoxy)copper(II)	342
Magnetic properties of dimeric μ -(<i>p</i> -chlorophenolato)-mono(dizirconium-ennea-isopropoxy)copper(II)	343
Magnetic properties of polymeric μ -(chloro)-mono(dizirconium-ennea-methoxy)copper(II)	344

Magnetic properties of dimeric μ -(chloro)-mono(dizirconium-ennea-ethoxy)copper(II)	345
Magnetic properties of dimeric μ -(chloro)-mono(dizirconium-ennea-n-butoxy) copper(II)	346
Magnetic properties of dimeric μ -(chloro)-mono(dizirconium-ennea-s-butoxy) copper(II)	347
Magnetic properties of dimeric μ -(chloro)-mono(dizirconium-tris-isopropoxy-hexa- <i>t</i> -butoxy) copper(II)	348
Magnetic properties of acetylpyrazine-3-azabicyclo[3.2.2]nonylthiosemicarbazone complex of copper(II) bromide	349
Magnetic properties of acetylpyrazine-3-azabicyclo[3.2.2]nonylthiosemicarbazone complex of copper(II) chloride	350
Magnetic properties of copper(II) chloride complex of 2-acetylpyridine ⁴ N-(2-methylpyridinyl)thiosemicarbazone	351
Magnetic properties of copper(II) bromide complex of 2-acetylpyridine ⁴ N-(2-methylpyridinyl)thiosemicarbazone	352
Magnetic properties of copper(II) chloride complex of 2-acetylpyridine ⁴ N-(2-ethylpyridinyl)thiosemicarbazone	353
Magnetic properties of copper(II) bromide complex of 2-acetylpyridine ⁴ N-(2-ethylpyridinyl)thiosemicarbazone	354
Magnetic properties of copper(II) chloride complex of 2-acetylpyridine ⁴ N-methyl(2-ethylpyridinyl)thiosemicarbazone	355
Magnetic properties of copper(II) bromide complex of 2-acetylpyridine ⁴ N-methyl(2-ethylpyridinyl)thiosemicarbazone	356
Magnetic properties of copper(II) complex of N-salicylidene anthranilamide	357
Magnetic properties of 6-nitrobenzimidazolate bridged copper(II) complex	358
Magnetic properties of 6-nitrobenzimidazolate bridged copper(II) complex	360
Magnetic properties of 6-nitrobenzimidazolate bridged copper(II) complex	362
Magnetic properties of imidazolate bridged copper(II) complex	363
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 8-aminoquinoline and salicylaldehyde	364
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 8-aminoquinoline and 2-hydroxy-1-naphthaldehyde	365
Magnetic properties of copper(II) complex of Schiff base derived from 8-aminoquinoline and 2, 4-dihydroxybenzaldehyde	367
Magnetic properties of copper(II) complex of Schiff base derived from 8-aminoquinoline and 2, 3-dihydroxybenzaldehyde	368
Magnetic properties of dinuclear copper(II) complex of Schiff base derived from 8-aminoquinoline and 5-chlorosalicylaldehyde	369
Magnetic properties of allopurinol complex with copper(II) perchlorate	370
Magnetic properties of complex of copper(II) with tridentate Schiff base derived by the condensation of salicylaldehyde and furfuraldehyde with <i>o</i> -phenyldiamine	371
Magnetic properties of copper(II) chelate of acetoacetanilide thiosemicarbazone	372

Magnetic properties of copper(II) complex with acetylacetone bis(benzoyl-hydrazone)	373
Magnetic properties of copper(II) complex with acetylacetone bis-(isonicotinoyl hydrazone)	374
Magnetic properties of complex of copper(II) with Schiff base derived from 2-aminocyclopent-1-ene-1-dithiocarboxylic acid and benzaldehyde	375
Magnetic properties of complex of copper(II) chloride with Schiff base derived from 2-aminocyclopent-1-ene-1-dithiocarboxylic acid and salicylaldehyde	376
Magnetic properties of complex of copper(II) bromide with Schiff-base derived from 2-aminocyclopent-1-ene-dithiocarboxylic acid and salicylaldehyde	377
Magnetic properties of copper(II) chloride complex of 2, 6-diacetyl-pyridinesalicylaldazine	378
Magnetic properties of copper(II) complex of 2, 6-diacetylpyridine salicylaldazine	379
Magnetic properties of copper(II) complex of 1, 3, 8, 10, 15, 17, 22, 24-octanitrophthalocyanine	380
Magnetic properties of copper(II) chloride complex of 1, 3-bis(aminomethyl)cyclohexane- N, N, N', N'-tetrakis-benzimidazole	382
Magnetic properties of copper(II) perchlorate complex of 1, 3-bis-aminomethylcyclohexane- N, N, N', N'-tetrakis-benzimidazolato-copper(II) chloride	383
Magnetic properties of copper(II) chloride complex of pyridine-2-carboxaldehyde Schiff-base of S-methyl- β -N-(2-aminophenyl) methylenedithiocarbazate	384
Magnetic properties of copper(II) chloride complex of 6-methyl-pyridine-2-carboxaldehyde Schiff-base of S-methyl- β -N-(2-aminophenyl) methylenedithiocarbazate	385
Magnetic properties of copper(II) nitrate complex of pyridine-2-carboxaldehyde Schiff-base of S-methyl- β -N-(2-aminophenyl) methylenedithiocarbazate	386
Magnetic properties of copper(II) bromide complex of pyridine-2-carboxaldehyde Schiff-base of S-methyl- β -N-(2-aminophenyl) methylenedithiocarbazate	387
Magnetic properties of copper(II) complex with 2-pyridinone	388
Magnetic properties of copper(II) complex with 1-hydroxy-2-pyridine-thione	389
Magnetic properties of copper(II) complex with 2-pyridinemethane thiole-1-oxide	390
Magnetic properties of copper(II) mixed ligand complex with 2-pyridinone and phenanthroline . . .	391
Magnetic properties of benzoylacetoneisonicotinoylhydrazone complex of copper(II)	392
Magnetic properties of copper(II) complex derived from N-isonicotinamido-N'- benzoylthiocarbamide	394
Magnetic properties of copper(II) complex derived from N-isonicotinamido-N'- benzoylthiocarbamide	395
Magnetic properties of complex of copper(II) chloride with tetradentate Schiff-base ligand 1, 2-bis(2'-pyridyl)methyleneimino)benzene	396
Magnetic properties of copper(II) complex of 2-benzylthio-4-formyl-1- <i>p</i> -methoxyphenylimidazole	397
Magnetic properties of 1, 3-pyrimidinyl(2, 4, 6-pyrimidinetrione) chelate of copper(II)	399
Magnetic properties of <i>o</i> -vanillin-isonicotinoyl hydrazone complex of copper(II)	400

Magnetic properties of copper(II) complex of N, N'-bis(1-ferrocenyl-1-oxo-3-methyl) propenylethylenediamine	401
Magnetic properties of dinuclear copper(II) acetate complex with ligand derived from the condensation of triazolo-2, 4-pentanedione with 1, 6-diaminohexane	402
Magnetic properties of dinuclear copper(II) chloride complex with ligand derived from the condensation of triazolo-2, 4-pentanedione with 1, 6-diaminohexane	403
Magnetic properties of dinuclear copper(II) acetate complex with ligand derived from the condensation of thiazolo-2, 4-pentanedione with 1, 6-diaminohexane	404
Magnetic properties of dinuclear copper(II) chloride complex with ligand derived from the condensation of thiazolo-2, 4-pentanedione with 1, 6-diaminohexane	405
Magnetic properties of praseodymium(III) complex of an 18-membered dioxatetraaza macrocycle ligand	406
Magnetic properties of mixed ligand complex of praseodymium(III) with bis(phenylsulphinyl) propane and bipyridyl	407
Magnetic properties of neodymium(III) complex of 18-membered dioxatetraaza macrocycle ligand	409
Magnetic properties of mixed ligand complex of neodymium(III) with bis(phenylsulphinyl) propane and bipyridyl	410
Magnetic properties of samarium(III) complex of 18-membered dioxatetraaza macrocycle ligand	412
Magnetic properties of europium(III) complex of 18-membered dioxatetraaza macrocycle ligand	413
Magnetic properties of binuclear complex of europium(II) with the bridging naphthalene dianion	414
Magnetic properties of binuclear complex of europium(II) with the bridging naphthalene dianion	415
Magnetic properties of mixed ligand complex of europium(III) with bis(phenylsulphinyl) propane and bipyridyl	416
Magnetic properties of gadolinium(III)-copper(II) squarate	418
Magnetic properties of mixed ligand complex of gadolinium(III) with bis(phenylsulphinyl) propane and bipyridyl	419
Magnetic properties of heterodinuclear nickel(II)-gadolinium(III)-complex with an acyclic Schiff base	421
Magnetic properties of heterodinuclear copper(II)-gadolinium(III)-complex with an acyclic Schiff base	423
Magnetic properties of heterotetranuclear copper(II)-gadolinium(III)-complex with an acyclic Schiff base	425
Magnetic properties of mixed ligand complex of terbium(III) with bis(phenylsulphinyl)propane and bipyridyl	427
Magnetic properties of mixed ligand complex of holmium(III) with bis(phenylsulphinyl)propane and bipyridyl	429

Magnetic properties of mixed ligand complex of erbium(III) with bis(phenylsulphinyl)propane and bipyridyl	431
Magnetic properties of tris diazabutadiene ytterbium complex	433
Magnetic properties of mixed ligand complex of ytterbium(III) with bis(phenylsulphinyl)propane and bipyridyl	435
Magnetic properties of polymeric, two-dimensional, multidecker complex of bis(cyclopentadienyl)vanadium with naphthaleneytterbium	437
Magnetic properties of uranium triiodide tetrakis(acetonitrile)	439
Magnetic properties of uranyl(VI) complex derived from N-isonicotinamido-N'-benzoylthiocarbamide	440
Magnetic properties of dichloro(phthalocyano)silicon	441
Magnetic properties of bis(<i>p</i> -nitrophenoxy)(phthalocyanino)silicon	443
Magnetic properties of trimetallic lanthanum(III)-europium(III), La ₂ -Eu complex of 12-membered tetraaza macrocycle	445
Magnetic properties of trimetallic lanthanum(III)-gadolinium(III), La ₂ -Gd complex of 12-membered tetraaza macrocycle	446
Magnetic properties of trimetallic lanthanum(III)-terbium(III), La ₂ -Tb complex of 12-membered tetraaza macrocycle	447
Magnetic properties of trimetallic lanthanum(III)-dysprosium(III), La ₂ -Dy complex of 12-membered tetraaza macrocycle	448
Magnetic properties of heterodinuclear titanium lithium chloro complex supported by amide ligand	449
Magnetic properties of heterodinuclear titanium lithium methyl complex supported by amide ligand	450
Magnetic properties of dinuclear chlorotitanium(III) derivative with bridging μ -(dimethylsilyl) biscyclopentadienyl ligand	451
Magnetic properties of <i>trans</i> -bis(4-picoline)tetratolylporphyrinato)- titanium(II)	453
Magnetic properties of μ -(dichloro)bis[(trimethylsilyl)benzamidinato]-N, N, N', N'-tetramethylethylenediaminetitanium(III)-lithium(I)	454
Magnetic properties of bis[(trimethylsilyl)benzamidinato]titanium(III) borohydride	456
Magnetic properties of bis[(trimethylsilyl)benzamidinato]titanium(III)- η^3 -allyl complex	458
Magnetic properties of trisaminopyridinatotitanium(III) complex	459
Magnetic properties of dimeric titanium(III) complex with chloride and tetrahydrofuran	461
Magnetic properties of oxysulfide of mixed metals (Nd ₂ Ti ₂)	462
Magnetic properties of complex obtained by intercalation of η -cycloheptatrienyl molybdenum into ZrS ₂	464
Magnetic properties of complex obtained by intercalation of η -cycloheptatrienyl tungsten into ZrS ₂	466
Magnetic properties of hafnium(III) nitride	468
Magnetic properties of carbonylbis(pentadienyl)vanadium(II)	469

Magnetic properties of (triethylphosphine)bis(pentadienyl)vanadium(II)	470
Magnetic properties of (trimethyl phosphite)bis(pentadienyl) vanadium(II)	471
Magnetic properties of (trimethylphosphine)bis(2, 4-dimethylpentadienyl) vanadium(II)	472
Magnetic properties of (trimethyl phosphite)bis(2, 4-dimethylpentadienyl) vanadium(II)	473
Magnetic properties of triethylphosphine adduct of half-open, cyclopentadienyl (3-methylpentadienyl)vanadocene	474
Magnetic properties of trimethylphosphine adduct of half-open, cyclopentadienyl(2, 4- dimethylpentadienyl)vanadocene	475
Magnetic properties of triphenylphosphine adduct of half-open, cyclopentadienyl(2, 4- dimethylpentadienyl)vanadocene	476
Magnetic properties of trifluorophosphine adduct of half-open, cyclopentadienyl(2, 4- dimethylpentadienyl)vanadocene	477
Magnetic properties of carbonyl adduct of half-open, cyclopentadienyl (3-methylpentadienyl) vanadocene	478
Magnetic properties of trimethyl phosphite adduct of half-open, cyclopentadienyl(pentadienyl) vanadocenes	479
Magnetic properties of trimethyl phosphite adduct of half-open, cyclopentadienyl (3-methylpentadienyl)vanadocene	480
Magnetic properties of trimethylphosphite adduct of half-open, cyclopentadienyl(2, 4- dimethylpentadienyl)vanadocene	481
Magnetic properties of 4-methyl-2, 6, 7-trioxa-1-phosphabicyclo [2.2.2]octane adduct of half-open, cyclopentadienyl(2, 4-dimethyl pentadienyl)vanadocene	482
Magnetic properties of triphenyl phosphite adduct of half-open, cyclopentadienyl(2, 4- dimethylpentadienyl)vanadocene	483
Magnetic properties of <i>t</i> -butylcyanide adduct of half-open, cyclopentadienyl (2, 4-dimethylpentadienyl)vanadocene	484
Magnetic properties of <i>t</i> -butylcyanide adduct of half-open, 2, 4-dimethylpentadienylvanadocene	485
Magnetic properties of binuclear hexamethyldisilazanato vanadium(III) compound	486
Magnetic properties of four membered vanadacycle ring supported by bulky silazanate	487
Magnetic properties of pyridine adduct of four membered vanadacycle ring supported by bulky silazanate	488
Magnetic properties of diphenylacetylene adduct of six membered vanadacycle ring supported by bulky silazanate	490
Magnetic properties of bis(trimethylsilyl)lacetylene adduct of six membered vanadacycle ring supported by bulky silazanate	491
Magnetic properties of asymmetric dinuclear oxocitrato vanadate(IV)	492
Magnetic properties of hexavanadium(IV) polyoxo alkoxide anion cluster with rubidium cation . . .	494
Magnetic properties of mixed valence hexavanadium(V, IV) polyoxo alkoxide anion cluster with trimethylammonium cation	496
Magnetic properties of a new perovskite-related vanadate with V ^{III} /V ^{IV} ordering	498

Magnetic properties of oxovanadium(IV) complex with 3-salicyliden- acetylacetonediimino-N, N-bis(anthranilic acid)	500
Magnetic properties of oxovanadium(IV) complex of Schiff-base ligand	502
Magnetic properties of oxovanadium(IV) sulphate complex with 2, 3-dihydrazinoquinoxaline	504
Magnetic properties of mixed ligand oxovanadium(IV) complex with salicylic acid and bipyridine	505
Magnetic properties of mixed ligand oxovanadium(IV) complex with salicylic acid and 1, 10-phenanthroline	506
Magnetic properties of heptachlorodivanadium(III) as tetraethyl- ammonium salt	507
Magnetic properties of trisaminopyridinatovanadium(III) complex	509
Magnetic properties of oxovanadium(IV) complex of kojic acid	510
Magnetic properties of vanadyl(IV) complex with thioglycolic acid	511
Magnetic properties of vanadyl(IV) complex of Schiff-base derived from anthranilic acid and acetylacetone	512
Magnetic properties of vanadyl(IV) complex with Schiff-base derived from anthranilic acid and salicylaldehyde	514
Magnetic properties of vanadyl(IV) complex with Schiff-base derived from anthranilic acid and 5-bromo-salicylaldehyde	516
Magnetic properties of vanadyl(IV) complex with Schiff-base derived from anthranilic acid and 3-methoxy-salicylaldehyde	518
Magnetic properties of triclinic form of decahydrated Keggin anion	520
Magnetic properties of monoclinic form of nano-hydrated Keggin anion	521
Magnetic properties of oxovanadium complex with dinucleating tetraaminodiphenol	522
Magnetic properties of oxovanadium complex with dinucleating tetraaminodiphenol	524
Magnetic properties of dimeric vanadium(III) complex with chloride and tetrahydrofuran	526
Magnetic properties of electrogenerated oxo-vanadium complex with a tridentate ONO ligand	527
Magnetic properties of electrogenerated oxo-vanadium complex with a tridentate ONO ligand	529
Magnetic properties of electrogenerated oxo-vanadium complex with azobenzene-2, 2'-diolate	531
Magnetic properties of diiodo vanadium(IV) complex with a Schiff-base	533
Magnetic properties of oxovanadium(IV) complex with a Schiff-base	534
Magnetic properties of diiodo vanadium(IV) complex with a Schiff-base	535
Magnetic properties of diiodovanadium(IV) complex with Schiff-base	536
Magnetic properties of diiodovanadium(IV) complex with a Schiff-base	537
Magnetic properties of chloro-bridged vanadium(III) complex with tetrahydrofuran	538
Magnetic properties of dinuclear copper(II) complex anion with oxovanadium cation	540
Magnetic properties of octanuclear mixed-metal (vanadium-tungsten) cluster	542
Magnetic properties of oxo-vanadium(IV) complex with tetradentate Schiff-base	543

Magnetic properties of vanadium(IV) complex with tetradentate Schiff-base as tetrafluoroborate salt	544
Magnetic properties of oxo-vanadium(IV) complex with tetradentate Schiff-base as tetrafluoroborate salt	545
Magnetic properties of bis(μ -oxo) vanadium(IV) complex with tetradentate Schiff-base as tetrafluoroborate salt	546
Magnetic properties of tris(μ -oxo) vanadium(IV) complex with tetradentate Schiff-base as tetrafluoroborate salt	547
Magnetic properties of tris(μ -oxo) vanadium(IV) complex with tetradentate Schiff-base, as tetrafluoroborate salt	549
Magnetic properties of tris(μ -oxo) vanadium(IV) complex with tetradentate Schiff-base, as tetraphenylborate salt	550
Magnetic properties of tris(μ -oxo) vanadium(IV) complex with tetradentate Schiff-base, as trifluoromethanesulfonate salt	552
Magnetic properties of bis(μ -oxo) vanadium(III, IV) complex with tetradentate Schiff-base, as triiodate salt	553
Magnetic properties of hexakis(acetonitrile)vanadium(II) tetraiodide	554
Magnetic properties of mixed valence organic-based layered vanadium oxide	555
Magnetic properties of arsenic rich polyoxovanadium anion	557
Magnetic properties of homobimetallic vanadium d^1 - d^1 complex	559
Magnetic properties of homobimetallic vanadium d^1 - d^1 complex	561
Magnetic properties of homobimetallic vanadium d^2 - d^2 complex with triyne	563
Magnetic properties of vanadium(IV) complex with kojic acid	565
Magnetic properties of vanadyl-biguanide complex	566
Magnetic properties of vanadyl-metformin complex	568
Magnetic properties of vanadyl-phenformin complex	570
Magnetic properties of vanadyl complex with metformin and dimethylformin	572
Magnetic properties of vanadyl(IV) sulphate	573
Magnetic properties of vanadyl complex with 2-mercaptopyridine-N-oxide	574
Magnetic properties of mononuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone and 2, 2'-bipyridine	575
Magnetic properties of mononuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone and 2, 2'-bipyridine	577
Magnetic properties of mononuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone and 1, 10-phenanthroline	578
Magnetic properties of mononuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone and 1, 10-phenanthroline	580
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 2, 2'-bipyridine a heterocyclic nitrogen base	582

Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone 2, 2'-bipyridine and 2, 6-diaminopyridine	584
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 2, 2'-bipyridine and <i>m</i> -phenylenediamine	586
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 2, 6-diaminopyridine and 2, 2'-bipyridine	588
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 1, 10-phenanthroline and <i>m</i> -phenylenediamine	590
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 1, 10-phenanthroline and 2, 6-diaminopyridine	592
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 1, 10-phenanthroline and <i>m</i> -phenylenediamine	594
Magnetic properties of binuclear mixed ligand complex of oxovanadium(IV) with heterocyclic β -diketone, 1, 10-phenanthroline and 2, 6-diaminopyridine	596
Magnetic properties of bis(acetylacetonato)-chloro-(4-nitrophenoxo)- vanadium(IV)	598
Magnetic properties of bis(acetylacetonato)-chloro-(4-methoxyphenoxo)-vanadium(IV)	599
Magnetic properties of bis(acetylacetonato)-chloro-{4-methyl-2,6-di(<i>tert</i> -butyl)phenoxo}-vanadium(IV)	600
Magnetic properties of bis(acetylacetonato)-bis(4-methoxyphenoxo)- vanadium(IV)	601
Magnetic properties of vanadium(IV) complex with Schiff-base ligand derived from benzoylacetophenone and ethanolamine	602
Magnetic properties of vanadium(IV) complex with Schiff-base ligand derived from benzoylacetone and ethanolamine	603
Magnetic properties of vanadium(IV) complex with Schiff-base ligand derived from acetylacetone and ethanolamine	604
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from benzoylacetophenone and ethanolamine	605
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from benzoylacetone and ethanolamine	606
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from acetylacetone and ethanolamine	607
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from benzoylacetophenone and <i>o</i> -aminophenol	608
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from benzoylacetone and <i>o</i> -aminophenol	609
Magnetic properties of oxovanadium(IV) complex with Schiff-base ligand derived from acetylacetone and <i>o</i> -aminophenol	610
Magnetic properties of pyridine adduct of oxovanadium(IV) complex of Schiff-base complex derived from benzoylacetophenone and <i>o</i> -aminophenol	611
Magnetic properties of <i>n</i> -pentylamine adduct of oxovanadium(IV) Schiff-base complex derived from benzoylacetophenone and <i>o</i> -aminophenol	612

Magnetic properties of thiophene adduct of oxovanadium(IV) Schiff-base complex derived from benzoylacetophenone and <i>o</i> -aminophenol	613
Magnetic properties of vanadium(III) complex of salicylaldehyde-4-phenyl thiosemicarbazone	614
Magnetic properties of vanadium(III) complex of salicylaldehyde-4- <i>p</i> -bromophenyl-thiosemicarbazone	615
Magnetic properties of vanadium(III) complex of salicylaldehyde-4- <i>p</i> -chlorophenyl-thiosemicarbazone	616
Magnetic properties of vanadium(III) complex of salicylaldehyde-4- <i>p</i> -tolyl-thiosemicarbazone	617
Magnetic properties of vanadium(III) complex of 5-bromo-salicylaldehyde- <i>p</i> -tolyl-thiosemicarbazone	618
Magnetic properties of vanadium(III) complex of 5-chloro-salicylaldehyde- <i>p</i> -tolyl-thiosemicarbazone	619
Magnetic properties of first oxalate-bridged dinuclear Co ^{II} -V ^{IV} O complex	620
Magnetic properties of first oxalate-bridged dinuclear Co ^{II} - V ^{IV} O complex	622
Magnetic properties of niobium(IV) tetrachloride adduct of 7-azaindole	624
Magnetic properties of tetrakis(hexafluoroacetylacetato)niobium(IV)	625
Magnetic properties of tetrakis(acetylacetato)niobium(IV)	626
Magnetic properties of niobium(IV) complex with anionic organic amides	627
Magnetic properties of tetravalent niobium phenyl stabilized by anionic organic amides	628
Magnetic properties of tetravalent niobium methyl stabilized by anionic organic amides	629
Magnetic properties of tetravalent niobium enolate stabilized by anionic organic amides	630
Magnetic properties of 3-indole carboxylate complex of niobocene dichloride	631
Magnetic properties of mononuclear tantalum(0) carbonyl compound	632
Magnetic properties of monomeric chromium(II) aryl	634
Magnetic properties of monomeric pyridine adduct of chromium(II) aryl	635
Magnetic properties of dimeric chromium(II) aryl	636
Magnetic properties of monomeric chromium(III) aryl	637
Magnetic properties of monomeric chromium(III) aryl	638
Magnetic properties of polycyclophosphane complex of cyclopentadienyl chromium(I)	639
Magnetic properties of heteronuclear sulfide bridged chromium and rhenium containing cluster (<i>cis</i> -isomer)	640
Magnetic properties of heteronuclear sulfide bridged chromium and rhenium containing cluster (<i>trans</i> -isomer)	642
Magnetic properties of tetranuclear sulfide bridged chromium and rhenium cluster containing carbonyl and nitrosyl ligands	644
Magnetic properties of triangular sulfide bridged chromium and rhenium cluster containing carbonyl and nitrosyl ligands	646

Magnetic properties of tetranuclear sulfide bridged chromium and rhenium cluster containing carbonyl ligand	648
Magnetic properties of triangular sulfide and oxothiolate bridged chromium and rhenium cluster containing chloro, carbonyl and nitrosyl ligands	650
Magnetic properties of sulfide and thiolate bridged triangular cluster with CrRe_2 core containing carbonyl and nitrosyl ligands	651
Magnetic properties of sulfide and thiolate bridged triangular cluster with CrRe_2 core containing carbonyl and nitrosyl ligands	652
Magnetic properties of sulfide and thiolate bridged antiferromagnetic cluster with Cr_2Re core containing carbonyl and nitrosyl ligands	653
Magnetic properties of <i>fac</i> (S)-tris(2-aminoethanethiolato)chromium(III)	655
Magnetic properties of S-bridged zinc bromide polynuclear complex composed of <i>fac</i> -(S)-[Cr(aet) ₃] units	656
Magnetic properties of S-bridged nickel bromide polynuclear complex composed of <i>fac</i> -(S)-[Cr(aet) ₃] units	658
Magnetic properties of S-bridged zinc nitrate polynuclear complex composed of <i>fac</i> -(S)-[Cr(aet) ₃] units	660
Magnetic properties of S-bridged nickel nitrate polynuclear complex composed of <i>fac</i> -(S)-[Cr(aet) ₃] units	661
Magnetic properties of imidodiphosphinoselenido complex of chromium(III)	662
Magnetic properties of chromium(III) complex with 5-(2-hydroxy-benzylidene) aminobenzopyrazole	664
Magnetic properties of chromium(III) telluroate as 1, 2-bis(dimethyl- phosphino)ethane	665
Magnetic properties of mixed ligand complex of chromium(III) picrates with isonicotinic acid hydrazide	666
Magnetic properties of chromium(III) complex with indan-1, 3-dione-2-isonicotinoyl hydrazone	667
Magnetic properties of chromium(III) complex of monophenylbutazone	668
Magnetic properties of chromium(III) complex with tetrasodium calix[4]arene sulphonate (a phenolic macrocycle)	669
Magnetic properties of chromium(III) complex with oxyphenbutazone	671
Magnetic properties of chromium(III) complex of bis(bismuthiol-1-yl)- borate	672
Magnetic properties of chromium(III) complex with 3-hydrazino-5, 6-diphenyl-1, 2, 4-triazine	673
Magnetic properties of chromium(III) complex with diclofenac sodium	674
Magnetic properties of tetracyanonotrosylchromate(I) complex with 8-hydroxyquinoline	675
Magnetic properties of tetracyanonotrosylchromate(I) complex with benzothiazole	676
Magnetic properties of tetracyanonotrosylchromate(I) complex with 2-methylbenzothiazole	677
Magnetic properties of tetracyanonotrosylchromate(I) complex with imidazole	678
Magnetic properties of tetracyanonotrosylchromate(I) with phenir amine maleate	679

Magnetic properties of chromium(III) chloride complex with 1, 3-disuccinimide(4-phenyl-5-ethoxycarbonyl-6-methylpyrimidine-2-one)	680
Magnetic properties of polynuclear chromium(III) chloride complex with 2, 3-dihydrazinoquinoxaline	681
Magnetic properties of Cr(III) complex of 3-(3, 5-dimethylpyrazol-1-yl)quinoxaline-2-one	683
Magnetic properties of heterotetranuclear $\text{Cr}_2^{\text{III}}\text{-Fe}_2^{\text{III}}$ complex of 1, 4, 7-trimethyl-1, 4, 7-triazacyclononane and salicylaldehyde oxime	684
Magnetic properties of a novel 14-membered metallamacrobicycle containing two chromium(III) ions as part of the ring skeleton	686
Magnetic properties of a novel heterotetranuclear $[\text{Cr}^{\text{III}}_2\text{Mn}^{\text{III}}_2(\mu_3\text{-O})_2]^{8+}$ core with an $S_T=0$ spin ground state	688
Magnetic properties of 5-fluorouracil complex with Cr(III)	691
Magnetic properties of mixed ligand, 5-fluorouracil and histidine complex with Cr(III)	692
Magnetic properties of trisaminopyridinatochromium(III) complex	693
Magnetic properties of heterodinuclear copper(II)-chromium(III) complex with N, N'-bis(3-aminopropyl)oxamide and 2, 2'-bipyridine	695
Magnetic properties of heterodinuclear copper(II)-chromium(III) complex with N, N'-bis(3-aminopropyl)oxamide and 1, 10-phenanthroline	697
Magnetic properties of heterodinuclear copper(II)-chromium(III) complex with N, N'-bis(3-aminopropyl)oxamide and 5-nitro-1, 10-phenanthroline	699
Magnetic properties of heterodinuclear copper(II)-chromium(III) complex with N, N'-bis(3-aminopropyl)oxamide and 5-chloro-1, 10-phenanthroline	701
Magnetic properties of 2, 2'-bipyridyl-oxalate bridged heterotrinnuclear complex, NiCr_2	703
Magnetic properties of chromium(II) methylphosphonate	705
Magnetic properties of chromium(III) complex with L(-)cysteine	707
Magnetic properties of dinuclear chromium(III) complex with L(-)cysteine	708
Magnetic properties of chromium(III) complex with L(-)cystine	709
Magnetic properties of dinuclear chromium(III) complex with L(-)cystine	710
Magnetic properties of chromium(III) mixed ligand 5, 5'-dithiobisnitrobenzoic acid and L(-)cysteine	711
Magnetic properties of chromium(III) chloride complex with dibenzylidysulphide	712
Magnetic properties of chromium(III) complex with α -mercaptosuccinic acid	713
Magnetic properties of dinuclear chromium(III) complex with 5, 5'-dithiobisnitrobenzoic acid	714
Magnetic properties of a molecular cluster compound $\text{Cr}_6\text{Te}_8(\text{PEt}_3)_6$	715
Magnetic properties of chromium(III) trispicolate	717
Magnetic properties of di- μ -hydroxotetrapicolinatodichromium(III)	718
Magnetic properties of chromium(II) complex with permethylindenyl	719
Magnetic properties of chromium(III) complex with permethylindenyl	721

Magnetic properties of chromium(II) diphenylamide	722
Magnetic properties of chromium(II) dicyclohexylamide	723
Magnetic properties of chromium(II) di-isopropylamide	724
Magnetic properties of tetrahydrofuran complex with chromium(II) diphenylamide	725
Magnetic properties of pyridine complex with chromium(II) diphenylamide	727
Magnetic properties of tetrahydrofuran complex with chromium(II) diphenylamide	728
Magnetic properties of chromium(II) complex with phenothiazinyl and excess tetrahydrofuran . . .	729
Magnetic properties of sodium diphenylamidochromate(II)	730
Magnetic properties of lithium diethylamidochromate(II)	731
Magnetic properties of lithium diethylamidochromate(II)	732
Magnetic properties of dimeric chromium(II) complex with 2-methylaminopyridine	733
Magnetic properties of dimeric chromium(II) complex with aminopyridine	734
Magnetic properties of dithioxalato-bridged heterometallic chromium(III)-trinickel(II) complex	735
Magnetic properties of binuclear bis(acetylacetonato)chromium(III) complex bridged by imidazole-4, 5-dicarboxylate	737
Magnetic properties of binuclear bis(acetylacetonato)chromium(III) complex bridged by 1, 2, 3-triazole-4, 5-dicarboxylate	739
Magnetic properties of binuclear bis(acetylacetonato)chromium(III) complex bridged by pyrazole 3, 5-dicarboxylate	741
Magnetic properties of chromium(III) aldonate complex	743
Magnetic properties of chromium(III) complex with methylhydrazine- carboxylate ligand	745
Magnetic properties of binuclear bis(acetylacetonato)chromium(III) complex bridged by pyrazolate and hydroxide groups	746
Magnetic properties of dinuclear chromium(III) complex with bis(2-diisopropylphosphinoethyl) amidolithium reagent	748
Magnetic properties of tetra- <i>n</i> -butylammonium tetrakisothiocyanato chromate(II) (red-form)	750
Magnetic properties of tetra- <i>n</i> -butylammonium tetrakisothiocyanato chromate(II) (blue form)	751
Magnetic properties of half-sandwich, imidochromium(V) compound	753
Magnetic properties of half-sandwich imidochromium(V) compound	754
Magnetic properties of imidochromium(V) Schiff-base compound	755
Magnetic properties of imidochromium(V) Schiff-base compound	756
Magnetic properties of octanuclear mixed metal (chromium-molybdenum) cluster	757
Magnetic properties of octanuclear mixed-metal (chromium-tungsten) cluster	758
Magnetic properties of octanuclear mixed-metal (Cr + Fe-Mo) cluster	759
Magnetic properties of μ -oxo, tetranuclear Cr ₂ -Mn ₂ complex with 1, 4, 7-triazacyclononane and salicylaldehyde oxime	760

Magnetic properties of pentanuclear μ -cyano $\text{Cr}^{\text{III}}\text{Ni}^{\text{II}}$ (radical) ₂ complex	762
Magnetic properties of dodecanuclear chromium(III) cage with carboxylate ligand	764
Magnetic properties of two-dimensional, cyano-bridged Ni(II)-Cr(I) complex with 3, 10-dimethyl-1, 3, 5, 8, 10, 12-hexaazacyclotetradecane	766
Magnetic properties of tetraacetatodimolybdenum(II)	768
Magnetic properties of molybdenum complex of the anticancer drug 6-mercaptopurine(II)	770
Magnetic properties of molybdenum complex of the anticancer drug 6-mercaptopurine(II)	772
Magnetic properties of molybdenum(V) complex with polystyrene-anchored Schiff base ligand	774
Magnetic properties of oxomolybdenum(V) chloride complex of salicylaldehydemorpholine-N-thiohydrazone	775
Magnetic properties of oxomolybdenum(V) chloride complex of 2-hydroxyacetophenonemorpholine-N-thiohydrazone	776
Magnetic properties of oxomolybdenum(V) thiocyanate complex of salicylaldehydemorpholine-N-thiohydrazone	777
Magnetic properties of hydrotris(pyrazolyl)borate complex of molybdenum carbonyl	778
Magnetic properties of hydrotris (pyrazolyl) borate complex of molybdenum carbonyl	779
Magnetic properties of cyclopentadienylmolybdenum(IV) chloride	780
Magnetic properties of cyclopentadienylmolybdenum(IV) bromide	781
Magnetic properties of cyclopentadienylmolybdenum(IV) iodide	782
Magnetic properties of trinuclear complex MoCu_2 possessing a hexathiolatomolybdenum(IV) core	783
Magnetic properties of triiodo molybdenum(III) complex with 1, 4, 7-trimethyl-1, 4, 7-triazacyclononane	784
Magnetic properties of diiodo oxomolybdenum(V) complex with 1, 4, 7-trimethyl-1, 4, 7-triazacyclononane	785
Magnetic properties of dimethoxy oxomolybdenum(V) complex with 1, 4, 7-trimethyl-1, 4, 7-triazacyclononane	786
Magnetic properties of dinuclear mixed valent (Mo^{V} , Mo^{VI}) oxo bridged iodo dioxomolybdenum complex with 1, 4, 7-trimethyl-1, 4, 7-triazacyclononane	787
Magnetic properties of η -cycloheptatrienyl derivative of molybdenum	788
Magnetic properties of η -cycloheptatrienyl derivative of molybdenum	790
Magnetic properties of η -cycloheptatrienyl derivative of molybdenum	792
Magnetic properties of oxomolybdenum(V) complex with 2-aminocyclopent-1-ene-1-carbodithiolate	794
Magnetic properties of mononuclear oxomolybdenum(V) complex with N-ethyl-2-aminocyclopent-1-ene-1-carbodithiolate	795
Magnetic properties of mononuclear oxomolybdenum(V) complex with N-propyl-2-aminocyclopent-1-ene-1-carbodithiolate	796

Magnetic properties of mononuclear oxomolybdenum(V) complex with N-butyl-2-aminocyclopent-1-ene-1-carbodithiolate	797
Magnetic properties of (η -cyclopentadienyl)imido derivative of molybdenum	798
Magnetic properties of mononuclear, 15-electron molybdenum complex with 1, 2-bis (diisopropylphosphino)ethane	800
Magnetic properties of mononuclear, 16-electron molybdenum complex with 1, 2-bis (diisopropylphosphino)ethane	801
Magnetic properties of oxomolybdenum(V) complex with hydrazone derived from 6-methyl-4-hydroxy-2-hydrazinopyrimidine and salicylaldehyde	802
Magnetic properties of oxomolybdenum(V) complex with hydrazone derived from 6-methyl-4-hydroxy-2-hydrazinopyrimidine and 5-methyl- salicylaldehyde	803
Magnetic properties of oxomolybdenum(V) complex with hydrazone derived from 6-methyl-4-hydroxy-2-hydrazinopyrimidine and 5-chloro-salicylaldehyde	804
Magnetic properties of oxomolybdenum(V) complex with hydrazone derived from 6-methyl-4-hydroxy-2-hydrazinopyrimidine and 5-bromo- salicylaldehyde	805
Magnetic properties of oxomolybdenum(V) complex with hydrazone derived from 6-methyl-4-hydroxy-2-hydrazinopyrimidine and 3-methoxy- salicylaldehyde	806
Magnetic properties of bimetallic, phosphido-bridged complex of tungsten	808
Magnetic properties of <i>mer</i> -trichlorotris(tetrahydrofuran)tungsten(III)	809
Magnetic properties of trinuclear complex WCu_2 possessing a hexathiolatotungsten(II)	811
Magnetic properties of (2, 4-dimethylpentadienyl)bis(cyclopentadienyl)- dimanganese(II)	812
Magnetic properties of dinuclear manganese complex of N, N'-ethylenebis(acetylacetimine)	813
Magnetic properties of methyl manganese Schiff base complex	814
Magnetic properties of hexamesityltrimanganese(II)	815
Magnetic properties of dimeric phenylmanganese(II) ion-pair with tetraphenylborate	817
Magnetic properties of monomeric mesitylmanganese(II) ion-pair with tetraphenylborate	819
Magnetic properties of manganese(II) complex of isoorotic acid	820
Magnetic properties of manganese(II) complex of nitronyl nitroxide carboxylic acid	821
Magnetic properties dinuclearmanganese(II) complex with 2, 2'-bipyrimidine and thiocyanate ligands	823
Magnetic properties dinuclearmanganese(II) complex with 2, 2'-bipyrimidine and selenocyanate ligands	824
Magnetic properties of dimer of trinuclear manganese complex	825
Magnetic properties of octadecanuclear manganese(II) complex	826
Magnetic properties of diaqua-bis(1, 3, 6, 7-tetramethyl-pterdine-2, 4-(1H, 3H)-dione- O^4 , N^5)manganese(II) perchlorate	828
Magnetic properties of manganese(II) tellurostannates	829
Magnetic properties of 2-(4-chlorophenyl)-4, 4, 5, 5-tetramethyl- imidazoline-1-oxyl-3-oxide-bis(hexafluoroacetylacetonato)manganese(II)	831

Magnetic properties of hexa[2-(4-methylphenyl)-4, 4, 5, 5-tetramethyl imidazoline-1-oxyl-3-oxide-bis(hexafluoroacetylacetonato)manganese(II)]	833
Magnetic properties of di[1-hydroxo-2-furfural-4, 4, 5, 5-tetramethyl-4, 5-dihydro-1H-imidazoline-bis(hexafluoroacetylacetonato)manganese(II)]	835
Magnetic properties of di(2-furfural-4, 4, 5, 5-tetramethylimidazoline-1-oxyl-3-oxide)-bis(hexafluoroacetylacetonato)manganese(II)	837
Magnetic properties of dinuclear copper(II)-manganese(II) complex of a phenol-based macrocycle	839
Magnetic properties of one-dimensional azide-bridged manganese(III) uniform chain	841
Magnetic properties of tris{[bis(triphenylphospho)amide]hexacyano- manganate(III)}	843
Magnetic properties of oxo-centered trinuclear manganese complex	845
Magnetic properties of oxo-centered trinuclear manganese complex	847
Magnetic properties of oxo-centered trinuclear manganese complex	849
Magnetic properties of oxo-centered trinuclear manganese complex	851
Magnetic properties of oxo-centered trinuclear manganese complex	853
Magnetic properties of one dimensional chain compound <i>catena</i> -poly[μ -4, 4'-bis-1, 2, 4-triazole-N1N1]aquabis(4, 4'-bis-1, 2, 4-triazole-N1)(thiocyanato-N)manganese(II) thiocyanato]	855
Magnetic properties of oximate-bridged heterobinuclear $[\text{Cu}^{\text{II}}\text{Mn}^{\text{II}}]_2$ complex	856
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and phthalic acid	858
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and phthalic acid	860
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and phthalic acid	861
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and phthalic acid	862
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and malonic acid	863
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and malonic acid	864
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and malonic acid	865
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and succinic acid	866
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and succinic acid	867
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and succinic acid	868
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and succinic acid	869

Magnetic properties of manganese(III) complex containing tetradentate Schiff base and glutaric acid	870
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and glutaric acid	871
Magnetic properties of manganese(III) complex containing tetradentate Schiff base and glutaric acid	872
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxyphenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamide	873
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-5-bromophenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamide	874
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3-ethoxyphenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamide	875
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3-methoxyphenyl-5-bromophenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamide	876
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3,5-dichlorophenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamido	877
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3,5-dibromophenyl)-N-(2-hydroxy-4-methylphenyl)but-3-enamido	878
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3-bromo-5-chlorophenyl)-N-(2-hydroxy-4-methyl phenyl)but-3-enamido	879
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxy-3-bromo-5-nitrophenyl)-N-(2-hydroxy-4-methyl phenyl)but-3-enamido	880
Magnetic properties of 1-D- μ_1 , $_3$ -bridged Mn(II) compound with 2, 13-dimethyl-3, 6, 9, 12, 18-pentaazabicyclo{12.3.1}octadeca-1(18), 2, 12, 14, 16-pentaene	881
Magnetic properties of (μ -oxo)(μ -carboxylato)dimanganese(III) polyimidazole complex	883
Magnetic properties of ($\pm$)-5, 7, 7, 12, 14, 14-hexamethyl-1, 4, 8, 11-tetraazacyclotetradecane-N, N'-bis(3-aminopropyl)oxamide- manganese(II)copper(II) triflate	885
Magnetic properties of manganese(III) N, N'-(ethylene)bis(salicylidene- imine)selenocyanate complex	887
Magnetic properties of manganese(III) N, N'-(propylene)bis(salicylidene- imine)selenocyanate complex	888
Magnetic properties of manganese(III) complex with Schiff base ligand, 3-aza-4-(2-hydroxyphenyl)-N-(2-hydroxy-5-chlorophenyl)but-3-enamide	889
Magnetic properties of manganese(III) N, N'-(<i>o</i> -phenylene)bis- (salicylideneimine) selenocyanate complex	890
Magnetic properties of manganese(III) 1, 2-dihydro-1, 5-dimethyl-2-phenyl-4-(2'-hydroxynaphthylazo)3H-pyrazol-3-one thiocyanate complex	891
Magnetic properties of manganese(III) 1, 2-dihydro-1, 5-dimethyl-2-phenyl-4-(2'-hydroxynaphthylazo)3H-pyrazol-3-one selenocyanate complex	892
Magnetic properties of manganese(II) complex with 5-(2-hydroxy- benzylidene) aminobenzopyrazole	893

Magnetic properties of manganese(II) tellurolate as 1, 2-bis(dimethyl-phosphino)ethane	894
Magnetic properties of manganese(II) tellurolate as 4- <i>tert</i> -butylpyridine adduct	895
Magnetic properties of heterodinuclear manganese-copper complex of <i>o</i> -phenylenebis-(oxamic acid)	896
Magnetic properties of a tetranuclear rhomblike cluster of manganese(II)	898
Magnetic properties of manganese(III) complex of doubly-hydrated hexafluoroacetylacetone	900
Magnetic properties of dodecanuclear, $Mn_{11}Cr$ complex of acetic acid	902
Magnetic properties of manganese(II) complex with 3-salicylidene acetylacetonediimino-N, N-bis(anthranilic acid)	904
Magnetic properties of manganese(II) complex with 2, 2'-dihydroxy- oxanilide	906
Magnetic properties of hetero trimetallic ($Mn-Zr_2$) alkoxide derivative	908
Magnetic properties of chloro-4-(2-pyridyl)-1-(2, 4-dihydroxy- benzaldehyde)-3-thiosemicarbazonomanganese(II)	909
Magnetic properties of manganese(III) complex of monophenylbutazone	910
Magnetic properties of manganese(II) complex with oxyphenbutazone	911
Magnetic properties of manganese(II) complex of bis(bismuthiol-1-yl)- borate	912
Magnetic properties of manganese(II) complex with 3-hydrazino-5, 6-diphenyl-1, 2, 4-triazine	913
Magnetic properties of manganese(II) complex of a tripodal ligand tris(2-benzimidazolymethyl)amine	914
Magnetic properties of manganese(II) complex of a tripodal ligand N,N',N''-triethyl-tris(2-benzimidazolymethyl)amine	915
Magnetic properties of trimeric manganese(II) complex of a tripodal ligand tris (2-benzimidazolymethyl)amine	916
Magnetic properties of trimeric manganese(II) complex of a tripodal ligand N.N',N''-triethyl-tris(2-benzimidazolymethyl)amine	918
Magnetic properties of dimeric manganese(II) complex of a tripodal ligand tris (2-benzimidazolymethyl)amine	919
Magnetic properties of manganese(II) complex with diclofenac sodium	920
Magnetic properties of dimeric manganese(II) complex of diacetylmonoximeisonicotinoyl hydrazone	922
Magnetic properties of manganese(II) chloride complex with 1, 3-disuccinimide(4-phenyl-5-ethoxycarbonyl-6-methylpyrimidine-2-one)	924
Magnetic properties of polynuclear manganese(II) chloride complex with 2, 3-dihydrazinoquinoxaline	925
Magnetic properties of mononuclear manganese(II) complex with 13-membered macrocycle, 1, 5, 8, 11-tetra-azacyclotridecane, 2, 4-dione	927
Magnetic properties of trinuclear lanthanum(III)-manganese(II), La_2-Mn , complex with 13-membered macrocycle, 1, 5, 8, 11-tetra-azacyclo tridecane-2, 4-dione	928

Magnetic properties of manganese(III) thiocyanate complex with hexadentate Schiff base derived from 1-phenyl-3-methyl-4-benzoyl-2-pyrazoline-5-one and triethylenetetraamine	929
Magnetic properties of manganese(III) nitrate complex with hexadentate Schiff base derived from 1-phenyl-3-methyl-4-benzoyl-2-pyrazoline-5-one and triethylenetetraamine	930
Magnetic properties of manganese(III) perchlorate complex with hexadentate Schiff base derived from 1-phenyl-3-methyl-4-benzoyl-2-pyrazoline-5-one and triethylenetetraamine	931
Magnetic properties of manganese(III) thiocyanate complex with hexadentate Schiff base derived 1-phenyl-3-methyl-4-acetyl-2-pyrazoline-5-one and triethylenetetraamine	932
Magnetic properties of manganese(III) nitrate complex with hexadentate Schiff base derived from 1-phenyl-3-methyl-4-acetyl-2-pyrazoline-5-one and triethylenetetraamine	933
Magnetic properties of manganese(III) perchlorate complex with hexadentate Schiff base derived from 1-phenyl-3-methyl-4-acetyl-2-pyrazoline-5-one and triethylenetetraamine	934
Magnetic properties of manganese(II) complex with 3-(3, 5-dimethylpyrazol-1-yl) quinoxaline-2-one	935
Magnetic properties of chloranilate-bridged manganese(II) chain compound	936
Magnetic properties of bis[6-(<i>tert</i> -butyliminomethyl)phenylamido] manganese(II)	938
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	939
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	941
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	943
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	945
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	947
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	949
Magnetic properties of manganese(III) complex with tetradentate aryl substituted Schiff-base ligand	951
Magnetic properties of oxalate bridged dinuclear Mn ^{II} -V ^{IV} O complex with 1, 10-phenanthroline as auxillary ligand	953
Magnetic properties of tris(quinoxaline-2, 3-dithiolato)manganate(IV) as tetraphenylphosphate salt	955
Magnetic properties of manganese(II) dimethylformamide complex cation with bis-[di(quinoxaline-2, 3-dithiolato)cuprate(II)] anion	957
Magnetic properties of manganese(III) complex with chiral salen-type ligand	959
Magnetic properties of manganese(III) complex with chiral salen-type ligand	961
Magnetic properties of manganese(II) complex with asymmetric tetraiminodiphenol macrocyclic ligand	963
Magnetic properties of di- μ -oxo-dimanganese(IV) complex with Schiff-base	965

Magnetic properties of di- μ -oxo-dimanganese(IV) complex with Schiff-base	967
Magnetic properties of tetradentate di- μ -oxo-dimanganese(IV) complex with Schiff-base	969
Magnetic properties of tetradentate di- μ -oxo-dimanganese(IV) complex with Schiff-base	971
Magnetic properties of di- μ -oxo-dimanganese(IV) complex with tetradentate Schiff-base	973
Magnetic properties of dinuclear manganese(II) hexafluoropentane-2, 4-dione complex with pyrazine	975
Magnetic properties of mononuclear manganese(II) hexafluoropentane-2, 4-dione complex with pyrimidine	977
Magnetic properties of mononuclear manganese(II) hexafluoropentane-2, 4-dione complex with pyridazine	978
Magnetic properties of mononuclear manganese(II) hexafluoropentane-2, 4-dione complex with 3-methylpyridazine	980
Magnetic properties of hexacyanomanganate(IV)	982
Magnetic properties of manganese tricyanomethamide	983
Magnetic properties of manganese(II) complex with diclofenac	985
Magnetic properties of one dimensional polymeric bicinate manganese complex	986
Magnetic properties of dinuclear bis- μ -(1, 3- N_3) bridged manganese(II) complex containing macrocyclic amine ligand	988
Magnetic properties of hexanuclear Mn^{III} cluster with $[Mn_6O_4X_4]^{6+}$ core	990
Magnetic properties of hexanuclear Mn^{III} cluster with $[Mn_6O_4X_4]^{6+}$ core	992
Magnetic properties of manganese(II) chloride adduct with 2-(2-pyridyl)-4, 4, 5, 5-tetramethyl-4, 5-dihydro-1 <i>H</i> -imidazolyl-1-oxy-3-oxide	993
Magnetic properties of manganese(II) hexafluoroacetylacetonate complex with 2-(2-pyridyl)-4, 4, 5, 5-tetramethyl-4, 5-dihydro-1 <i>H</i> -imidazolyl-1-oxy-3-oxide	995
Magnetic properties of methoxy bridged manganese(III) dimer with salicylic acid	997
Magnetic properties of heterometallic, Mn-Cu, complex with 1, 3-propylenediamine-N, N, N', N'-tetraacetate	999
Magnetic properties of end-to-end azido bridged binuclear complex of manganese(II), Mn_2 , with tris(6-methyl-2-pyridylmethyl)amine	1001
Magnetic properties of hepta coordinate manganese complex with bicine	1003
Magnetic properties of mononuclear manganese(II) complex with 2, 3-pyrazinedicarboxylic acid	1005
Magnetic properties of polynuclear manganese(II) complex with 2, 3-pyrazinecarboxylic acid	1007
Magnetic properties of dodecanuclear manganese aggregate containing heterocarboxylate ligands	1009
Magnetic properties of manganese(II) complex with D-Gluconic acid	1012
Magnetic properties of dinuclear manganese(II) complex of 2, 6-bis[2-(dimethylamino)-ethyliminomethyl]-4-methylphenolate(-1) and acetate	1013

Magnetic properties of dinuclear manganese(II) complex of 2, 6-bis[2-(diethylamino)-ethyliminomethyl]-4-methylphenolate(-1) and acetate	1015
Magnetic properties of dinuclear manganese(II) complex of 2, 6-bis[2-(dimethylamino)-ethyliminomethyl]-4-methylphenolate(-1) and benzoate	1017
Magnetic properties of dinuclear manganese(II) complex of 2, 6-bis[2-(diethylamino)-ethyliminomethyl]-4-methylphenolate(-1) and benzoate	1019
Magnetic properties of dinuclear chloro bridged manganese(II) chloro complex of 2, 6-bis[2-(dimethylamino)-ethyliminomethyl]-4-methylphenolate(-1)	1021
Magnetic properties of dinuclear chloro bridged manganese(II) chloro complex of 2, 6-bis[2-(diethylamino)-ethyliminomethyl]-4-methylphenolate(-1)	1023
Magnetic properties of manganese(III) complex of tripodal tris[2-(salicylideneamino)ethyl]amine	1025
Magnetic properties of manganese(III) complex of tripodal tris[2-(5-chlorosalicylideneamino)ethyl]amine	1026
Magnetic properties of manganese(III) complex of tripodal tris[2-(5-chlorosalicylideneamino)ethyl]amine	1028
Magnetic properties of tetranuclear manganese(III) complex of tripodal tris[2-(salicylideneamino)ethyl]amine	1030
Magnetic properties of tetranuclear manganese(III) complex of tripodal tris[2-(5-chlorosalicylideneamino)ethyl]amine	1032
Magnetic properties of manganese(III) complex with unsymmetrical Schiff-base	1034
Magnetic properties of manganese(III) complex with unsymmetrical Schiff-base	1036
Magnetic properties of manganese(III) complex with unsymmetrical Schiff-base	1038
Magnetic properties of manganese(III) complex with unsymmetrical Schiff-base	1040
Magnetic properties of manganese(III) complex with unsymmetrical Schiff-base	1042
Magnetic properties of mixed-valence binuclear $\text{Mn}^{\text{IV}}\text{-Mn}^{\text{III}}$ complex with methyl [2-(2-pyridyl)ethyl](2-pyridyl-methyl)amine)	1044
Magnetic properties of manganese(II) diiodo complex with organophosphorous compound	1046
Magnetic properties of macrocyclic tetranuclear manganese(II) complex of a “dimer of dimers” type	1047
Magnetic properties of macrocyclic tetranuclear manganese(II) complex of a “dimer of dimers” type	1049
Magnetic properties of dinuclear manganese(II) complex of unsymmetric phenol based dinucleating ligand with amino and imino chelating arms	1051
Magnetic properties of dinuclear manganese(II) complex of unsymmetric phenol based dinucleating ligand with amino and imino chelating arms	1053
Magnetic properties of dinuclear manganese(II) complex of unsymmetric phenol based dinucleating ligand with amino and imino chelating arms	1055
Magnetic properties of dinuclear manganese(II) complex of unsymmetric phenol based dinucleating ligand with amino and imino chelating arms	1057

Magnetic properties of manganese(III) complex with N, N'-bis(3-ethoxysalicylidene)propane-1, 3-diamine	1059
Magnetic properties of dimanganese(III) with tetranucleating macrocyclic Schiff-base ligand	1060
Magnetic properties of manganese(IV) complex with salicylaldehyde thiosemicarbazone	1062
Magnetic properties of manganese(IV) complex with 5-methyl-salicylaldehyde thiosemicarbazone	1063
Magnetic properties of manganese(IV) complex with 5-bromo-salicylaldehyde thiosemicarbazone	1064
Magnetic properties of manganese(IV) complex with salicylaldehyde-4-phenylthiosemicarbazone	1065
Magnetic properties of manganese(IV) complex with 5-methyl-salicylaldehyde-4-phenylthiosemicarbazone	1066
Magnetic properties of manganese(IV) complex with 5-bromo-salicylaldehyde-4-phenylthiosemicarbazone	1067
Magnetic properties of manganese(III) complex with S-methyl-3-[(2-hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1068
Magnetic properties of manganese(III) complex with S-methyl-3-[(5-methyl-2-hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1069
Magnetic properties of manganese(III) complex with S-methyl-3-[(5-bromo-2-hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1070
Magnetic properties of manganese(III) complex with S-benzyl-3-[(2- hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1071
Magnetic properties of manganese(III) complex with S-benzyl-3-[5- methyl(2-hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1072
Magnetic properties of manganese(III) complex with S-benzyl-3-[5- bromo(2-hydroxyphenyl)methylene]dithiocarbazon and acetic acid	1073
Magnetic properties of manganese(III) complex with S-benzyl-3-[(2- hydroxyphenyl)methylene]dithiocarbazon and acetylacetone	1074
Magnetic properties of terephthalate bridged manganese(II) complex	1075
Magnetic properties of Ruddlesden-Popper Mn^{III}/Mn^{IV} oxides	1077
Magnetic properties of mixed valence manganese cluster (Mn_7) with 2-hydroxymethylpyridine . . .	1079
Magnetic properties of dimanganese imidazol complex with bridging aqua and acetato groups . . .	1080
Magnetic properties of manganese(II) complex with azide and pyrazine ligands	1082
Magnetic properties of dimanganese(II) complex with squaric acid	1084
Magnetic properties of nonamanganese(II) 3×3 portcullis-like grid generated by strict self assembly of metal cation and heptadentate ligand	1086
Magnetic properties of heterometallic, manganese-antimony cubane	1088
Magnetic properties of heterometallic, manganese-arsenic cubane	1089
Magnetic properties of two dimensional ferrimagnet cyanide-bridged, iron(III)-manganese(II) complex with a partial cubane motif	1090

Magnetic properties of new oxide fluoride of mixed metals (LaSrMn)	1092
Magnetic properties of anionic metal dicyanamide with tris(bipyridyl)- iron(II) cation	1094
Magnetic properties of anionic metal dicyanamide with tris(bipyridyl) nickel(II) cation	1096
Magnetic properties of dimanganese(III) complex with enantiopure proline-derived triazacyclononane	1098
Magnetic properties of first mixed valence $Mn^{II, III}$ gallium phosphate phase	1099
Magnetic properties of manganese(II) complex of diclofenac	1101
Magnetic properties of manganese(II) complex with uric acid	1102
Magnetic properties of manganese(III) complex with acetylacetone	1103
Magnetic properties of manganese(II) chloro complex with neutral malonamide	1104
Magnetic properties of manganese(II) bromo complex with neutral malonamide	1105
Magnetic properties of manganese(II) iodo complex with neutral malonamide	1106
Magnetic properties of manganese(II) complex with methylhydrazine- carboxylate ligand	1107
Magnetic properties of manganese(II) complex with 1, 1-dimethylhydrazine carboxylate ligand . . .	1108
Magnetic properties of manganese(II) complex of benzene-1,3,5-tricarboxylic acid	1109