

Contents

Volume 4A

Introduction	1
Part I Cu	15
Magnetic properties of copper(I) complex with verdazyl radical	17
Magnetic properties of copper(II) tellurite complex	20
Magnetic properties of copper(II) complex with 1-(2-chloroethyl)tetrazole	22
Magnetic properties of copper(II) complex with methyl[2-(pyridylmethyl)- {2-((2-pyridylmethyl)amino)ethyl}amine]- copper(II) perchlorate	24
Magnetic properties of copper(II) complex with methyl[2-((pyridylmethyl) amino)-ethyl]amine	26
Magnetic properties of copper(II) complex with bis(2-pyridylmethyl)amine	28
Magnetic properties of dimeric copper(II) complex with tripod ligand, tris- [2-((2-pyridylmethyl)amino)ethyl]amine	30
Magnetic properties of mononuclear copper(II) complex with azacyano-carbanion and pyridine ligand	32
Magnetic properties of dimeric copper(II) complex with tetracyanoquinodimethane and pyridine ligand	34
Magnetic properties of bis-(1,5-diaminotetrazole) dichlorocopper(II)	36
Magnetic properties of copper(II) complex with 2-(2-pyridyl)imino- N-(2-thiazolin-2-yl)-thiazolidine	39

vii

Magnetic properties of copper(II) complex with sulfurated triazoline derivative of glycine	41
Magnetic properties of copper(II) complex with macrocyclic ligand containing thiosemicarbazone moiety	44
Magnetic properties of copper(II) sulphato complex with macrocyclic ligand containing thiosemicarbazone moiety	46
Magnetic properties of copper(II) nitrate complex with macrocyclic ligand containing thiosemicarbazone moiety	48
Magnetic properties of copper(II) chloro complex with macrocyclic ligand containing thiosemicarbazone moiety	50
Magnetic properties of copper(II) complex with chelating resin containing tridentate Schiff-base	52
Magnetic properties of copper(II) complex with N-4-acetophenyl-3-carboxy-salicylaldehyde	54
Magnetic properties of copper(II) complex with N-4-acetophenyl-5-bromo- salicylaldehyde	56
Magnetic properties of copper(II) complex with N-3-acetophenyl-salicylaldehyde	58
Magnetic properties of copper(II) complex with N-4-acetophenyl-salicylaldehyde	60
Magnetic properties of mixed ligand complex of copper(II) with salicylidenevalinate and imidazole	62
Magnetic properties of mononuclear copper(II) complex with salicylidenevalinate	64
Magnetic properties of copper(II) complex with 8-(2-azobenzothiazolyl)-7-hydroxy-4-methylcoumarin	66
Magnetic properties of copper(II) complex with N-isonicotinoyl-N'-p-hydroxythiobenzhydrazine	68
Magnetic properties of copper(II) complex with 8-(2-azothiazolyl)-7-hydroxy-4-methylcoumarin	70
Magnetic properties of ion-pair complex containing 2-amino-5-bromopyridinium cation and tetrabromocuprate anion	72
Magnetic properties of mononuclear copper(II) complex with 3, 5-pyrazoledicarboxylic acid	74
Magnetic properties of copper(II) complex with cyclam	76

Magnetic properties of mixed-ligand copper(II) complex with ethanol and (tetradecachloro-4-carboxytriphenyl)-methyl radical	78
Magnetic properties of mixed-ligand copper(II) complex with pyridine and (tetradecachloro-4-carboxytriphenyl)-methyl radical	80
Magnetic properties of mixed-ligand copper(II) complex with water and (tetradecachloro-4-carboxytriphenyl)-methyl radical	82
Magnetic properties of copper(II) picolinate	84
Magnetic properties of copper(II) complex with N-salicylidene-n-ethanoylhydrazine	88
Magnetic properties of copper(II) ternary complex of 2-hydroxy-acetophenone- 4-hydroxybenzoic acid hydrazone and 2,2'-bipyridine	90
Magnetic properties of copper(II) ternary complex of 2-hydroxy-acetophenone- 4-hydroxybenzoic acid hydrazone and 1,10-phenanthroline	92
Magnetic properties of copper(II) complex with N-salicylidene-n-pentanoylhydrazine	94
Magnetic properties of copper(II) complex with N-salicylidene-n-hexanoylhydrazine	96
Magnetic properties of copper(II) complex with N-salicylidene-n-heptanoylhydrazine	98
Magnetic properties of mixed ligand mononuclear copper(II) complex with phenanthroline and substituted tetradecane	100
Magnetic properties of copper(II) complex with the dicyanoacetic acid methyl ester anion	102
Magnetic properties of copper(II) complex with 2-oxo-pyrimidinate	104
Magnetic properties of copper(II) complex with pyridinecarboxylic acid	106
Magnetic properties of copper(II) complex with pyridinecarboxylic acid	108
Magnetic properties of mixed ligand complex of copper(II) with 2,2'-bipyrimidine and malonic acid	110
Magnetic properties of copper(II) complex with 3,3-bis(2-imidazolyl)propionate	112

Magnetic properties of copper(II) complex with 1,4-diazacycloheptane	115
Magnetic properties of copper(II) 2-methylthionicotinate adduct with nicotinamide	117
Magnetic properties of copper(II) 2-methylthionicotinate adduct with N-methylnicotinamide	119
Magnetic properties of copper(II) chloro complex with substituted hydrazine	121
Magnetic properties of copper(II) complex with cyanodithioformate ...	123
Magnetic properties of copper(II) chloro, thiocyanato complex with an azo dye derived from 14-aminoantipyrine and <i>p</i>-chlorophenol	125
Magnetic properties of copper(II) perchlorato complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-chlorophenol	127
Magnetic properties of copper(II) sulphato complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-chlorophenol	129
Magnetic properties of copper(II) nitrato complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-chlorophenol	131
Magnetic properties of copper(II) acetato complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-chlorophenol	133
Magnetic properties of copper(II) complex with poly-Schiff-base of 5-vinylsalicylaldehyde	135
Magnetic properties of copper(II) complex with antibiotic drug, dicluxacillin	137
Magnetic properties of copper(II) complex with salicylidene-2-aminothiophenol and valine	139
Magnetic properties of copper(II) complex with salicylidene-2-aminothiophenol and histidine	141
Magnetic properties of copper(II) complex with salicylidene-2-aminothiophenol and alanine	143
Magnetic properties of copper(II) complex with salicylidene-2-aminothiophenol and glycine	145
Magnetic properties of copper(II) chloro complex with 5-(2-hydroxy-phenylazo)-2-thiohydantoin	147
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	149

Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	151
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	153
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	155
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	157
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	159
Magnetic properties of copper(II) chlorobenzoate adduct with substituted piperidine	161
Magnetic properties of copper(II) acetato complex with 5-(phenylazo)-2-thiohydantoin	163
Magnetic properties of copper(II) chloro complex with 2-amino-3,5-dichloropyridine	165
Magnetic properties of copper(II) bromo complex with 2-amino-3,5-dichloropyridine	167
Magnetic properties of copper(II) chloro complex with 2-amino-3,5-dibromopyridine	169
Magnetic properties of copper(II) bromo complex with 2-amino-3,5-dibromopyridine	171
Magnetic properties of copper(II) chlorobenzoate adducts with substituted piperidine	173
Magnetic properties of copper(II) chlorobenzoate adducts with substituted piperidine	175
Magnetic properties of mixed ligand copper(II) thiocarboxylate	177
Magnetic properties of mixed ligand copper(II) thiocarboxylate	179
Magnetic properties of mixed ligand copper(II) thiocarboxylate	181
Magnetic properties of mixed ligand copper(II) thiocarboxylate	183
Magnetic properties of copper(II) fluoride complex with imidazole ...	185
Magnetic properties of copper(II) complex with polystyrene supported tridentate Schiff-base	187
Magnetic properties of mixed ligand complex of copper(II) with cytidine and L-tryptophan	189

Magnetic properties of mixed ligand complex of copper(II) with cytidine and L-phenylalanine	191
Magnetic properties of mixed ligand complex of copper(II) with cytidine and L-alanine	193
Magnetic properties of copper(II) chloro complex with <i>cis</i>-3,7-dimethyl-2, 6-octadienthiosemicarbazone	195
Magnetic properties of copper(II) chloro complex with 18-membered [N₄]-macrocycle	197
Magnetic properties of copper(II) chloro complex with 16-membered [N₄]-macrocycle	199
Magnetic properties of copper(II) chloro complex with 18-membered [N₄]-macrocycle	201
Magnetic properties of mixed ligand copper(II) thiocarboxylate	203
Magnetic properties of mixed ligand copper(II) thiocarboxylate	205
Magnetic properties of mixed ligand copper(II) thiocarboxylate	207
Magnetic properties of copper(II) complex with N-benzoyl-N'-<i>p</i>-hydroxythiobenzhydrazine	209
Magnetic properties of copper(II) complex with N-nicotinoyl-N'-thiobenzoyl hydrazine	211
Magnetic properties of copper(II) chloro complex with N-nicotinoyl-N'-thiobenzoyl hydrazine	213
Magnetic properties of copper(II) complex with 1,2-bis(4-pyridyl)ethane and dicynamide	215
Magnetic properties of copper(II) complex with 1,2-bis(4-pyridyl)ethane	218
Magnetic properties of copper(II) chloro complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-chlorophenol	221
Magnetic properties of copper(II) chloro, thiocyanate complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	223
Magnetic properties of copper(II) perchlorato with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	225
Magnetic properties of copper(II) sulphato with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	227
Magnetic properties of copper(II) nitrato with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	229

Magnetic properties of copper(II) acetato complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	231
Magnetic properties of copper(II) chloro complex with an azo dye derived from 4-aminoantipyrine and <i>p</i>-cresol	233
Magnetic properties of copper(II) complex with Mannich base, N-(morpholinobenzyl)-benzamide	235
Magnetic properties of copper(II) complex with a Schiff-base	237
Magnetic properties of salt consisting of ammonium-crown ether supramolecular cation with bis(maleonitriledithiolato)-cuprate	239
Magnetic properties of salt consisting of ammonium-benzo-crown ether supramolecular cation with bis-(maleonitriledithiolato)cuprate	242
Magnetic properties of copper(II) complex with bis(2-hydroxy-1-naphthaldehyde)malonoyldihydrazone	245
Magnetic properties of copper(II) complex with Schiff-base	247
Magnetic properties of copper(II) complex with a Schiff-base	249
Magnetic properties of copper(II) complex with tripod ligand tris[2-((2-pyridylmethyl)-amino)ethyl]amine	251
Magnetic properties of copper(II) complex with bis[2-((2-pyridylmethyl)-amino)ethyl]-amine	253
Magnetic properties of copper(II) complex with tetraazamacrocyclic and tetracyanoquinodimethane anion-radical ligands	255
Magnetic properties of copper(II) complex with tetraazamacrocyclic and tetracyanoquinodimethane anion-radical ligands	257
Magnetic properties of copper(II) complex with tetraazamacrocyclic and tetracyanoquinodimethane anion-radical ligands	259
Magnetic properties of copper(II) complex with N,N,N-tris(2-aminoethyl)-amine and tetracyanoquinodimethane anion-radical ligands	261
Magnetic properties of copper(II) complex with 1,4,7,10-tetraazadecane and tetracyanoquinodimethane anion-radical	263
Magnetic properties of copper(II) complex with name of N₄ tetraaza-macrocyclic and tetracyanoquinodimethane anion-radical ligands	265

Magnetic properties of copper(II) complex with tetraazamacrocyclic and tetracyanoquinodimethane anion-radical ligands	267
Magnetic properties of two-dimensional copper(II) complex with 1,3-propylenebis-(oxamate)	269
Magnetic properties of copper(II) complex with 6-aminouracil	271
Magnetic properties of mixed-ligand copper(II) with pyridine and <i>o</i>-H-(tetradecachlorotriphenyl)methane-4-carboxylic acid	273
Magnetic properties of helical, one-dimensional copper(II) coordination polymer based on <i>N</i>-(2-hydroxybenzyl)glycine	275
Magnetic properties of helical, one-dimensional copper(II) coordination polymer based on <i>N</i>-(2-hydroxybenzyl)-L-alanine	277
Magnetic properties of dicopper(II) complex with bis(2-imidazolyl)-bis-(methoxycarbonyl)methylmethane	279
Magnetic properties of dicopper(II) complex with malonic acid and 4,4'-bipyridine	281
Magnetic properties of copper(II) complex with a Schiff-base	284
Magnetic properties of copper(II) complex with 2,2'-dipyridylamine and monovalent tetraoxophosphate anion	286
Magnetic properties of copper(II) complex with ethylenediphosphonate	288
Magnetic properties of copper(II) complex with methylenediphosphonate	290
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and 3-methylisoquinoline	292
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and isoquinoline	294
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and 4-methylquinoline	296
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and quinoline	298
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and 3,5-dimethylpyridine	300
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and pyridine	302
Magnetic properties of mixed ligand copper(II) complex with salicylideneglycinate and urea	304

Magnetic properties of mixed ligand copper(II) complex with salicylidene glycinate and 2,4-dimethylpyridine	306
Magnetic properties of copper(II) complex with dinucleating ligand	308
Magnetic properties of alkoxy bridged binuclear copper(II) complex with thiocyanate and 2-(1',1'-dimethylamino)- 2-methyl-1-propanol ...	310
Magnetic properties of alkoxy bridged binuclear copper(II) complex with isocyanate and 2-(2-hydroxyethyl)pyridine	312
Magnetic properties of alkoxy bridged binuclear copper(II) complex with acetate and 6-methyl-2-pyridylmethanol	314
Magnetic properties of alkoxy bridged binuclear copper(II) complex with thiocyanate and 2-(hydroxymethyl)pyridine	316
Magnetic properties of alkoxy bridged binuclear copper(II) complex with thiocyanate and 2-(1',1'-dimethylamino)-propanol	318
Magnetic properties of μ-alkoxo, μ-pyrazolato bridged dicopper(II) complex with 1,3-bis-(4,6-dimethoxysalicylideneamino)-2-propanol ...	320
Magnetic properties of μ-alkoxo, μ-pyrazolato bridged dicopper(II) complex with 1,3-bis(3-fluorosalicylideneamino)-2-propanol	322
Magnetic properties of oxalate bridged binuclear copper(II) complex with 2,2'-dipyridylamine	324
Magnetic properties of mixed ligand copper(II) complex with nitronyl nitroxide and nitrate ligands	327
Magnetic properties of mixed ligand copper(II) complex with nitronyl nitroxide and acetate ligands	329
Magnetic properties of mixed ligand copper(II) complex with nitronyl nitroxide and dicyanamide ligands (as dihydrate)	331
Magnetic properties of mixed ligand copper(II) complex with nitronyl nitroxide and dicyanamide ligands (as triacetonitrilate)	333
Magnetic properties of mixed ligand copper(II) complex with nitronyl nitroxide and dicyanamide ligands	335
Magnetic properties of alkoxy bridged binuclear copper(II) complex with chloro and 2-(1',1'-dimethylamino)-2-methyl-1-propanol	338
Magnetic properties of phenolato bridged binuclear copper(II) complex with Schiff-base	340
Magnetic properties of oxalato bridged binuclear copper(II) complex with Schiff-base	342

Magnetic properties of hydroxo-bridged binuclear copper(II) aqua compound with 2,2'-dipyridylamine	345
Magnetic properties of one-dimensional bromo-bridged copper(II) compound with 1,4-diazacycloheptane	347
Magnetic properties of pseudohalide bridged polymeric copper(II) complex with 2,3-bis(2-pyridyl)pyrazine	350
Magnetic properties of polymeric tetrakis-(pyrazolyl)borate copper(II) complex	352
Magnetic properties of polymeric tetrakis-(pyrazolyl)borate copper(II) complex	354
Magnetic properties of polymeric tetrakis-(pyrazolyl)borate copper(II) complex	356
Magnetic properties of copper(II) 2-methylthionicotinate adduct with butylnicotinate	358
Magnetic properties of hydroxo-bridged binuclear copper(II) triflate compound with 2,2'-dipyridylamine	360
Magnetic properties of hydroxo-bridged Binuclear copper(II) bromo compound with 2,2'-dipyridylamine	362
Magnetic properties of binuclear copper(II) complex with 2-(2-pyridyl)imino-N-(2-thiazolin-2-yl)thiazolidine	364
Magnetic properties of ion-pair complex consisting of copper(II) cation with bis-(2-pyridylcarbonyl)amidate and copper(II) anion having bis(2-pyrimidylcarbonyl)amidate as well as sulphate ligands	367
Magnetic properties of triply bridged binuclear copper(II) complex containing 2,2'-dipyridylamine	369
Magnetic properties of triply bridged binuclear copper(II) complex containing 2,2'-dipyridylamine	372
Magnetic properties of doubly bridged binuclear copper(II) complex containing 2,2'-dipyridylamine	375
Magnetic properties of binuclear copper(II) complex with bis(2-pyrimidyl-carbonyl)amidate	377
Magnetic properties of binuclear copper(II) complex with azacyano-carbanion and pyridine ligand	380
Magnetic properties of croconato bridged binuclear copper(II) complex with bis-(2-pyrimidylcarbonyl)amidate	382

Magnetic properties of squarato bridged binuclear copper(II) complex with bis(2-pyrimidylcarbonyl)amidate	384
Magnetic properties of oxalate bridged binuclear copper(II) complex with bis-(2-pyrimidylcarbonyl)amidate	386
Magnetic properties of binuclear triplybridged copper(II) complex with di-2-pyridylamine	388
Magnetic properties of binuclear triplybridged copper(II) complex with di-2-pyridylamine	390
Magnetic properties of binuclear triplybridged copper(II) complex with di-2-pyridylamine	392
Magnetic properties of imidazolate bridged binuclear copper(II), Cu₂ complex with salicylidenevalinate	394
Magnetic properties of homodinuclear copper(II), Cu₂ complex with glutathione	396
Magnetic properties of homobimetallic Cu(II)-Cu(II) complex with bis(2-hydroxy-1-naphthaldehyde)malonoyldihydrazone	398
Magnetic properties of μ-oxalato bridged binuclear copper(II) complex with 1-(2-aminoethyl)pyrrolidine	400
Magnetic properties of μ-oxalato bridged binuclear copper(II) complex with 1-(2-aminoethyl)piperidine	403
Magnetic properties of μ-oxalato bridged binuclear copper(II) complex with 4-(2-aminoethyl)morpholine	406
Magnetic properties of copper(II) complex with 2-[2'-hydroxy-benzalidene-5'-(4''-phenyl, 2''-thiazolylazo)]benzoic acid	408
Magnetic properties of copper(II) complex with oximethiosemicarbazone	410
Magnetic properties of copper(II) complex with oxime-thiosemicarbazone	412
Magnetic properties of copper(II) complex with 2-[2'-hydroxy-benzalidene-5'-(4''-phenyl, 2''-thiazolylazo)]phenol	414
Magnetic properties of copper(II) 2-methylthionicotinate adduct with pyridine	416
Magnetic properties of copper(II) complex with 2-methylthionicotinate	418
Magnetic properties of dicopper(II) nitrato complex with thioether functionalized pyrazolate ligand	420

Magnetic properties of dicopper(II) perchlorate complex with thioether functionalized pyrazolate ligand	422
Magnetic properties of binuclear copper(II) complex with 1,4,7-trithia-cyclononane and TCNQ	424
Magnetic properties of dicopper(II) chloro complex with 5-(phenylazo)-2-thiohydantoin	426
Magnetic properties of mixed ligand copper(II) thiocarboxylate	428
Magnetic properties of chloro bridged binuclear copper(II) complex with Schiff-base	430
Magnetic properties of chloro bridged binuclear copper(II) complex with Schiff-base	432
Magnetic properties of phenoxo bridged binuclear copper(II) complex with Schiff-base	434
Magnetic properties of phenoxo bridged binuclear copper(II) complex with Schiff-base	436
Magnetic properties of binuclear copper(II) complex with N-(2-hydroxybenzyl)ethanolamine	438
Magnetic properties of copper(II) complex with phenanthroline and maleic acid	440
Magnetic properties of binuclear copper(II) complex with N,N'-disubstituted dithiooxamides derived from α-amino acid	442
Magnetic properties of binuclear copper(II) complex with N,N'-disubstituted dithiooxamide derived from α-amino acid	444
Magnetic properties of binuclear copper(II) complex with N,N'-disubstituted dithiooxamide derived from α-amino acid	446
Magnetic properties of binuclear copper(II) complex with N,N'-bis(1-carboxy-3-methylbutyl)dithiooxamide	448
Magnetic properties of μ-oxalato bridged binuclear copper(II) complex with 1,8-bis(2-pyridyl)-3,6-dithiaoctane	450
Magnetic properties of μ-imidazolato-bridged copper(II) complex with diethylenetriamine	452
Magnetic properties of μ-imidazolato-bridged copper(II) complex with diethylenetriamine	454
Magnetic properties of μ-imidazolato-bridged copper(II) complex with diethylenetriamine	456

Magnetic properties of copper(II) 2-methylthionicotinate adduct with ethylnicotinate	458
Magnetic properties of chloro-bridged binuclear copper(II) complex with 5-aminomethyl-3-methylpyrazole	460
Magnetic properties of dimeric copper(II) 2-methylthionicotinate adduct with dimethylsulfoxide	462
Magnetic properties of dimeric copper(II) 2-methylthionicotinate adduct with dimethylformamide	464
Magnetic properties of μ-hydroxo bridged binuclear copper(II) complex with 2-aminopyrimidine	467
Magnetic properties of dicopper(II) complex with 2-nitrobenzoic acid	469
Magnetic properties of dicopper(II) chloro complex with N-salicylidene-<i>n</i>-myristoylhydrazine	471
Magnetic properties of dicopper(II) chloro complex with N-salicylidene-<i>n</i>-alkanoylhydrazine	473
Magnetic properties of neutral dicopper(II) complex with N-salicylidene-<i>n</i>-ethanoylhydrazine	475
Magnetic properties of neutral dicopper(II) chloro complex with N-salicylidene-<i>n</i>-pentanoylhydrazine	477
Magnetic properties of neutral dicopper(II) complex with N-salicylidene-<i>n</i>-hexanoylhydrazine	479
Magnetic properties of neutral dicopper(II) complex with N-salicylidene-<i>n</i>-heptanoylhydrazine	481
Magnetic properties of neutral dicopper(II) complex with N-salicylidene-<i>n</i>-myristoylhydrazine	483
Magnetic properties of neutral dicopper(II) chloro complex with N-salicylidene-<i>n</i>-alkanoylhydrazine	485
Magnetic properties of copper chloro complex with 2-(dimethylamino-methyl)-3-hydroxypyridine	487
Magnetic properties of copper bromo complex with 2-(dimethylamino-methyl)-3-hydroxypyridine	489
Magnetic properties of NCO bridged dimeric copper(II) complex with phenanthroline	492
Magnetic properties of μ-1,1,3-azido bridged dicopper(II) complex with 1-(2-aminoethyl)pyrrolidine	494

Magnetic properties of binuclear carbonato-bridged Cu(II) complex with 2,2'-dipyridylamine	497
Magnetic properties of binuclear carbonato-bridged Cu(II) complex with 2,2'-dipyridylamine	499
Magnetic properties of mixed ligand binuclear copper(II) complex with 2,2'-bipyrimidine and malonic acid	501
Magnetic properties of dicopper(II) complex with 2-[(4-methyl-pyridin- 2-ylimino)-methyl]phenol	503
Magnetic properties of dicopper(II) chloro complex with unsymmetrical binucleating ligand	505
Magnetic properties of dicopper(II) bromo complex with unsymmetrical binucleating ligand	507
Magnetic properties of dimeric-copper(II) complex with 2-furoic hydrazide	509
Magnetic properties of binuclear copper(II) complex with macrocyclic ligand	512
Magnetic properties of oxamido-bridged binuclear copper(II) complex with imino nitroxide radical	514
Magnetic properties of oxamido-bridged binuclear copper(II) complex with nitrosyl nitroxide radical	517
Magnetic properties of dicopper(II) acetate complex with unsymmetrical binucleating ligand	519
Magnetic properties of binuclear copper(II) complex with 1,4-dinitro-2,3,5,6-tetracarboxylatobenzoic acid	521
Magnetic properties of μ-oxalato bridged binuclear copper(II) complex with 2,2'-dipyridylamine	523
Magnetic properties of tetracarboxylato bridged dimer, copper(II) formato dimethylformamide	525
Magnetic properties of μ-cyanodicopper(II,II) furyl linked azacryptlate	527
Magnetic properties of an imidazole-bridged dicopper complex of a macrocyclic ligand with two hydroxyethyl pendants	529
Magnetic properties of μ-cyanodicopper(II,II) <i>para</i>-linked azacryptlate	532
Magnetic properties of μ-cyanodicopper(II,II) <i>meta</i>-linked azacryptlate	534

Magnetic properties of copper(II) complex of tetraaldehyde phenylhydrazone	536
Magnetic properties of binuclear 2,3,5, 6-tetrakis(2-pyridyl)pyrazine containing copper(II) complex	538
Magnetic properties of mixed-valence μ-cyanodicopper(I,II) cryptlate	541
Magnetic properties of 3D arrays of binuclear copper(II) complex with 3, 5-pyrazoledicarboxylic acid	544
Magnetic properties of binuclear copper(II) complex with 3, 5-pyrazoledicarboxylic acid	546
Magnetic properties of mixed-ligand binuclear copper(II) complex with (tetradecachloro-4-carboxytriphenyl)-methyl radical	548
Magnetic properties of oxalato bridged binuclear copper(II) complex with NNO donor Schiff-base ligand	551
Magnetic properties of oxalato bridged binuclear copper(II) complex with NNO donor Schiff-base ligand	553
Magnetic properties of oxamidato bridged binuclear copper(II) complex with NNO donor Schiff-base ligand	556
Magnetic properties of endogenously bridged binuclear copper(II) complex formed by 1,3-bis[1,4,7-triazacyclonon- 1-yl]-propan-2-ol	558
Magnetic properties of binuclear μ_2-1,1-N_3 copper(II) complex with tridentate Schiff-base	560
Magnetic properties of binuclear μ_2-1,1-N_3 copper(II) complex with tridentate Schiff-base	563
Magnetic properties of binuclear μ_2-1,1-N_3 copper(II) complex with tridentate Schiff-base	566
Magnetic properties of binuclear copper(II) di-2-pyridylamine complex with dual end-to-end and end-on cyanate bridges	569
Magnetic properties of a dual μ-[N(CN)$_2$] bridged copper(II) supramolecule with binuclear unit	572
Magnetic properties of copper(II) complex with macrocyclic multidentate ligand and 4,4'-bipyridine	575
Magnetic properties of μ-phenylphosphinato bridged binuclear copper(II) complex with bipyridine	578

Magnetic properties of μ-acetate bridged binuclear Cu(II) complex with <i>N, N, N', N'</i>-bis-[(2-hydroxybenzyl)(<i>N</i>-methylimidazolyl)]-2-ol-1,3-propanediamine	580
Magnetic properties of copper(II) complex with picolinic-N-oxide built from Cu₂O₂ dimer units	582
Magnetic properties of copper(II) binuclear complex with μ-hydroxo and μ-pyrazolato bridges	585
Magnetic properties of binuclear Cu(II)-Cu(II) complex containing macrocyclic oxamide and phenanthroline ligands	587
Magnetic properties of heterometallic copper(II)-potassium(I) complex with macrocyclic ligand	590
Magnetic properties of alkoxo bridged bimetallic, copper(II)-Al(III) complex with Schiff-base	592
Magnetic properties of alkoxo bridged bimetallic, Cu(II)-Al(III) complex with Schiff-base	594
Magnetic properties of binuclear Cu(II)-Mn(II) complex containing macrocyclic oxamide and bipyridine ligands	596
Magnetic properties of heteronuclear (Cu-Co), thiocyanate bridged complex with macrocyclic ligand	599
Magnetic properties of heteronuclear (Cu-Co), thiocyanate bridged complex with macrocyclic ligand	602
Magnetic properties of oxo-bridged hetero-binuclear, Cu(II)-Co(II) complex with compartmental Schiff-base	605
Magnetic properties of binuclear Cu(II)-Ni(II) complex containing macrocyclic oxamide and phenanthroline ligands	607
Magnetic properties of imidazolate bridged heterodinuclear copper(II)-nickel(II), Cu(II)-Ni(II), complex with salicylidenevalinate	610
Magnetic properties of ion-pair (salt) complex containing bis(ethylenediamine)- copper(II) cation and tetracyanopalladate(II) anion	612
Magnetic properties of heterodinuclear copper(II)-zinc (II) complex with glutathione	614
Magnetic properties of heterobimetallic Zn(II)-Cu(II) complex with bis(2-hydroxy- 1-naphthaldehyde)malonyldihydrazone	616

Magnetic properties of imidazolate bridged heterodinuclear copper(II)-zinc(II), Cu(II)-Zn(II), complex with salicylidenevalinate	618
Magnetic properties of two-dimensional copper(II) complex with mellitic acid	620
Magnetic properties of 3D arrays of trinuclear copper(II) complex with 3, 5-pyrazoledicarboxylic acid	622
Magnetic properties of trinuclear copper(II) complex containing 2,3,5,6-tetrakis(2-pyridyl)pyrazine	624
Magnetic properties of μ_2-acetato bridged trinuclear copper(II) complex with 2,6-bis-(pyridine-2-carboxamido)pyridine	627
Magnetic properties of trinuclear copper(II) complex with tris-cyclam based on 1,3, 5-trimethylenebenzene	630
Magnetic properties of pyrazolato-bridged trinuclear copper(II) complex with methanol	632
Magnetic properties of pyrazolato-bridged trinuclear copper(II) complex with N,N-dimethylethylenediamine	634
Magnetic properties of tricopper(II) azide complex containing diethylenetriamine as co-ligand	636
Magnetic properties of trinuclear copper(II) complex with malic acid	639
Magnetic properties of μ_3-OMe bridged tricopper(II) chloro complex with substituted pyrazole	642
Magnetic properties of μ_3-OMe bridged tricopper(II) bromo complex with substituted pyrazole	645
Magnetic properties of trinuclear copper(II) complex with tris-cyclam based on phloroglucinol	648
Magnetic properties of trinuclear cyanamidonitrate-copper(II) complex bridged by imidazolato ligand	650
Magnetic properties of trinuclear copper(II) complex with semi-coordinated iodides formed through ligand sharing	653
Magnetic properties of linear trinuclear copper(II) complex bridged by sulfosalicylate ligand	656
Magnetic properties of trimeric copper(II) complex simultaneously containing salicylato mono and dianions	659