

Contents

Volume 5A

Introduction	1
Part I La	13
Magnetic properties of lanthanide-manganese oxide	15
Part II Ti	17
Magnetic properties of polychelate of titanium(III) with Schiff-base 4,4'-bis[(N-phenylsalicylaldimine-5)azo]biphenyl	19
Part III V	21
Magnetic properties of oxovanadium(IV) complex with salicylaldehyde-2-pyrazinoyl hydrazone	23
Magnetic properties of non-oxo vanadium(IV) complex with aminebis(phenolate) ligand	25
Magnetic properties of non-oxo vanadium(IV) complex with aminebis(phenolate) ligand	27
Magnetic properties of oxovanadium(IV) complex with hydrazone of isonicotinic acid hydrazide	29
Magnetic properties of oxovanadium(II) complex with bis(<i>S</i> -methyldithio-carbazate)-5-acetyl-2,4-dihydroxyacetophenone ...	31
Magnetic properties of oxovanadium(IV) complex with N,N'-bis(4-hydroxy-salicylidene)-1,2-diaminoethane	33
Magnetic properties of oxovanadium(IV) complex with N,N'-bis(4-hydroxysalicylidene)-1,3-diaminopropane	35
Magnetic properties of oxovanadium(IV) complex with N,N'-(4-hydroxy-salicylidene)-1,4-diaminobutane	37

vii

Magnetic properties of oxalato-bridged oxovanadium(IV) dimer using L-ascorbic acid as oxalate precursor	39
Magnetic properties of oxalato-malonato complex of oxovanadium(IV) using L-ascorbic acid as oxalate precursor	42
Magnetic properties of dinuclear oxovanadium(IV) complex with salicylaldehyde-2-pyrazinoyl hydrazone	45
Magnetic properties of dinuclear oxovanadium(IV) complex with bis (salicylaldehyde)-2,3-pyrazinoyldihydrazone	47
Magnetic properties of dimeric oxovanadium(II) complex with tetrathio-carbamate of <i>o</i>-phenylenediamine	49
Magnetic properties of dimeric oxovanadium(II) complex with tetrathiocarbamate of <i>m</i>-phenylenediamine	51
Magnetic properties of dimeric oxovanadium(II) complex with tetrathiocarbamate of <i>p</i>-phenylenediamine	53
Magnetic properties of polychelate of oxovanadium(II) with Schiff-base 4,4'-bis[(N-phenylsalicylaldimine-5)azo]biphenyl	55
Magnetic properties of chiral mixed-valence vanadium oxide	57
Part IV Cr	61
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base, N,N'-(2-hydroxy)propylenebis-[(2-imino-3-oximino)butane] and pyridine	63
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base N,N'-(2-hydroxy)propylenebis-(acetylacetonimine) and pyridine	65
Magnetic properties of mixed ligand chromium(III) complex involving bidentate Schiff-base N,N'-propylenebis(benzaldimine) and pyridine	67
Magnetic properties of mixed ligand chromium(III) complex involving bidentate Schiff-base N,N'-propylenebis(benzaldimine) and bipyridine	69
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base, N,N'-(2-hydroxy)propylenebis-[(2-imino-3-oximino)butane] and ammonia	71
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base N,N'-(2-hydroxy)propylenebis-(acetylacetonimine) and ammonia	73

Magnetic properties of mixed ligand chromium(III) complex involving bidentate Schiff-base, N,N'-propylenebis(benzaldimine) and ammonia	75
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base, N,N'-(2-hydroxy)propylenebis-[(2-imino-3-oximino)butane] and thiocyanate	77
Magnetic properties of mixed ligand complex of chromium(III) involving quadridentate Schiff-base, N,N'-(2-hydroxy)propylenebis-(acetylacetoneimine) and thiocyanate	79
Magnetic properties of mixed ligand chromium(III) complex involving bidentate Schiff-base N,N'-propylenebis(benzaldimine) and thiocyanate	81
Magnetic properties of chromium(III) complex with hydrazone of isonicotinic acid hydrazide	83
Magnetic properties of polychelate of chromium(III) with Schiff-base 4,4'-bis[(N-phenylsalicylaldehyde-5-azo)biphenyl	85
Magnetic properties of chromium(III) nitrate complex with 14-membered tetraazamacrocyclic	87
Magnetic properties of chromium(III) nitrate complex with 16-membered tetraazamacrocyclic	89
Magnetic properties of mixed ligand chromium(III) complex with alanine and 2-acetylpyridine	91
Magnetic properties of heterometallic Mg-Cr₂ complex with nitrilotri-acetate ligand	93
Magnetic properties of heterometallic Ca-Cr₂ complex with nitrilotri-acetate ligand	95
Magnetic properties of heterometallic Sr-Cr₂ complex with nitrilotriacetate ligand	98
Magnetic properties of heterometallic Ba-Cr₂ complex with nitrilotri-acetate ligand	100
Magnetic properties of oxalato-bridged hetero binuclear Cu(II)-Cr(III) complex with 2,2'-bipyridine and 3-[N-2-(pyridylethyl)formimidoyl]salicylic acid	102
Magnetic properties of oxalato-bridged hetero binuclear Mn(II)-Cr(III) complex with 1,10-phenanthroline and azide ligands ...	105

Part V Mo	107
Magnetic properties of molybdenum cluster cyanide complex with selenide and bromide ligands in the cluster core	109
Magnetic properties of molybdenum cluster cyanide complex with selenide and chloride ligands in the cluster core	111
Part VI Mn	113
Magnetic properties of mixed ligand complex of manganese(II) with oxydiacetate and 2,2'-bipyridine	115
Magnetic properties of mixed ligand complex of manganese(II) with oxydiacetate and 2,2':6,2''-terpyridine	117
Magnetic properties of mixed ligand manganese(II) complex with acetylacetone and bipyridine	119
Magnetic properties of manganese(II) complex with tripodal aminopyridyl chelator	121
Magnetic properties of manganese(II) complex with tripodal aminopyridyl chelator	124
Magnetic properties of manganese(II) complex with 2-trithiocarbonatoethyl xanthate	127
Magnetic properties of manganese(II) complex with polystyrene-anchored Schiff-base	129
Magnetic properties of manganese(II) complex with bis(<i>S</i> -methyldithio-carbazate)-5-acetyl-2,4-dihydroxyacetophenone	131
Magnetic properties of mixed ligand complex of manganese(II) with 2-pyridinealdoxime and benzimidazole	133
Magnetic properties of mixed ligand complex of manganese(II) with 2-pyridinealdoxime and benzotriazole	135
Magnetic properties of mixed ligand complex of manganese(II) with 2-pyridinealdoxime and triazole	137
Magnetic properties of mixed ligand complex of manganese(II) with 2-pyridinealdoxime and carbazole	139
Magnetic properties of manganese(II) complex with quinazoline ligand	141
Magnetic properties of manganese(II) complex with 2-pyridine-2-yl-3-(pyridine-2-carboxylideneamino)quinazolin-4(3H)-one	143

Magnetic properties of manganese(II) complex with 2-thiophene-2-yl-3-(thiophene-2-carboxylideneamino)-1,2-dihydroquinazolin-4(3H)-one	145
Magnetic properties of manganese(II) complex with N-(2-hydroxy-1-naphthalidene)phenylglycine Schiff-base	147
Magnetic properties of manganese(II) complex with a tetradentate NOOO- donor Schiff-base ligand	149
Magnetic properties of three dimensional coordination polymer having diaquamanganese(II) complex cation and octacyanomolybdate complex anion	151
Magnetic properties of dinuclear manganese(II) complex with phenyl-cyanamide and phenanthroline ligands	153
Magnetic properties of thiocyanato-bridged copper(II)-manganese(II) bimetallic coordination polymer	156
Magnetic properties of trinuclear dipicolinato manganese complex ...	159
Magnetic properties of mixed valent oxo-centered trinuclear manganese cluster	162
Magnetic properties of trinuclear manganese(II) benzoate complex with phenyl (2-pyridyl) ketone	165
Magnetic properties of neutral cubane cluster of manganese(II) with acetate, dicyanamide and di-2-pyridyl-hemiacetal ligands	167
Magnetic properties of oxamido-bridged tetranuclear Cu₃Mn complex with substituted diene and thiocyanate	170
Magnetic properties of oxamido-bridged tetranuclear Cu₃Mn complex with substituted diene and dicyanamide	174
Magnetic properties of three dimensional hydroxide sulfate of manganese(II)	177
Magnetic properties of octanuclear mixed valent manganese benzoate cluster with phenyl(2-pyridyl)oxime ligand	180
Magnetic properties of coordination polymer of manganese(II) maleate / α,α'-diimine system	183
Magnetic properties of coordination polymer of manganese(II) maleate / α,α'-diimine system	186
Magnetic properties of polynuclear manganese(II) complex with phenyl-cyanamide and 4-benzoylpyridine	188

Magnetic properties of polymeric manganese(II) complex with phenyl-cyanamide and 4,4'-bipyridine	191
Magnetic properties of manganese(II) complex with oxydiacetate ligand	193
Magnetic properties of polymeric mixed ligand complex of manganese(II) with oxydiacetate and 1,10-phenanthroline	196
Magnetic properties of manganese(III) complex with methyl salicylimidate	199
Magnetic properties of manganese(III) complex with hydrazone of isonicotinic acid hydrazide	201
Magnetic properties of manganese(III) complex with salicylidine-4, 4-dimethyl-3-thiosemicarbazide	203
Magnetic properties of manganese(III) complex with salicylidine-morpholine-N-thiohydrazide	205
Magnetic properties of manganese(III) complex with salicylidine-piperidine-N-thiohydrazide	207
Magnetic properties of manganese(III) chloro complex with aza-crown macrocyclic ligand	209
Magnetic properties of manganese(III) chloro complex with aza-crown macrocyclic ligand	212
Magnetic properties of manganese(III) chloro complex with aza-crown macrocyclic ligand	215
Magnetic properties of manganese(III) chloro complex with morpholino Schiff-base	218
Magnetic properties of manganese(III) chloro complex with morpholino Schiff-base	220
Magnetic properties of manganese(III) chloro complex with morpholino Schiff-base	222
Magnetic properties of manganese(III) chloro complex with benzo-10-aza-crown ether macrocyclic ligand	224
Magnetic properties of manganese(III) chloro complex with morpholino Schiff-base	227
Magnetic properties of dinuclear manganese(III) complex with ethyl salicylimidate	229

Magnetic properties of coordination complex having manganese(II) cation with 1,4,7-trimethyl-1,4,7-triazacyclononane and manganese(III) anion with aminebis(phenolate)	232
Magnetic properties of polychelate of manganese(III) with Schiff-base 4,4'-bis[(N-phenylsalicylaldimine-5)azo]biphenyl	235
Part VII Re	237
Magnetic properties of heterometallic rhenium(IV)-copper(II) complex with macrocyclic ligand	239
Magnetic properties of heterometallic rhenium(IV)-copper(II) complex with macrocyclic ligand	242
Magnetic properties of phenylimido rhenium(VI) complex with substituted amide	246
Magnetic properties of phenylimido rhenium(VI) complex with substituted amide	248
Magnetic properties of phenylimido rhenium(VI) complex with substituted amide	250
Part VIII Fe	253
Magnetic properties of iron(II) quinoline-2-carboxylate complex	255
Magnetic properties of iron(II) quinoline-2-carboxylate complex	257
Magnetic properties of iron(II) polymer containing 1,2,4,5-tetrakis-(diphenylphosphanyl)benzene ligand	259
Magnetic properties of iron(II) formate	261
Magnetic properties of iron(II) phosphite	263
Magnetic properties of iron(II) complex with dianionic [N₄]macrocyclic ligand	266
Magnetic properties of methanol adduct of iron(II) complex with dianionic [N₄]macrocyclic ligand	268
Magnetic properties of iron(II) complex with dianionic [N₄]macrocyclic ligand	270
Magnetic properties of iron(II) complex with tripodal aminopyridyl chelator	273
Magnetic properties of iron(II) complex with tripodal aminopyridyl chelator	275

Magnetic properties of 1,4-diazabicyclo[2.2.2]octane bridges adduct of dinuclear iron(II) complex with dianionic [N₄]macrocyclic ligand . .	278
Magnetic properties of 1,4-diazabicyclo[2.2.2]octane bridged adduct of dinuclear iron(II) complex with dianionic [N₄]-macrocyclic ligand	280
Magnetic properties of 1,4-diazabicyclo[2.2.2]octane bridged adduct of dinuclear iron(II) complex with dianionic [N₄]-macrocyclic ligand	283
Magnetic properties of iron(II) complex with bis(<i>S</i>-methyldithiocarbazate)-5-acetyl-2,4-dihydroxyacetophenone	286
Magnetic properties of polymeric iron(II) complex with 4,4'-bipyridine and bridging dichromate anion	288
Magnetic properties of iron(III) complex with 1,5-bis(<i>N</i>-methylaceto-hydroxamic acid) derivative	291
Magnetic properties of mixed ligand iron(III) complex with octamethyltetraphenyl-prophyrin and dimethylaminopyridine	293
Magnetic properties of mixed ligand iron(III) complex with octaethyltetraphenyl-prophyrin and pyridine	296
Magnetic properties of mixed ligand iron(III) complex with octamethyltetraphenyl-prophyrin and pyridine	299
Magnetic properties of mixed ligand iron(III) complex with octaethyltetraphenyl-prophyrin and dimethylaminopropane	301
Magnetic properties of hexakis(thiocyanato)ferrate(III) ion	304
Magnetic properties of iron(III) chloro complex with dianionic [N₄]-macrocyclic ligand	307
Magnetic properties of iron(III) iodo complex with dianionic [N₄]-macrocyclic ligand	310
Magnetic properties of iron(III) nitrosyl complex with dianionic [N₄]-macrocyclic ligand	312
Magnetic properties of iron(III) nitrite complex with dianionic [N₄]-macrocyclic ligand	314
Magnetic properties of iron(III) chloro complex with dianionic [N₄]-macrocyclic ligand	317
Magnetic properties of iron(III) complex with hydrazine of isonicotinic acid hydrazide	319
Magnetic properties of iron(III) complex with polystyrene-anchored Schiff-base	321

Magnetic properties of iron(III) complex with 2-trithiocarbonatoethyl xanthate	323
Magnetic properties of iron(III) nitrate complex with 14-membered tetraazamacrocyclic	325
Magnetic properties of iron(III) nitrate complex with 16-membered tetraazamacrocyclic	327
Magnetic properties of iron(III) chloro complex with N,N'-bis(4-hydroxy-salicylidene)-1,2-diaminoethane	329
Magnetic properties of iron(III) chloro complex with N,N'-bis(4-hydroxy-salicylidene)-1,3-diaminopropane	331
Magnetic properties of iron(III) chloro complex with N,N'-(4-hydroxy-salicylidene)-1,4-diaminobutane	333
Magnetic properties of iron(III) complex with 12-membered tetraimine macrocyclic ligand	335
Magnetic properties of mixed ligand iron(III) complex with alanine and 2-acetylpyridine	337
Magnetic properties of μ-oxo-bridged dinuclear Fe(III) complex with picolinamide	339
Magnetic properties of oximate-bridged heterodinuclear iron(III)-iron(II) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclononane	342
Magnetic properties of oximate-bridged heterodinuclear iron(III)-manganese(II) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclononane	345
Magnetic properties of oximate-bridged heterodinuclear iron(II)-cobalt(III) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclononane	348
Magnetic properties of oximate-bridged heterodinuclear iron(III)-nickel(II) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclononane	351
Magnetic properties of ion-pair coordination complex having iron(III) Schiff-base complex cation and nickel(III) thiolate anion	354
Magnetic properties of oximate-bridged heterodinuclear iron(III)-copper(II) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclononane	356

Magnetic properties of oximate-bridged heterodinuclear iron(III)-zinc(II) complex with pyridine-2-aldoxime and 1,4,7-trimethyl-1,4,7-triazacyclo- nonane	359
Magnetic properties of ion-pair coordination complex having iron(III) Schiff-base complex cation and palladium(III) thiolate anion	362
Magnetic properties of ion-pair coordination complex having iron(III) Schiff-base complex cation and platinum(III) thiolate anion	364
Magnetic properties of 2,4,6-tris(4-carboxyphenylimino-4-formylphenoxy)-1,3,5-triazine bridged trinuclear iron(III) saloph Schiff-base complex	366
Magnetic properties of 2,4,6-tris(4-carboxyphenylimino-4-formylphenoxy)-1,3,5-triazine bridged trinuclear iron(III) salen Schiff-base complex	369
Magnetic properties of 2,4,6-tris(2,5-dicarboxyphenylimino-4-formyl-phenoxy)-1,3,5-triazine bridged trinuclear iron(III) saloph Schiff-base complex	372
Magnetic properties of 2,4,6-tris(2,5-dicarboxyphenylimino-4-formyl-phenoxy)-1,3,5-triazine bridged trinuclear iron(III) salen Schiff-base complex	375
Magnetic properties of cyano bridged bimetallic copper(II)-iron(III) complex	378
Magnetic properties of cyano bridged bimetallic copper(II)-iron(III) complex	380
Magnetic properties of μ-azido-bridged heterobimetallic Fe(III)₂-Cu(II) complex with picolinamide	382
Magnetic properties of polychelate of iron(III) with Schiff-base 4,4'-bis[(N-phenylsalicylalimine-5)azo]biphenyl	385
Part IX Ru	387
Magnetic properties of mononuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, adenosine	389
Magnetic properties of mononuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, cytidine	391
Magnetic properties of ruthenium(III) complex with 2-(2'-hydroxyphenyl)-benzimidazole	393

Magnetic properties of ruthenium(III) complex with 2-(2'-mercaptphenyl)-benzimidazole	395
Magnetic properties of ruthenium(III) complex with 2-(2'-hydroxynaphthyl)-benzimidazole	397
Magnetic properties of ruthenium(III) chloro complex with N,N'-bis(4-hydroxysalicylidene)-1,2-diaminoethane	399
Magnetic properties of ruthenium(III) chloro complex with N,N'-bis(4-hydroxysalicylidene)-1,3-diaminopropane	401
Magnetic properties of ruthenium(III) chloro complex with N,N'-(4-hydroxysalicylidene)-1,4-diaminobutane	403
Magnetic properties of dinuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, adenine	405
Magnetic properties of dinuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, guanosine	407
Magnetic properties of dinuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, guanine	409
Magnetic properties of dinuclear ruthenium(III) complex with α-alanine and nucleic acid constituent, cytosine	411
Magnetic properties of aqua chlorodiruthenium(II, III) complex with 3-phenoxypropionic acid	413
Magnetic properties of bromodiruthenium(II, III) complex with 3-phenoxypropionic acid	416
Magnetic properties of iododiruthenium (II, III) complex with 3-phenoxypropionic acid	418
Magnetic properties of aquaiododiruthenium(II, III) complex with 3-phenoxypropionic acid	420
Magnetic properties of aquachlorodiruthenium(II, III) complex with 2,2-diphenylpropionic acid	423
Magnetic properties of bromodiruthenium(II, III) complex with 2,2-diphenylpropionic acid	426
Magnetic properties of aquabromodiruthenium(II, III) complex with 2,2-diphenylpropionic acid	428
Magnetic properties of iododiruthenium(II, III) complex with 2,2-diphenylpropionic acid	430
Magnetic properties of aquaiododiruthenium(II, III) complex with 2,2-diphenylpropionic acid	432

Magnetic properties of chlorodiruthenium(II, III) complex with 3-phenoxypropionic acid	434
Magnetic properties of mixed valence trinuclear complex of ruthenium with trifluoroperazine	436
Magnetic properties of mixed valence trinuclear complex of ruthenium with prochlorperazine	438
Magnetic properties of mixed valence trinuclear complex of ruthenium with butaperazine	440
Magnetic properties of mixed valence trinuclear complex of ruthenium with methoxypromazine	442
Part X Os	445
Magnetic properties of mixed ligand osmium(III) complex with 2,2-bipyridine and 2,2'-dipyridylamine	447
Magnetic properties of mixed ligand osmium(III) complex with 8-hydroxy-quinoline and 2,2'-dipyridylamine	449
Magnetic properties of mixed ligand osmium(III) complex with 2-methyl-8-hydroxyquinoline and 2,2'-dipyridylamine	451
Magnetic properties of mixed ligand osmium(IV) complex with 8-hydroxy-quinoline and 2,2'-dipyridylamine	453
Magnetic properties of mixed ligand osmium(IV) complex with 2-methyl-8-hydroxyquinoline and 2,2'-dipyridylamine	455
Part XI Co	457
Magnetic properties of cobalt(II) chloro complex with 4-(phenyl)thiazolyl-hydrazone of <i>o</i>-anisaldehyde	459
Magnetic properties of cobalt(II) bromo complex with 4-(phenyl)thiazolyl-hydrazone of <i>o</i>-anisaldehyde	461
Magnetic properties of cobalt(II) dinitrato complex with 4-(phenyl)-thiazolylhydrazone of <i>o</i>-anisaldehyde	463
Magnetic properties of cobalt(II) chloro complex with 4-(<i>p</i>-bromophenyl)-thiazolylhydrazone of <i>o</i>-anisaldehyde	465
Magnetic properties of cobalt(II) bromo complex with 4-(<i>p</i>-bromophenyl)-thiazolylhydrazone of <i>o</i>-anisaldehyde	467
Magnetic properties of cobalt(II) dinitrato complex with 4-(<i>p</i>-bromophenyl)thiazolylhydrazone of <i>o</i>-anisaldehyde	469

Magnetic properties of cobalt(II) chloro complex with macrocyclic ligand derived from 2,3-butanedione and its hydrazide	471
Magnetic properties of cobalt(II) bromo complex with macrocyclic ligand derived from 2,3-butanedione and its hydrazide	473
Magnetic properties of cobalt(II) nitrato complex with macrocyclic ligand derived from 2,3-butanedione and its hydrazide	475
Magnetic properties of cobalt(II) thiocyanato complex with macrocyclic derived ligand from 2,3-butanedione and its hydrazide	477
Magnetic properties of three dimensional coordination polymer having diaquacobalt(II) complex cation and octacyanomolybdate complex anion	479
Magnetic properties of cobalt(II) complex with 2-pyridine-2-yl-3-(pyridine-2-carboxylideneamino)quinazolin-4(3H)-one	481
Magnetic properties of cobalt(II) complex with salicylaldehyde thiazolyl hydrazone	483
Magnetic properties of cobalt(II) complex with salicylaldehyde thiazolyl hydrazone	486
Magnetic properties of cobalt(II) complex with 2-thiophene-2-yl-3(thiophene-2-carboxylideneamino)-1,2-dihydroquinazolin-4(3H)-one ..	489
Magnetic properties of cobalt(II) complex with propranolol dithiocarbamate	491
Magnetic properties of cobalt(II) complex with morpholino Schiff-base	493
Magnetic properties of cobalt(II) chloro complex with benzo-10-aza-crown ether macrocyclic ligand	496
Magnetic properties of cobalt(II) complex with N-(2-hydroxy-1-naphthalidene)phenylglycine Schiff-base	499
Magnetic properties of cobalt(II) complex N-(ethyl-4-amino-1-piperidine carboxylate)glyoxime	501
Magnetic properties of cobalt(II) complex with N-(ethyl-4-amino-1-piperidine carboxylate)phenylglyoxime	503
Magnetic properties of cobalt(II) complex with <i>vic</i>-dioxime ligand, N,N-bis(2-[(2,2-dimethyl-1,3-dioxolan-4-yl)methyl]amino}butyl) dihydroxy-ethanediimidamide	505

Magnetic properties of cobalt(II) complex with polystyrene-anchored Schiff-base	507
Magnetic properties of cobalt(II) complex with bis(<i>S</i>-methyldithio-carbazate)-5-acetyl-2,4-dihydroxyacetophenone	509
Magnetic properties of mixed ligand complex of cobalt(II) with 2-pyridinealdoxime and carbazole	511
Magnetic properties of mixed ligand complex of cobalt(II) with 2-pyridinealdoxime and triazole	513
Magnetic properties of mixed ligand complex of cobalt(II) with 2-pyridinealdoxime and benzotriazole	515
Magnetic properties of mixed ligand cobalt(II) complex with 2-pyridinealdoxime and benzimidazole	517
Magnetic properties of cobalt(II) complex with heterocyclic N₄O₂ aldimine	519
Magnetic properties of cobalt(II) complex with 2-trithiocarbonatoethyl xanthate	521
Magnetic properties of cobalt(II) nitrate complex with 16-membered tetraazamacrocyclic	523
Magnetic properties of cobalt(II) nitrate complex with 14-membered tetraazamacrocyclic	525
Magnetic properties of cobalt(II) Schiff-base complex with N-[2-thienyl-methylidene]-2-hydroxy-1-aminoethane as bidentate ligand	527
Magnetic properties of cobalt(II) Schiff-base complex with N-[2-thienyl-methylidene]-2-hydroxy-1-aminoethane as tridentate ligand	529
Magnetic properties of cobalt(II) quinoline-2-carboxylate complex ...	531
Magnetic properties of cobalt(II) bis(dicyanobenzenedithiolate) complex	533
Magnetic properties of cobalt(II) complex with <i>p</i>-<i>tert</i>-butylcalix[4]arene having phosphinoyl pendant arms	535
Magnetic properties of cobalt(II) phosphite	537
Magnetic properties of mononuclear cobalt(II) complex with phthalic acid and 2,2'-bipyridine	540
Magnetic properties of cobalt(II) thiocyanato complex with N,N,N',N'-tetraethylpyridine-2,6-dithiocarboxamide	543

Magnetic properties of cobalt(II) bromo complex with N,N,N'-N'-tetraethylpyridine-2,6-dithiocarboxamide	546
Magnetic properties of cobalt(II) iodo complex with N,N,N'-N'-tetraethylpyridine-2,6-dithiocarboxamide	549
Magnetic properties of cobalt(II) complex with N,N'-bis(ethyl-4-amino-1-piperidine carboxylate)glyoxime	552
Magnetic properties of cobalt(II) complex with acesulfamate	554
Magnetic properties of cobalt(II) complex with azo-linked Schiff-base	556
Magnetic properties of cobalt(II) complex with azo-linked Schiff-base	558
Magnetic properties of cobalt(II) complex with N,N'-bis(4-hydroxysalicylidene)-1,2-diaminoethane	560
Magnetic properties of cobalt(II) complex with N,N'-bis(4-hydroxysalicylidene)-1,3-diaminopropane	562
Magnetic properties of cobalt(II) complex with N,N'-(4-hydroxysalicylidene)-1,4-diaminobutane	564
Magnetic properties of cobalt(II) chloro complex with macrocyclic ligand derived from thiocarbohydrazide and benzil	566
Magnetic properties of cobalt(II) acetato complex with macrocyclic ligand derived from thiocarbohydrazide and benzil	568
Magnetic properties of cobalt(II) nitrate complex with macrocyclic ligand derived from thiocarbohydrazide and benzil	570
Magnetic properties of cobalt(II) complex with a tetradentate NOOO-donor Schiffbase ligand	572
Magnetic properties of cobalt(II) complex with mono-Schiff-base aza-crown macrocyclic pendant	574
Magnetic properties of cobalt(II) complex with mono-Schiff-base aza-crown macrocyclic pendant	577
Magnetic properties of cobalt(II) complex with mono-Schiff-base aza-crown macrocyclic pendant	580
Magnetic properties of cobalt(II) complex with morpholino macrocyclic ligand	583
Magnetic properties of cobalt(II) complex with morpholino macrocyclic ligand	585

Magnetic properties of cobalt(II) complex with morpholino macrocyclic ligand	587
Magnetic properties of mixed ligand cobalt(II) complex with alanine and 2-acetylpyridine	589
Magnetic properties of cobalt(II) complex with 12-membered tetraimine macrocyclic ligand	591
Magnetic properties of cobalt(II) complex with quinazoline ligand	593
Magnetic properties of five membered cobaltocycle with C,P-chelating ligand	595
Magnetic properties of dimeric cobalt(II) complex with tetrathiocarbamate of <i>p</i>-phenylenediamine	597
Magnetic properties of dimeric cobalt(II) complex with tetrathiocarbamate of <i>m</i>-phenylenediamine	599
Magnetic properties of dimeric cobalt(II) complex with tetrathiocarbamate of <i>o</i>-phenylenediamine	601
Magnetic properties of dinuclear cobalt(II) complex with bis-bidentate azo dye	603
Magnetic properties of dinuclear cobalt(II) complex with bis-tridentate azo dye	605
Magnetic properties of homo-bimetallic cobalt(II) complex with 4-isobutyl-3,5-di(2-pyridyl)-4H-1,2,4-triazole	607
Magnetic properties of dinuclear cobalt(II) complex with phthalic acid and 2,2'-bipyridine	609
Magnetic properties of dinuclear cobalt(II) complex with 3-methylsulfanylpropionate and thioacetate ligands	612
Magnetic properties of phenoxo and chloro bridged binuclear cobalt(II) complex with Schiff-base derived from 2,6-diformyl-4-methylphenol and 2-hydroxy-3-hydrazinoquinoxiline	614
Magnetic properties of phenoxo and chloro bridged binuclear cobalt(II) complex with Schiff-base derived from 2,6-diformyl-4-methylphenol and 4-amino-3-antipyrine	616
Magnetic properties of trinuclear cobalt cluster having 1,2,4,5-benzene-tetracarboxylate ligand	619
Magnetic properties of neutral cubane cluster of cobalt(II) with acetate, dicyanamide and di-2-pyridyl-hemiacetal ligands	622

Magnetic properties of tetranuclear cobalt(II) complex with ditopic ligand	624
Magnetic properties of polymeric cobalt(II) complex with 4,4'-bipyridine and bridging dichromate anion	627
Magnetic properties of cobalt(II) complex with benzene-1,2,4,5-tetracarbo-1,2:4,5-dihydrazide and 1,10-phenanthroline	630
Magnetic properties of cobalt(II) complex with benzene-1,2,4,5-tetracarbo-1,2:4,5-dihydrazide	633
Magnetic properties of coordination polymer of cobalt(II) with malic acid	635
Magnetic properties of polynuclear cobalt(II) complex with phthalic acid and 2,2'-bipyridine	638
Magnetic properties of dinuclear cobalt(III) complex with salicylidinepiperidine-N-thiohydrazide	641
Magnetic properties of dinuclear cobalt(III) complex with salicylidinemorpholine-N-thiohydrazide	643
Magnetic properties of dinuclear cobalt(III) complex with salicylidine-4,4-dimethyl-3-thiosemicarbazide	645
Magnetic properties of polynuclear cobalt cluster with N-(2-hydroxybenzyl)ethanolamine	647