

# Contents

## Volume 7A

|                                                                                                                                                              |           |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>Introduction</b> .....                                                                                                                                    | <b>1</b>  |
| <b>Part I V</b> .....                                                                                                                                        | <b>13</b> |
| <b>Magnetic properties of sulfato-bridged dinuclear vanadium(III) complex with 2-oxo-N,N'-bis(2-pyridylmethyl)1,3-propanediamine-N,N'-diacetate</b> .....    | <b>15</b> |
| <b>Magnetic properties of sulfato-bridged tetranuclear vanadium(III) complex with 2-oxo-N,N'-bis(2-pyridylmethyl)1,3-propanediamine-N,N'-diacetate</b> ..... | <b>17</b> |
| <b>Magnetic properties of oxovanadium(IV) complex with 1,4-phenyldiphosphonic acid as piperazinium salt</b> .....                                            | <b>19</b> |
| <b>Magnetic properties of fluoro oxovanadium(IV) complex with 1,4-phenyl-diphosphonic acid as 1,4-diazabicyclo- [2.2.2]octanium salt</b> .....               | <b>21</b> |
| <b>Magnetic properties of oxovanadium(IV) complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                                               | <b>23</b> |
| <b>Magnetic properties of vanadium(IV) complex with 2,3-dihydroxypyridine</b> .....                                                                          | <b>25</b> |
| <b>Magnetic properties of oxovanadium(IV) complex with 14-membered tetraazamacrocyclic ligand</b> .....                                                      | <b>27</b> |
| <b>Magnetic properties of oxoxvanadium(IV) complex with 16-membered tetraazamacrocyclic ligand</b> .....                                                     | <b>29</b> |
| <b>Magnetic properties of oxovanadium(IV) complex with 14-membered tetraazamacrocyclic ligand</b> .....                                                      | <b>31</b> |
| <b>Magnetic properties of oxovanadium(IV) complex with 14-membered tetraazamacrocyclic ligand</b> .....                                                      | <b>33</b> |

|                                                                                                                                                       |    |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| <b>Magnetic properties of vanadium(IV) complex with 2-pyridineformamide thiosemicarbazone</b> .....                                                   | 35 |
| <b>Magnetic properties of vanadium(IV) complex with 2-pyridineformamide thiosemicarbazone</b> .....                                                   | 37 |
| <b>Magnetic properties of vanadium(IV) complex with 2-pyridineformamide thiosemicarbazone</b> .....                                                   | 39 |
| <b>Magnetic properties of vanadium(IV) complex with 2-pyridineformamide thiosemicarbazone</b> .....                                                   | 41 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from 2,6-diacetylpyridine and semicarbazide</b> .....                      | 43 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and ethanol amine</b> .....  | 45 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and ethanol amine</b> .....  | 47 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and ethanol amine</b> .....  | 49 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and propanol amine</b> ..... | 51 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and propanol amine</b> ..... | 53 |
| <b>Magnetic properties of oxovanadium(IV) complex with Schiff-base derived from heterocyclic <math>\beta</math>-diketone and propanol amine</b> ..... | 55 |
| <b>Magnetic properties of dinuclear vanadium(IV) complex with 24-membered macrocycle</b> .....                                                        | 57 |
| <b>Magnetic properties of dinuclear vanadium(IV) complex with 24-membered macrocycle</b> .....                                                        | 60 |
| <b>Magnetic properties of dinuclear oxovanadium(IV) complex with Schiff-base derived from 2,6-diacetylpyridine and thiosemicarbazide</b> .....        | 63 |
| <b>Magnetic properties of germanium-polyoxovanadate cluster</b> .....                                                                                 | 65 |
| <b>Magnetic properties of polymeric chain assemblies of oxovanadium(IV) with bispyrazolone</b> .....                                                  | 67 |

|                                                                                                                                                                                           |            |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Part II Cr .....</b>                                                                                                                                                                   | <b>69</b>  |
| <b>Magnetic properties of chormium(III) complex with ethylene cross-bridged tetraazamacrocyclic ligand (cyclam) .....</b>                                                                 | <b>71</b>  |
| <b>Magnetic properties of chromium(III) complex with benzimidazole thiocarbamate .....</b>                                                                                                | <b>73</b>  |
| <b>Magnetic properties of chromium(III) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol .....</b>                                                           | <b>75</b>  |
| <b>Magnetic properties of chromium(III) complex with 3-methyl-5-oxo-N,1-diphenyl-4,5-dihydro-1-H-pyrazol-4-carbothioamide .....</b>                                                       | <b>77</b>  |
| <b>Magnetic properties of homo-bimetallic Cr<sub>2</sub>(II) complex with [N<sub>10</sub>] macrocyclic ligand modified with tetramide function .....</b>                                  | <b>79</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-bimetallic Cr-Ni complex with bis(2-pyridylmethyl)-N,N'-diethylenetriamine ....</b>                                       | <b>82</b>  |
| <b>Magnetic properties of heterobimetallic Cu(II)-Cr(III) ion-pair complex having ammonia, ethylenediamine and thiocyanate ligands .....</b>                                              | <b>84</b>  |
| <b>Magnetic properties of heterobimetallic Cu(II)-Cr(III) complex having ammonia, ethylenediamine and thiocyanate ligand .....</b>                                                        | <b>86</b>  |
| <b>Magnetic properties of heterobimetallic Cu(II)-Cr(III) ion-pair complex having ammonia, ethylenediamine and thiocyanate ligand ...</b>                                                 | <b>88</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-trimetallic Cr-Ni<sub>2</sub> complex with tris(2-aminoethyl)-N,N,N'-ethyl diamine .....</b>                              | <b>90</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-trimetallic Cr-Ni<sub>2</sub> complex with tetraethylenepentamine .....</b>                                               | <b>93</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-trimetallic Cr-Ni<sub>2</sub> complex with bis(2-pyridylmethyl)-N,N'-diethylenetriamine as BF<sub>4</sub> salt .....</b>  | <b>95</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-trimetallic Cr-Ni<sub>2</sub> complex with bis(2-pyridylmethyl)-N,N'-diethylenetriamine as ClO<sub>4</sub> salt .....</b> | <b>98</b>  |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hetero-tetrametallic Cr-Ni<sub>3</sub> complex with bis(2-pyridylmethyl)-N,N'-dipropylenetriamine .....</b>                      | <b>100</b> |
| <b>Magnetic properties of <math>\mu</math>-cyano-bridged hexametallic Cr-Ni<sub>5</sub> complex with tetraethylenepentamine .....</b>                                                     | <b>102</b> |
| <b>Magnetic properties of heptametallic Cr-Ni<sub>6</sub> complex with tetraethylenepentamine .....</b>                                                                                   | <b>104</b> |

|                                                                                                                                                                |            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of heptametallic Cr-Ni<sub>6</sub> complex with tetraethylenepentamine</b> .....                                                        | <b>106</b> |
| <b>Part III Mo</b> .....                                                                                                                                       | <b>109</b> |
| <b>Magnetic properties of ion-pair complex having cubane type Mo<sub>3</sub>CoS<sub>4</sub> cluster cation and maleonitriledithiolato nickelate anion</b> .... | <b>111</b> |
| <b>Part IV Mn</b> .....                                                                                                                                        | <b>113</b> |
| <b>Magnetic properties of tetrachloromanganate(II) anion as bdt-ttp based cation radical salt</b> .....                                                        | <b>115</b> |
| <b>Magnetic properties of manganese(II) complex with chiral nitronyl nitroxide radical</b> .....                                                               | <b>117</b> |
| <b>Magnetic properties of manganese(II) complex with photochromic phenanthroline ligand</b> .....                                                              | <b>120</b> |
| <b>Magnetic properties of manganese(II) complex with pyrophosphate as ammonium salt</b> .....                                                                  | <b>122</b> |
| <b>Magnetic properties of manganese(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                                | <b>124</b> |
| <b>Magnetic properties of manganese(II) polymeric complex with N,N'-bis(carboxymethyl)dithiooxamide</b> .....                                                  | <b>126</b> |
| <b>Magnetic properties of manganese(II) complex with benzimidazole thiocarbamate</b> .....                                                                     | <b>129</b> |
| <b>Magnetic properties of mixed ligand manganese(II) complex with 2-mercapto-quinazolin-4-one and 1,10-phenanthroline</b> .....                                | <b>131</b> |
| <b>Magnetic properties of mixed ligand manganese(II) complex with 2-mercapto-quinazolin-4-one and ethylenediamine</b> .....                                    | <b>133</b> |
| <b>Magnetic properties of mixed ligand manganese(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and 1,10-phenanthroline</b> .....                       | <b>135</b> |
| <b>Magnetic properties of mixed ligand manganese(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and ethylenediamine</b> .....                           | <b>137</b> |
| <b>Magnetic properties of manganese(II) complex with 2,6-pyridinedicarboxylic acid derivative</b> .....                                                        | <b>139</b> |
| <b>Magnetic properties of manganese(II) complex with 1-formylpiperazinyl-dithiocarbamate</b> .....                                                             | <b>141</b> |
| <b>Magnetic properties of manganese(II) complex with Schiff-base derived from 4-aminoantipyrine and 2-aminobenzoic acid</b> .....                              | <b>143</b> |

|                                                                                                                                    |     |
|------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of manganese(II) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol</b> .....    | 145 |
| <b>Magnetic properties of manganese(II) complex with 22-membered tetramide macrocyclic ligand</b> .....                            | 147 |
| <b>Magnetic properties of manganese(II) complex with 24-membered tetramide macrocyclic ligand</b> .....                            | 149 |
| <b>Magnetic properties of manganese(II) complex with sulphonylurea glibenclamide</b> .....                                         | 151 |
| <b>Magnetic properties of dinuclear chloroacetato bridged manganese(II) complex with 1,10-phenanthroline</b> .....                 | 153 |
| <b>Magnetic properties of cyano-bridged manganese(II)-molybdenum(IV) complex</b> .....                                             | 155 |
| <b>Magnetic properties of heterobimetallic Cu-Mn complex with ethylenediamine and chloro ligands (as dmsolvate)</b> .....          | 158 |
| <b>Magnetic properties of heterobimetallic Cu-Mn complex with ethylenediamine and chloro ligands (as dmfsolvate)</b> .....         | 160 |
| <b>Magnetic properties of heterobimetallic Cu-Mn complex with 1,10-phenanthroline and chloro ligands (as dmfsolvate)</b> .....     | 162 |
| <b>Magnetic properties of heterobimetallic Cu-Mn complex with 1,10-phenanthroline and chloro ligands (as hydratesolvate)</b> ..... | 164 |
| <b>Magnetic properties of heterobimetallic Mn(II)-Ag(I) complex with 4-methylpyridine</b> .....                                    | 166 |
| <b>Magnetic properties of heterobimetallic Mn(II)-Ag(I) complex with 3-methylpyridine</b> .....                                    | 168 |
| <b>Magnetic properties of heterobimetallic Mn(II)-Ag(I) coordination polymer having pyridine-4-aldoxime</b> .....                  | 170 |
| <b>Magnetic properties of heterobimetallic Mn(II)-Ag(I) coordination polymer having 4-cyanopyridine ligand</b> .....               | 172 |
| <b>Magnetic properties of trinuclear trithiocyanurato-bridged manganese(II) involving 1,10-phenanthroline</b> .....                | 175 |
| <b>Magnetic properties of trinuclear trithiocyanurato-bridged manganese(II) involving 4,4'-dimethyl-2,2'-bipyridine</b> .....      | 178 |
| <b>Magnetic properties of trinuclear trithiocyanurato bridged manganese(II) involving 2,2'-bipyridine</b> .....                    | 181 |
| <b>Magnetic properties of trinuclear chloroacetato bridged manganese(II) complex with 2,2'-bipyridine</b> .....                    | 183 |

|                                                                                                                                                                                               |     |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of cyano-bridged hetero-trimetallic Mn<sub>2</sub>Nb complex with imidazole</b> .....                                                                                  | 186 |
| <b>Magnetic properties of hexametallic manganese cluster with 1-naphthoic acid and 2-hydroxyphenylpropanone oxime</b> .....                                                                   | 188 |
| <b>Magnetic properties of hexametallic manganese cluster with anthracene-9-carboxylic acid and 2-hydroxyphenylpropanone oxime</b> .....                                                       | 190 |
| <b>Magnetic properties of hexametallic manganese cluster with anthracene-4-ethynylbenzoic acid and 2-hydroxyphenylpropanone oxime</b> .....                                                   | 192 |
| <b>Magnetic properties of homometallic heptanuclear mixed-valent manganese (Mn<sub>4</sub><sup>III</sup>Mn<sub>3</sub><sup>II</sup>) complex with Schiff-base</b> .....                       | 195 |
| <b>Magnetic properties of hybridized complex of Mn<sub>2</sub><sup>III</sup> single molecule magnet and nickel thiolate complex</b> .....                                                     | 197 |
| <b>Magnetic properties of polymeric chain assemblies of manganese(II) with bis-pyrazolone</b> .....                                                                                           | 200 |
| <b>Magnetic properties of polynuclear chloroacetato bridged manganese(II) complex with 1,10-phenanthroline</b> .....                                                                          | 202 |
| <b>Magnetic properties of acetato bridged dinuclear manganese(III, III) complex with ligand having three phenolate groups</b> .....                                                           | 204 |
| <b>Magnetic properties of acetato bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> .....           | 207 |
| <b>Magnetic properties of diphenylglycolate bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> ..... | 210 |
| <b>Magnetic properties of triphenylacetate bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> .....  | 213 |
| <b>Magnetic properties of benzoato bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> .....          | 216 |
| <b>Magnetic properties of chloroacetato bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> .....     | 219 |
| <b>Magnetic properties of propionate bridged tetranuclear complex with [Fe<sub>2</sub>Mn<sub>2</sub>]<sup>8+</sup> core having triazacyclononane and salicylaldoxime ligands</b> .....        | 222 |

|                                                                                                                                                                                             |            |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of diphenylglycolate bridged tetranuclear complex with <math>[\text{Mn}_4\text{O}_2]^{8+}</math> core having triazacyclononane and salicylaldoxime ligands</b> ..... | 225        |
| <b>Magnetic properties of triphenylacetate bridged tetranuclear complex with <math>[\text{Mn}_4\text{O}_2]^{8+}</math> core having triazacyclononane and salicylaldoxime ligands</b> .....  | 228        |
| <b>Part V Re</b> .....                                                                                                                                                                      | <b>231</b> |
| <b>Magnetic properties of hexa(isothiocyanato)rhenate(IV)</b> .....                                                                                                                         | 233        |
| <b>Part VI Fe</b> .....                                                                                                                                                                     | <b>235</b> |
| <b>Magnetic properties of iron(II) complex with thioether ligand</b> .....                                                                                                                  | 237        |
| <b>Magnetic properties of iron(II) complex with pyrazolylbis{(thio)methyl}-borate</b> .....                                                                                                 | 239        |
| <b>Magnetic properties of iron(II) complex with 2-ethyl-5-methylimidazol-4-carbaldehyde azine</b> .....                                                                                     | 241        |
| <b>Magnetic properties of iron(II) complex with polynitrile ligand and 4,4'-bipyridine as co-ligand</b> .....                                                                               | 243        |
| <b>Magnetic properties of iron(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                                                                  | 245        |
| <b>Magnetic properties of iron(II) complex with 3,3-bis(1-methylimidazol-2-yl)propionate</b> .....                                                                                          | 247        |
| <b>Magnetic properties of mixed ligand of iron(II) complex with Schiff-base type ligand and di(4-picolyl)amine</b> .....                                                                    | 249        |
| <b>Magnetic properties of mixed ligand iron(II) complex with Schiff-base type ligand and 4,4'-bipyridine</b> .....                                                                          | 251        |
| <b>Magnetic properties of mixed ligand iron(II) complex with Schiff-base type ligand and <i>trans</i>-1,2-bis(4-pyridyl)ethylene</b> .....                                                  | 253        |
| <b>Magnetic properties of iron(II) complex with 2,6-pyridinedicarboxylic acid derivative</b> .....                                                                                          | 255        |
| <b>Magnetic properties of iron(II) complex with 1-formylpiperazinyldithiocarbamate</b> .....                                                                                                | 257        |
| <b>Magnetic properties of iron(II) complex with 4-(2-thiazolylazo)resorcinol</b> .....                                                                                                      | 259        |
| <b>Magnetic properties of iron(II) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol</b> .....                                                                  | 261        |
| <b>Magnetic properties of iron(II) complex with bis(pyrazolyl)pyridine based ligand (perchlorate salt)</b> .....                                                                            | 263        |

---

|                                                                                                                                    |            |
|------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of iron(II) complex with bis(pyrazolyl)pyridine based ligand (tetraphenylborate salt) .....</b>             | <b>266</b> |
| <b>Magnetic properties of dimeric iron(II) complex with Schiff-base, salophen .....</b>                                            | <b>269</b> |
| <b>Magnetic properties of dinuclear iron(II) complex with 2-ethyl-5-methylimidazol-4-carbaldehyde azine (plate crystals) .....</b> | <b>272</b> |
| <b>Magnetic properties of mixed ligand dinuclear iron(II) complex with Schiff-base type ligand and 1-methylimidazole .....</b>     | <b>274</b> |
| <b>Magnetic properties of mixed ligand iron(II) complex with Schiff-base type ligand and 4-dimethylaminopyridine .....</b>         | <b>277</b> |
| <b>Magnetic properties of heterobimetallic Fe(II)-Cu(II) salt of cyanoimino-methanedithiolate ligand .....</b>                     | <b>279</b> |
| <b>Magnetic properties of benzothiazole-2-thiolato-bridged heterobimetallic Fe(II)-Cu(II) complex .....</b>                        | <b>281</b> |
| <b>Magnetic properties of azido complex of nonplanar iron(III) porphyrin .....</b>                                                 | <b>283</b> |
| <b>Magnetic properties of octaethylporphyrin iron(III) complex containing phenyl cyanamide .....</b>                               | <b>285</b> |
| <b>Magnetic properties of octaethylporphyrin iron(III) complex containing 2-chlorophenylcyanamide .....</b>                        | <b>287</b> |
| <b>Magnetic properties of octaethylporphyrin iron(III) complex containing 2-methylphenylcyanamide .....</b>                        | <b>290</b> |
| <b>Magnetic properties of octaethylporphyrin iron(III) complex containing 2-4-dimethylphenylcyanamide .....</b>                    | <b>293</b> |
| <b>Magnetic properties of iron(III) complex with pyrophosphite .....</b>                                                           | <b>295</b> |
| <b>Magnetic properties of iron(III) complex with 2,6-diacetylpyridine-bis(thiosemicarbazone) .....</b>                             | <b>297</b> |
| <b>Magnetic properties of iron(III) complex with 2-acetylphenylthiosemi-carbazone-6-acetylpyridine .....</b>                       | <b>300</b> |
| <b>Magnetic properties of iron(III) complex with dipodal tridentate chelating agent .....</b>                                      | <b>302</b> |
| <b>Magnetic properties of iron(III) complex with dipodal tridentate chelating agent .....</b>                                      | <b>304</b> |
| <b>Magnetic properties of iron(III) complex with Schiff-base derived from 4-aminoantipyrine and 2-aminobenzoic acid .....</b>      | <b>306</b> |

|                                                                                                                                         |            |
|-----------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of iron(III) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol</b>                   | <b>308</b> |
| <b>Magnetic properties of iron(III) complex with 3-methyl-5-oxo-N,1-diphenyl-4,5-dihydro-1-H-pyrazol-4-carbothioamide</b>               | <b>310</b> |
| <b>Magnetic properties of iron(III) complex with sulphonylurea glibenclamide</b>                                                        | <b>312</b> |
| <b>Magnetic properties of binuclear iron(III) complex with N-(1-ethanol)-N,N-bis(3-<i>tert</i>-butyl-5-methyl-2-hydroxybenzyl)amine</b> | <b>314</b> |
| <b>Magnetic properties of binuclear iron(III) hydroxo complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> | <b>317</b> |
| <b>Magnetic properties of binuclear iron(III) chloro complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b>  | <b>320</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>323</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>325</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>327</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>329</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>331</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>333</b> |
| <b>Magnetic properties of <math>\mu</math>-oxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>           | <b>335</b> |
| <b>Magnetic properties of <math>\mu</math>-hydroxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>       | <b>337</b> |
| <b>Magnetic properties of <math>\mu</math>-hydroxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>       | <b>339</b> |
| <b>Magnetic properties of <math>\mu</math>-hydroxo-biscarboxylato dinuclear iron(III) complex with hydrotris(pyrazolyl)borate</b>       | <b>341</b> |
| <b>Magnetic properties of trinuclear oxo-centered Fe(III) complex with 1,4-di(2'-N-methylbenzimidazolyl)aminophthalazine</b>            | <b>343</b> |

|                                                                                                                                                                                                                                                         |     |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of heterobimetallic Fe<sub>2</sub>Mn coordination polymer based on tetracyanometallic building blocks</b> .....                                                                                                                  | 346 |
| <b>Magnetic properties of hetero-trimetallic Fe<sup>III</sup><sub>2</sub>-Ni<sup>II</sup> oxalate complex as <i>trans</i>-4-(4-dimethylaminostyryl)-1-methylpyridinium salt</b> .....                                                                   | 348 |
| <b>Magnetic properties of heterobimetallic Fe<sub>2</sub>Cu coordination polymer based on tetracyanometallic building blocks</b> .....                                                                                                                  | 350 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and acetate as terminal ligand</b> .....                 | 352 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and monochloro acetate as terminal ligand</b> ....       | 354 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and dichloro acetate as terminal ligand</b> .....        | 357 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and trifluoro acetate</b> .....                          | 360 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and pentafluoro propionate as terminal ligand</b> .....  | 363 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and benzoate as terminal ligand</b> .....                | 366 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and <i>p</i>-nitro benzoate as terminal ligand</b> ..... | 369 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and pentafluoro benzoate as terminal ligand</b> .....    | 372 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and diethylphosphate as terminal ligand</b> .....        | 375 |
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and diphenylphosphate as terminal ligand</b> ....        | 378 |

|                                                                                                                                                                                                                                                   |            |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of mixed ligand tetrairon(III) complex of di-<math>\mu</math>-oxo-bis[di-<math>\mu</math>-phenolatodiiron(III)] core with phenol based macrocyclic compartmental ligand and diethyl dithiophosphate as terminal ligand</b> | <b>381</b> |
| <b>Magnetic properties of tetrairon(III) complex with dipivaloylmethane acid and 2-hydroxymethyl-1,3-propanediol</b>                                                                                                                              | <b>384</b> |
| <b>Magnetic properties of diphenylglycolate bridged tetranuclear complex with <math>[\text{Fe}_4\text{O}_2]^{8+}</math> core having triazacyclononane and salicylaldoxime ligands</b>                                                             | <b>386</b> |
| <b>Magnetic properties of triphenylacetate bridged tetranuclear complex with <math>[\text{Fe}_4\text{O}_2]^{8+}</math> core having triazacyclononane and salicylaldoxime ligands</b>                                                              | <b>389</b> |
| <b>Part VII Ru</b>                                                                                                                                                                                                                                | <b>393</b> |
| <b>Magnetic properties of bifunctional material derived from polyoxo-molybdate and ruthenium mononitrosyl complex</b>                                                                                                                             | <b>395</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with 8-hydroxyquinoline and norborna-2, 5-diene</b>                                                                                                                                 | <b>397</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with 8-hydroxyquinoline and 4-picoline</b>                                                                                                                                          | <b>399</b> |
| <b>Magnetic properties of ruthenium(III) complex with 2-[(4-N,N'-dimethyl-aminophenylimino)-methyl]-4-chlorophenol</b>                                                                                                                            | <b>401</b> |
| <b>Magnetic properties of ruthenium(III) complex with 2-[(4-N,N'-dimethyl-aminophenylimino)-methyl]-4-bromophenol</b>                                                                                                                             | <b>403</b> |
| <b>Magnetic properties of ruthenium(III) complex with 2-[(4-N,N'-dimethyl-aminophenylimino)-methyl]-4-iodophenol</b>                                                                                                                              | <b>405</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with chloro, triphenylphosphine and Schiff-base</b>                                                                                                                                 | <b>407</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with bromo, tetraphenylphosphine and Schiff-base</b>                                                                                                                                | <b>409</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with chloro, triphenylarsine and Schiff-base</b>                                                                                                                                    | <b>411</b> |
| <b>Magnetic properties of mixed ligand ruthenium(III) complex with bromo, triphenylarsine and Schiff-base</b>                                                                                                                                     | <b>413</b> |
| <b>Part VIII Co</b>                                                                                                                                                                                                                               | <b>415</b> |
| <b>Magnetic properties of cobalt bromide complex supported by <i>m</i>-terphenyl isocyanide</b>                                                                                                                                                   | <b>417</b> |

|                                                                                                                                                 |     |
|-------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of cobalt iodide complex supported by <i>m</i>-terphenyl isocyanide</b> .....                                            | 420 |
| <b>Magnetic properties of cobalt chloride complex supported by <i>m</i>-terphenyl isocyanide</b> .....                                          | 423 |
| <b>Magnetic properties of cobalt(II) complex with pyrazolylbis{(thio)methyl}borate</b> .....                                                    | 426 |
| <b>Magnetic properties of cobalt(II) chloro complex with pyrazolylbis[(thio)<i>t</i>-butyl]borate</b> .....                                     | 428 |
| <b>Magnetic properties of mixed ligand cobalt(II) complex with pyrazolylbis[(thio)-<i>t</i>-butyl]borate</b> .....                              | 430 |
| <b>Magnetic properties of tetrachlorocobaltate(II) anion as bdt-ttp based cation radical salt</b> .....                                         | 432 |
| <b>Magnetic properties of cobalt(II) complex with photochromic phenanthroline ligand</b> .....                                                  | 434 |
| <b>Magnetic properties of cobalt(II) complex with polynitrile ligand and 4,4'-bipyridine as co-ligand</b> .....                                 | 437 |
| <b>Magnetic properties of cobalt(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 439 |
| <b>Magnetic properties of nanocavity zeolite-Y (host) encapsulated cobalt(II) complex with unsaturated 16-membered macrocyclic ligand</b> ..... | 441 |
| <b>Magnetic properties of cobalt(II) complex with 3-aminoflavone</b> .....                                                                      | 444 |
| <b>Magnetic properties of cobalt(II) complex with pyrophosphate as ammonium salt</b> .....                                                      | 446 |
| <b>Magnetic properties of cobalt(II) complex with diethyl(pyridin-4-ylmethyl)phosphate</b> .....                                                | 448 |
| <b>Magnetic properties of cobalt(II) complex with 2,6-diacetylpyridine-bis(thiosemicarbazone)</b> .....                                         | 450 |
| <b>Magnetic properties of cobalt(II) complex with (1R, 2R)-1,2-bis-(5-amino-1,3,4-thiadiazol-2-yl)ethane-1,2-diol</b> .....                     | 452 |
| <b>Magnetic properties of cobalt(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> .....                               | 454 |
| <b>Magnetic properties of cobalt(II) nitrate complex with hydrotris(3-<i>tert</i>-butyl-5-isopropylpyrazol-1-yl)borate</b> .....                | 457 |
| <b>Magnetic properties of cobalt(II) hydroxo complex with hydrotris(3-<i>tert</i>-butyl-5-isopropylpyrazol-1-yl)borate</b> .....                | 459 |

|                                                                                                                                       |     |
|---------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of dinuclear cobalt(II) complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                   | 462 |
| <b>Magnetic properties of cobalt(II) complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                             | 465 |
| <b>Magnetic properties of cobalt(II) complex with N,N'-bis(2-pyridinecarboxaldimine)-1,8-diaminonaphthalene</b> .....                 | 467 |
| <b>Magnetic properties of cobalt(II) bromo complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino) acetohydrazide</b> .....    | 469 |
| <b>Magnetic properties of cobalt(II) nitrate complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino) acetohydrazide</b> .....  | 471 |
| <b>Magnetic properties of cobalt(II) acetate complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino) acetohydrazide</b> .....  | 473 |
| <b>Magnetic properties of polymeric chain assemblies of cobalt(II) with bis-pyrazolone</b> .....                                      | 475 |
| <b>Magnetic properties of cobalt(II) chloro complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....                 | 477 |
| <b>Magnetic properties of cobalt(II) nitrate complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....                | 479 |
| <b>Magnetic properties of cobalt(II) complex with benzimidazole thiocarbamate</b> .....                                               | 481 |
| <b>Magnetic properties of mixed ligand cobalt(II) complex with 2-mercapto-quinazolin-4-one and 1,10-phenanthroline</b> .....          | 483 |
| <b>Magnetic properties of mixed ligand cobalt(II) complex with 2-mercapto-quinazolin-4-one and ethylenediamine</b> .....              | 485 |
| <b>Magnetic properties of mixed ligand cobalt(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and 1,10-phenanthroline</b> ..... | 487 |
| <b>Magnetic properties of mixed ligand cobalt(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and ethylenediamine</b> .....     | 489 |
| <b>Magnetic properties of cobalt(II) complex with 2,6-pyridinedicarboxylic acid derivative</b> .....                                  | 491 |
| <b>Magnetic properties of cobalt(II) complex with 1-formylpiperazinyldithiocarbamate</b> .....                                        | 493 |
| <b>Magnetic properties of cobalt(II) complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> .....          | 495 |

|                                                                                                                                                                                |     |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of cobalt(II) complex with Schiff-base derived from 4-aminoantipyrine and 2-aminobenzoic acid</b> .....                                                 | 497 |
| <b>Magnetic properties of cobalt(II) complex with 14-membered tetraaza-macrocyclic Schiff-base ligand</b> .....                                                                | 499 |
| <b>Magnetic properties of cobalt(II) complex with 16-membered tetraaza-macrocyclic Schiff-base ligand</b> .....                                                                | 502 |
| <b>Magnetic properties of cobalt(II) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol</b> .....                                                   | 504 |
| <b>Magnetic properties of cobalt(II) complex with Schiff-base derived from 2,6-diacetylpyridine and semicarbazide</b> .....                                                    | 506 |
| <b>Magnetic properties of cobalt(II) complex with 22-membered tetramide macrocyclic ligand</b> .....                                                                           | 508 |
| <b>Magnetic properties of cobalt(II) complex with 24-membered tetramide macrocyclic ligand</b> .....                                                                           | 510 |
| <b>Magnetic properties of cobalt(II) complex with sulphonylurea glibenclamide</b> .....                                                                                        | 512 |
| <b>Magnetic properties of cobalt(II) chloro complex with 14-membered octaazamacrocyclic ligand</b> .....                                                                       | 514 |
| <b>Magnetic properties of cobalt(II) nitrate complex with 14-membered octaazamacrocyclic ligand</b> .....                                                                      | 517 |
| <b>Magnetic properties of cobalt(II) acetate complex with 14-membered octaazamacrocyclic ligand</b> .....                                                                      | 520 |
| <b>Magnetic properties of cobalt(II) complex with <i>o</i>-aminothiophenolate</b> .....                                                                                        | 523 |
| <b>Magnetic properties of cobalt(II) complex with <i>o</i>-iminothiobenzosemi-quinonato <math>\pi</math>-radical ligand</b> .....                                              | 526 |
| <b>Magnetic properties of <math>\mu</math>-carbonato bridged dinuclear cobalt(II) hydroxo complex with hydrotris(3-<i>tert</i>-butyl-5-isopropylpyrazol-1-yl) borate</b> ..... | 529 |
| <b>Magnetic properties of cobalt(II) complex with N,N'-bis[(3,4-dichloro-phenyl)methylidene]cyclohexan-1,4-diamine</b> .....                                                   | 531 |
| <b>Magnetic properties of homo-bimetallic Co(II)<sub>2</sub> complex with [N<sub>10</sub>] macrocyclic ligand modified with tetramide function</b> .....                       | 533 |
| <b>Magnetic properties of binuclear cobalt(II) complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> .....                                         | 536 |

|                                                                                                                                                                                           |            |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of dinuclear cobalt(II) complex with 3-methyl-5-oxo-N,1-diphenyl-4,5-dihydro-1-H-pyrazol-4-carbothioamide</b> . . . . .                                            | <b>538</b> |
| <b>Magnetic properties of heterobimetallic Co(II)-Ni(II) salt of cyanoimino-methanedithiolate ligand</b> . . . . .                                                                        | <b>540</b> |
| <b>Magnetic properties of benzothiazole-2-thiolato-bridged heterobimetallic Co(II)-Ni(II) complex</b> . . . . .                                                                           | <b>542</b> |
| <b>Magnetic properties of heterobimetallic Co(II)-Cu(II) salt of cyanoimino-methanedithiolate ligand</b> . . . . .                                                                        | <b>544</b> |
| <b>Magnetic properties of linear chloro-bridged trinuclear cluster, CoU<sub>2</sub> having 3,5-dimethylpyrazole and cyclam ligands</b> . . . . .                                          | <b>546</b> |
| <b>Magnetic properties of 1D coordination polymer of cobalt(II) with <math>\pi</math>-radical ligand</b> . . . . .                                                                        | <b>549</b> |
| <b>Magnetic properties of cobalt(III) complex with 3,5-di-<i>tert</i>-butylsemiquinone and 2,2'-bipyridine</b> . . . . .                                                                  | <b>553</b> |
| <b>Magnetic properties of cobalt(III) bis complex with 3,5-di-<i>tert</i>-butyl-semiquinone and 2,2'-bipyridine</b> . . . . .                                                             | <b>555</b> |
| <b>Magnetic properties of hetero-tetranuclear dicobalt(III)-dicopper(II) complex with diethanolamine and diethanolamine as bridging ligands</b> . . . . .                                 | <b>557</b> |
| <b>Magnetic properties of hetero-tetranuclear dicobalt(III)-dicopper(II) complex with diethanolamine and diethanolamine as bridging ligands</b> . . . . .                                 | <b>559</b> |
| <b>Magnetic properties of hetero-tetranuclear dicobalt(III)-dicopper(II) complex with diethanolamine and diethanolamine as bridging ligands</b> . . . . .                                 | <b>561</b> |
| <b>Magnetic properties of hetero-tetranuclear dicobalt(III)-dicopper(II) complex with diethanolamine and diethanolamine as bridging ligands</b> . . . . .                                 | <b>563</b> |
| <b>Part IX Rh</b> . . . . .                                                                                                                                                               | <b>565</b> |
| <b>Magnetic properties of hetero-trimetallic Rh<sup>III</sup><sub>2</sub>-Ni<sup>II</sup> oxalate complex as <i>trans</i>-4-(4-dimethylaminostyryl)-1-methylpyridinium salt</b> . . . . . | <b>567</b> |
| <b>Volume 7B</b>                                                                                                                                                                          |            |
| <b>Part X Ni</b> . . . . .                                                                                                                                                                | <b>571</b> |
| <b>Magnetic properties of nickel(II) complex with pyrazolylbis{(thio)methyl}-borate</b> . . . . .                                                                                         | <b>573</b> |

|                                                                                                                       |            |
|-----------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of nickel(II) chloro complex with pyrazolylbis[(thio)<i>t</i>-butyl]borate</b>                 | <b>575</b> |
| <b>Magnetic properties of nickel(II) acetylacetato complex with pyrazolylbis-[(thio)<i>t</i>-butyl]borate</b>         | <b>577</b> |
| <b>Magnetic properties of nickel(II) coordination polymer with pyrimidine and pivalate bridges</b>                    | <b>579</b> |
| <b>Magnetic properties of nickel(II) complex with photochromic phenanthroline ligand</b>                              | <b>582</b> |
| <b>Magnetic properties of nickel(II) complex with polynitrile ligand and 4,4'-bipyridine as co-ligand</b>             | <b>585</b> |
| <b>Magnetic properties of nickel(II) complex with terdentate Schiff-base</b>                                          | <b>587</b> |
| <b>Magnetic properties of nickel(II) complex with terdentate Schiff-base</b>                                          | <b>589</b> |
| <b>Magnetic properties of nickel(II) complex with tetradentate Schiff-base and thiocyanate ligands</b>                | <b>591</b> |
| <b>Magnetic properties of nickel(II) complex with tetradentate Schiff-base and thiocyanate ligands</b>                | <b>593</b> |
| <b>Magnetic properties of nickel(II) complex with tetradentate Schiff-base azide ligands</b>                          | <b>595</b> |
| <b>Magnetic properties of nickel(II) complex with tetradentate Schiff-base and azide ligands</b>                      | <b>597</b> |
| <b>Magnetic properties of nickel(II) complex with 1,3-diaminopentane</b>                                              | <b>599</b> |
| <b>Magnetic properties of nickel(II) complex with <math>\beta</math>-iminodiketone</b>                                | <b>601</b> |
| <b>Magnetic properties of nickel(II) complex with 3-aminoflavone</b>                                                  | <b>603</b> |
| <b>Magnetic properties of nickel(II) complex with pyrophosphate as ammonium salt</b>                                  | <b>605</b> |
| <b>Magnetic properties of nickel(II) complex with diethyl(pyridin-4-ylmethyl)phosphate</b>                            | <b>607</b> |
| <b>Magnetic properties of nickel(II) complex with (1R, 2R)-1,2-bis-(5-amino-1,3,4-thiadiazol-2-yl)ethane-1,2-diol</b> | <b>609</b> |
| <b>Magnetic properties of nickel(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b>           | <b>611</b> |
| <b>Magnetic properties of nickel(II) chloro complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b>            | <b>614</b> |

|                                                                                                                                     |     |
|-------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of nickel(II) acetato complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                   | 616 |
| <b>Magnetic properties of polymeric chain assemblies of nickel(II) with bis-pyrazolone</b> .....                                    | 618 |
| <b>Magnetic properties of nickel(II) complex with N,N'-bis-(2-pyridinecarboxalimine)-1,8-diaminonaphthalene</b> .....               | 620 |
| <b>Magnetic properties of nickel(II) acetato complex with 2-hydroxybenzophenone substituted thiosemicarbazone</b> .....             | 622 |
| <b>Magnetic properties of nickel(II) complex with 2-hydroxybenzophenone substituted thiosemicarbazone</b> .....                     | 624 |
| <b>Magnetic properties of nickel(II) aqua complex with 2-hydroxy-4-methylbenzophenone substituted thiosemicarbazone</b> .....       | 626 |
| <b>Magnetic properties of nickel(II) acetato complex with 2-hydroxy-4-methylbenzophenone substituted thiosemicarbazone</b> .....    | 628 |
| <b>Magnetic properties of nickel(II) chloro complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> .....  | 630 |
| <b>Magnetic properties of nickel(II) nitrate complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> ..... | 632 |
| <b>Magnetic properties of nickel(II) acetato complex with N'-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> ..... | 634 |
| <b>Magnetic properties of nickel(II) chloro complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....               | 636 |
| <b>Magnetic properties of nickel(II) nitrate complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....              | 638 |
| <b>Magnetic properties of nickel(II) complex with benzimidazole thiocarbamate</b> .....                                             | 640 |
| <b>Magnetic properties of nickel(II) complex with dibenzo(b)1,8-naphthpyridine</b> .....                                            | 642 |
| <b>Magnetic properties of mixed ligand nickel(II) complex with benzo(b)-1,8-naphthpyridine and 1,10-phenanthroline</b> .....        | 644 |
| <b>Magnetic properties of mixed ligand nickel(II) complex with benzo(b)-1,8-naphthpyridine and bipyridine</b> .....                 | 646 |
| <b>Magnetic properties of mixed ligand nickel(II) complex with 2-mercapto-quinazolin-4-one and 1,10-phenanthroline</b> .....        | 648 |
| <b>Magnetic properties of mixed ligand nickel(II) complex with 2-mercapto-quinazolin-4-one and ethylenediamine</b> .....            | 650 |

---

|                                                                                                                                                             |            |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of mixed ligand nickel(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and 1,10-phenanthroline . . . . .</b>                   | <b>652</b> |
| <b>Magnetic properties of mixed ligand nickel(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and ethylenediamine . . . . .</b>                       | <b>654</b> |
| <b>Magnetic properties of nickel(II) chloro complex with 2,6-diacetylpyridine-bis(carbohydrazone) . . . . .</b>                                             | <b>656</b> |
| <b>Magnetic properties of nickel(II) nitrate complex with 2,6-diacetylpyridine-bis(carbohydrazone) . . . . .</b>                                            | <b>658</b> |
| <b>Magnetic properties of nickel(II) acetate complex with 2,6-diacetylpyridine-bis(carbohydrazone) . . . . .</b>                                            | <b>660</b> |
| <b>Magnetic properties of nickel(II) aqua complex with 2,6-diacetylpyridine-bis(carbohydrazone) . . . . .</b>                                               | <b>662</b> |
| <b>Magnetic properties of nickel(II) complex with Schiff-base derived from 4-aminoantipyrine and 2-aminobenzoic acid . . . . .</b>                          | <b>664</b> |
| <b>Magnetic properties of nickel(II) complex with 16-membered tetraaza-macrocyclic Schiff-base ligand . . . . .</b>                                         | <b>666</b> |
| <b>Magnetic properties of nickel(II) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol . . . . .</b>                            | <b>668</b> |
| <b>Magnetic properties of nickel(II) complex with 22-membered tetraamide macrocyclic ligand . . . . .</b>                                                   | <b>670</b> |
| <b>Magnetic properties of nickel(II) complex with 24-membered tetraamide macrocyclic ligand . . . . .</b>                                                   | <b>672</b> |
| <b>Magnetic properties of nickel(II) complex with sulphonylurea glibenclamide . . . . .</b>                                                                 | <b>674</b> |
| <b>Magnetic properties of nickel(II) chloro complex with 14-membered octaazamacrocyclic ligand . . . . .</b>                                                | <b>676</b> |
| <b>Magnetic properties of nickel(II) nitrate complex with 14-membered octaazamacrocyclic ligand . . . . .</b>                                               | <b>679</b> |
| <b>Magnetic properties of nickel(II) acetate complex with 14-membered octaazamacrocyclic ligand . . . . .</b>                                               | <b>682</b> |
| <b>Magnetic properties of homobimetallic Ni(II)-Ni(II) salt of cyanoimino-methanedithiolate ligand . . . . .</b>                                            | <b>685</b> |
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand . . . . .</b> | <b>687</b> |
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand . . . . .</b> | <b>690</b> |

|                                                                                                                                                          |     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand</b> .....  | 693 |
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand</b> .....  | 696 |
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand</b> .....  | 699 |
| <b>Magnetic properties of <math>\mu</math>-acetato-<math>\mu</math>-pyrazolato dinickel(II) complex with pyrazolato based binucleating ligand</b> .....  | 702 |
| <b>Magnetic properties of homo-bimetallic Ni(II)<sub>2</sub> complex with [N<sub>10</sub>] macrocyclic ligand modified with tetramide function</b> ..... | 705 |
| <b>Magnetic properties of binuclear nickel(II) complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> ....                    | 708 |
| <b>Magnetic properties of dinuclear nickel(II) complex with Schiff-base derived from 2,6-diacetylpyridine and semicarbazide</b> .....                    | 710 |
| <b>Magnetic properties of binuclear nickel(II) complex with N-functionalized cyclam</b> .....                                                            | 712 |
| <b>Magnetic properties of binuclear nickel(II) complex with N-functionalized cyclam</b> .....                                                            | 715 |
| <b>Magnetic properties of binuclear nickel(II) complex with N-functionalized cyclam</b> .....                                                            | 718 |
| <b>Magnetic properties of binuclear nickel(II) complex with N-functionalized cyclam</b> .....                                                            | 721 |
| <b>Magnetic properties of binuclear nickel(II) complex with N-functionalized cyclam</b> .....                                                            | 724 |
| <b>Magnetic properties of double stranded dinuclear nickel(II) helicate</b> .....                                                                        | 727 |
| <b>Magnetic properties of ion-pair complex having tris(bipyridyl)nickel(II) cation complex and octacyanomolybdate anion complex</b> .....                | 730 |
| <b>Magnetic properties of heterobimetallic Ni(II)-Cu(II) salt of cyanoimino-methanedithiolate ligand</b> .....                                           | 732 |
| <b>Magnetic properties of trinuclear nickel(II) complex with benzenetri-carboxylate ligand</b> .....                                                     | 734 |
| <b>Magnetic properties of ion-pair complex having bis(2-thioxo-1,3-dithiole-4,5-dithiolato)nickelate(III) anion and basic fuchsin cation</b> ...         | 737 |

|                                                                                                                                                                                                                           |            |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>Magnetic properties of ion-pair complex containing <math>\text{Ni}(\text{mnt})_2^-</math> monoanion and substituted 4-dimethylaminopyridinium cation</b> . . . . .                                                     | 739        |
| <b>Magnetic properties of ion-pair complex containing <math>\text{Ni}(\text{mnt})_2^-</math> monoanion and substituted 4-dimethylaminopyridinium cation</b> . . . . .                                                     | 741        |
| <b>Magnetic properties of <math>\text{Ni}(\text{III})</math> ion-pair complex having 1-(3',5'-dibromo-benzyl)pyridinium cation and bis(maleonitriledithiolato) nickalate anion</b> . . . . .                              | 743        |
| <b>Magnetic properties of heterotrinnuclear <math>\text{Ni}(\text{II})</math>-<math>\text{Tb}(\text{III})</math>-<math>\text{Ni}(\text{II})</math> complex with 1,1,1-tris[(salicylideneamino)methyl]ethane</b> . . . . . | 745        |
| <b>Magnetic properties of heterotrinnuclear <math>\text{Ni}(\text{II})</math>-<math>\text{Eu}(\text{III})</math>-<math>\text{Ni}(\text{II})</math> complex with 1,1,1-tris[(salicylideneamino)methyl]ethane</b> . . . . . | 748        |
| <b>Magnetic properties of heterotrinnuclear <math>\text{Ni}(\text{II})</math>-<math>\text{Gd}(\text{III})</math>-<math>\text{Ni}(\text{II})</math> complex with 1,1,1-tris[(salicylideneamino)methyl]ethane</b> . . . . . | 751        |
| <b>Magnetic properties of heterotrinnuclear <math>\text{Ni}(\text{II})</math>-<math>\text{Dy}(\text{III})</math>-<math>\text{Ni}(\text{II})</math> complex with 1,1,1-tris[(salicylideneamino)methyl]ethane</b> . . . . . | 754        |
| <b>Part XI Cu</b> . . . . .                                                                                                                                                                                               | <b>757</b> |
| <b>Magnetic properties of copper(II) complex with Schiff-base</b> . . . . .                                                                                                                                               | 759        |
| <b>Magnetic properties of copper(II) complex with imidazolyl substituted phenol</b> . . . . .                                                                                                                             | 761        |
| <b>Magnetic properties of polymeric copper(II) complex with imidazolyl substituted phenol</b> . . . . .                                                                                                                   | 763        |
| <b>Magnetic properties of <math>\mu</math>-azido and <math>\mu</math>-phenolato bridged copper(II) coordination polymer having salicylideneamine ligand</b> . . . . .                                                     | 765        |
| <b>Magnetic properties of polymeric copper(II) complex with pentafluoro benzoate and pyrazine</b> . . . . .                                                                                                               | 768        |
| <b>Magnetic properties of copper(II) complex with 1-amidino-O-methylurea</b> . . . . .                                                                                                                                    | 771        |
| <b>Magnetic properties of copper(II) complex with 1-amidino-O-ethylurea</b> . . . . .                                                                                                                                     | 773        |
| <b>Magnetic properties of copper(II) complex with 1-amidino-O-n-propylurea</b> . . . . .                                                                                                                                  | 775        |
| <b>Magnetic properties of copper(II) complex with 1-amidino-O-n-butylurea</b> . . . . .                                                                                                                                   | 777        |
| <b>Magnetic properties of copper(II) polymeric complex with 1,2-bis(2-pyrazinecarboxamido)-4,5-dimethylbenzene</b> . . . . .                                                                                              | 779        |

|                                                                                                                                                 |     |
|-------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of dicynamido bridged copper(II) complex with N,N,N',N''-pentamethylethylenetriamine</b> .....                           | 782 |
| <b>Magnetic properties of copper(II) pyrazine bromide</b> .....                                                                                 | 784 |
| <b>Magnetic properties of copper(II) pyrazine chloride</b> .....                                                                                | 786 |
| <b>Magnetic properties of copper(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 788 |
| <b>Magnetic properties of copper(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 790 |
| <b>Magnetic properties of copper(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 792 |
| <b>Magnetic properties of copper(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 794 |
| <b>Magnetic properties of copper(II) complex with <math>\beta</math>-iminodiketone</b> .....                                                    | 796 |
| <b>Magnetic properties of nanocavity zeolite-Y (host) encapsulated copper(II) complex with unsaturated 16-membered macrocyclic ligand</b> ..... | 798 |
| <b>Magnetic properties of copper(II) complex with 3-aminoflavone</b> .....                                                                      | 801 |
| <b>Magnetic properties of copper(III) complex with diethyl(pyridin-4-ylmethyl)phosphate</b> .....                                               | 804 |
| <b>Magnetic properties of mixed ligand copper(II) complex with pyridin-2,6-dicarboxylic acid-N-oxide and bipyridine</b> .....                   | 807 |
| <b>Magnetic properties of mixed ligand copper(II) complex with pyridin-2,6-dicarboxylic acid-N-oxide and 1,2-bis(4-pyridyl)ethene</b> ...       | 810 |
| <b>Magnetic properties of copper(II) complex with dimethyl aminomethyl-phosphine oxide</b> .....                                                | 812 |
| <b>Magnetic properties of copper(II) complex with (1R, 2R)-1,2-bis-(5-amino-1,3,4-thiadiazol-2-yl)ethane-1,2-diol</b> .....                     | 815 |
| <b>Magnetic properties of copper(II) complex of N-heteroaromatic hydrazone</b> .....                                                            | 817 |
| <b>Magnetic properties of copper(II) complex of N-heteroaromatic hydrazone</b> .....                                                            | 819 |
| <b>Magnetic properties of copper(II) chloro complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                                | 821 |
| <b>Magnetic properties of copper(II) chloro complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                                | 823 |
| <b>Magnetic properties of mixed ligand copper(II) chloro complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                   | 825 |
| <b>Magnetic properties of copper(II) complex derived from (3-oxo-2,3-dihydro-benzo[1,4]oxazin-4-yl)-acetic acid</b> .....                       | 827 |

|                                                                                                                                 |     |
|---------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of copper(II) complex with tetrazole-1-acetate</b> .....                                                 | 829 |
| <b>Magnetic properties of copper(II) complex with 5-nitriminotetrazole</b> .....                                                | 831 |
| <b>Magnetic properties of mixed ligand copper(II) complex with ammonia and 1-methyl-5-nitriminotetrazole</b> .....              | 833 |
| <b>Magnetic properties of copper(II) complex with 2-methyl-5-nitroamino-tetrazole</b> .....                                     | 835 |
| <b>Magnetic properties of mixed ligand copper(II) complex with ammonia and 2-methyl-5-nitroamino-tetrazole</b> .....            | 837 |
| <b>Magnetic properties of mixed ligand copper(II) complex with ammonia and 2-methyl-5-nitroamino-tetrazole (hydrated)</b> ..... | 839 |
| <b>Magnetic properties of copper(II) acetato complex with 2-(aminomethyl)pyridine</b> .....                                     | 841 |
| <b>Magnetic properties of copper(II) formato complex with 2-(aminomethyl)pyridine</b> .....                                     | 843 |
| <b>Magnetic properties of copper(II) chloro complex with 2-(aminomethyl)pyridine</b> .....                                      | 846 |
| <b>Magnetic properties of copper(II) acetato complex with 2-(aminomethyl)pyridine (hydrate)</b> .....                           | 848 |
| <b>Magnetic properties of copper(II) formato complex with 2-(aminomethyl)pyridine (hydrate)</b> .....                           | 850 |
| <b>Magnetic properties of polymeric chain assemblies of copper(II) with bis-pyrazolone</b> .....                                | 852 |
| <b>Magnetic properties of copper(II) complex with N,N'-bis-(2-pyridinecarboxalimine)-1,8-diaminonaphthalene</b> .....           | 854 |
| <b>Magnetic properties of copper(II) acetato complex with 2-hydroxybenzophenone substituted thiosemicarbazone</b> .....         | 856 |
| <b>Magnetic properties of copper(II) acetato complex with 2-hydroxybenzophenone substituted thiosemicarbazone</b> .....         | 858 |
| <b>Magnetic properties of copper(II) complex with 2-hydroxybenzophenone substituted thiosemicarbazone</b> .....                 | 860 |
| <b>Magnetic properties of copper(II) acetato complex with 2-hydroxy-4-methylbenzophenone substituted thiosemicarbazone</b> .... | 862 |
| <b>Magnetic properties of copper(II) acetato complex with 2-hydroxy-4-methylbenzophenone substituted thiosemicarbazone</b> .... | 864 |

|                                                                                                                                            |     |
|--------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of copper(II) chloro complex with <i>N'</i>-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> .....  | 866 |
| <b>Magnetic properties of copper(II) nitrate complex with <i>N'</i>-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> ..... | 868 |
| <b>Magnetic properties of copper(II) acetate complex with <i>N'</i>-(furan-3-ylmethylene)-2-(4-methoxyphenylamino)acetohydrazide</b> ..... | 870 |
| <b>Magnetic properties of copper(II) chloro complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....                      | 872 |
| <b>Magnetic properties of copper(II) nitrate complex with 14-membered tetraazamacrocyclic Schiff-base ligand</b> .....                     | 874 |
| <b>Magnetic properties of copper(II) chloro complex with benzimidazole thiocarbamate</b> .....                                             | 876 |
| <b>Magnetic properties of copper(II) sulphate complex with benzimidazole carbamate</b> .....                                               | 878 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-mercapto-quinazolin-4-one and 1,10-phenanthroline</b> .....               | 880 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-mercapto-quinazolin-4-one and ethylenediamine</b> .....                   | 882 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and 1,10-phenanthroline</b> .....      | 884 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-mercapto-3-phenyl-quinazolin-4-one and ethylenediamine</b> .....          | 886 |
| <b>Magnetic properties of copper(II) complex with 2,6-pyridinedicarboxylic acid derivative</b> .....                                       | 888 |
| <b>Magnetic properties of copper(II) chloro complex with 2,6-diacetylpyridine-bis(carbohydrazone)</b> .....                                | 890 |
| <b>Magnetic properties of copper(II) nitrate complex with 2,6-diacetyl- pyridine-bis(carbohydrazone)</b> .....                             | 892 |
| <b>Magnetic properties of copper(II) acetate complex with 2,6-diacetylpyridine-bis(carbohydrazone)</b> .....                               | 894 |
| <b>Magnetic properties of copper(II) aqua complex with 2,6-diacetylpyridine-bis(carbohydrazone)</b> .....                                  | 896 |
| <b>Magnetic properties of copper(II) complex with 1-formylpiperazinyl-dithiocarbamate</b> .....                                            | 898 |

|                                                                                                                                                                 |     |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of copper(II) complex with 2-pyridineformamide-derived thiosemicarbazone</b> .....                                                       | 900 |
| <b>Magnetic properties of copper(II) complex with 2-pyridineformamide-derived thiosemicarbazone</b> .....                                                       | 902 |
| <b>Magnetic properties of copper(II) complex with 4-(2-thiazolylazo) resorcinol</b> .....                                                                       | 904 |
| <b>Magnetic properties of copper(II) complex with Schiff-base derived from 4-aminoantipyrine and 2-aminobenzoic acid</b> .....                                  | 906 |
| <b>Magnetic properties of copper(II) complex with 14-membered tetraaza-macrocyclic Schiff-base ligand</b> .....                                                 | 908 |
| <b>Magnetic properties of copper(II) complex with 16-membered tetraaza-macrocyclic Schiff-base ligand</b> .....                                                 | 911 |
| <b>Magnetic properties of copper(II) complex with Schiff-base derived from <i>o</i>-phthaldehyde and 2-aminophenol</b> .....                                    | 914 |
| <b>Magnetic properties of copper(II) complex with Schiff-base derived from 2,6-diacetylpyridine and semicarbazide</b> .....                                     | 916 |
| <b>Magnetic properties of copper(II) complex with Schiff-base derived from 2,6-diacetylpyridine and thiosemicarbazide</b> .....                                 | 918 |
| <b>Magnetic properties of copper(II) complex with 22-membered tetramide macrocyclic ligand</b> .....                                                            | 920 |
| <b>Magnetic properties of copper(II) complex with 24-membered tetramide macrocyclic ligand</b> .....                                                            | 922 |
| <b>Magnetic properties of copper(II) complex with Schiff-base</b> .....                                                                                         | 924 |
| <b>Magnetic properties of copper(II) complex with sulphonylurea glibenclamide</b> .....                                                                         | 927 |
| <b>Magnetic properties of mixed ligand copper(II) chloro complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and ethylenediamine</b> .....     | 929 |
| <b>Magnetic properties of mixed ligand copper(II) chloro complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and 1,10-phenanthroline</b> ..... | 931 |
| <b>Magnetic properties of mixed ligand copper(II) azido complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and ethylenediamine</b> .....      | 933 |
| <b>Magnetic properties of mixed ligand copper(II) azido complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and 1,10-phenanthroline</b> .....  | 935 |

|                                                                                                                                                                      |     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| <b>Magnetic properties of mixed ligand copper(II) thiocyanate complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and ethylenediamine</b> .....     | 937 |
| <b>Magnetic properties of mixed ligand copper(II) thiocyanate complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and 1,10-phenanthroline</b> ..... | 939 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and ethylenediamine</b> .....                 | 941 |
| <b>Magnetic properties of mixed ligand copper(II) complex with 2-phenyl-3(benzylamino)1,2-dihydroquinazoline-4(3H)-one and 1,10-phenanthroline</b> .....             | 943 |
| <b>Magnetic properties of copper(II) chloro complex with 14-membered octaazamacrocyclic ligand</b> .....                                                             | 945 |
| <b>Magnetic properties of copper(II) nitrate complex with 14-membered octaazamacrocyclic ligand</b> .....                                                            | 948 |
| <b>Magnetic properties of copper(II) acetate complex with 14-membered octaazamacrocyclic ligand</b> .....                                                            | 951 |
| <b>Magnetic properties of 1D copper(II) coordination polymer containing fumarate and N,N,N',N'-tetramethylethane-1,2-diamine</b> .....                               | 954 |
| <b>Magnetic properties of copper(II) chloro complex with tripodal peralkylguanidine derivative</b> .....                                                             | 957 |
| <b>Magnetic properties of copper(II) acetonitrile complex with tripodal peralkylguanidine derivative</b> .....                                                       | 960 |
| <b>Magnetic properties of copper(II) complex with a rhenium chalcocyanide cluster</b> .....                                                                          | 963 |
| <b>Magnetic properties of copper(II) complex with a rhenium chalcocyanide cluster</b> .....                                                                          | 966 |
| <b>Magnetic properties of copper(II) complex with iminonitroxides diradical</b> .....                                                                                | 969 |
| <b>Magnetic properties of polymeric copper(II) complex with {2-[(hydroxylimino)methyl]phenoxy}acetate ligand</b> .....                                               | 972 |
| <b>Magnetic properties of di(<math>\mu</math>-phenolato) bridged dinuclear copper(II) complex with macrocyclic ligand</b> .....                                      | 975 |
| <b>Magnetic properties of di(<math>\mu</math>-phenolato) bridged dinuclear copper(II) complex with macrocyclic ligand</b> .....                                      | 978 |

|                                                                                                                                                                       |      |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| <b>Magnetic properties of dinuclear copper(II) complex with Schiff-base</b> .....                                                                                     | 981  |
| <b>Magnetic properties of dinuclear copper(II) complex with Schiff-base</b> .....                                                                                     | 983  |
| <b>Magnetic properties of dinuclear copper(II) complex with Schiff-base</b> .....                                                                                     | 985  |
| <b>Magnetic properties of dinuclear copper(II) complex with ligand bearing phthalocyanine acid Schiff-base coordination</b> .....                                     | 987  |
| <b>Magnetic properties of hydroxo bridged dinuclear copper(II) complex with Schiff-base</b> .....                                                                     | 990  |
| <b>Magnetic properties of dimeric copper(II) thiocyanato-isothiocyanato complex with tridentate N-donor ligand</b> .....                                              | 993  |
| <b>Magnetic properties of <math>\mu</math>-acetato bridged, dinuclear copper(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> .....         | 995  |
| <b>Magnetic properties of <math>\mu</math>-chloroacetato-bridged dinuclear copper(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> .....    | 998  |
| <b>Magnetic properties of <math>\mu</math>-trichloroacetato-bridged dinuclear copper(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> ..... | 1001 |
| <b>Magnetic properties of <math>\mu</math>-trifluoroacetato-bridged dinuclear copper(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> ..... | 1004 |
| <b>Magnetic properties of <math>\mu</math>-pyrazolato-bridged dinuclear copper(II) complex with N,N'-bis(2-tosylaminobenzylidene)-1,3-diaminopropanol</b> .....       | 1007 |
| <b>Magnetic properties of dinuclear copper(II) acetato complex with poly(dl-aspartic acid)</b> .....                                                                  | 1010 |
| <b>Magnetic properties of binuclear copper(II) complex of N-heteroaromatic hydrazone</b> .....                                                                        | 1013 |
| <b>Magnetic properties of dinuclear copper(II) chloro complex with 3-acetylcoumarin-N(4)-phenylthiosemicarbazone</b> .....                                            | 1016 |
| <b>Magnetic properties of dinuclear mixed ligand copper(II) complex with ammonia and 5-nitriminotetrazole</b> .....                                                   | 1019 |

|                                                                                                                                           |      |
|-------------------------------------------------------------------------------------------------------------------------------------------|------|
| <b>Magnetic properties of copper(II) complex with<br/>N,N'-bis[(3,4-dichloro-phenyl)methylidene]cyclohexan-<br/>1,4-diamine</b> .....     | 1021 |
| <b>Magnetic properties of dinuclear copper(II) chloro complex<br/>with 1,4-diformylpiperazinebis(carbohydrazone)</b> .....                | 1023 |
| <b>Magnetic properties of dinuclear copper(II) nitrato complex<br/>with 1,4-diformylpiperazine bis(carbohydrazone)</b> .....              | 1025 |
| <b>Magnetic properties of dinuclear copper(II) acetato complex<br/>with 1,4-diformylpiperazine bis(carbohydrazone)</b> .....              | 1027 |
| <b>Magnetic properties of dinuclear copper(II) sulphato complex<br/>with 1,4-diformylpiperazine bis(carbohydrazone)</b> .....             | 1029 |
| <b>Magnetic properties of dinuclear copper(II) complex<br/>with bisthiocarbohydrazone</b> .....                                           | 1031 |
| <b>Magnetic properties of dinuclear copper(II) chloro complex with<br/>bisthiocarbohydrazone</b> .....                                    | 1034 |
| <b>Magnetic properties of dinuclear copper(II) bromo complex with<br/>bisthiocarbohydrazone</b> .....                                     | 1036 |
| <b>Magnetic properties of dinuclear copper(II) complex with<br/>biscarbohydrazone</b> .....                                               | 1038 |
| <b>Magnetic properties of dinuclear copper(II) complex with<br/>biscarbohydrazone as perchlorate salt</b> .....                           | 1041 |
| <b>Magnetic properties of dinuclear copper(II) azido complex with<br/>biscarbohydrazone as perchlorate salt</b> .....                     | 1044 |
| <b>Magnetic properties of dinuclear copper(II) chloro complex with<br/>biscarbohydrazone</b> .....                                        | 1047 |
| <b>Magnetic properties of dinuclear copper(II) bromo complex with<br/>biscarbohydrazone</b> .....                                         | 1050 |
| <b>Magnetic properties of binuclear copper(II) complex with<br/>Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> .... | 1052 |
| <b>Magnetic properties of binuclear copper(II) complex with<br/>Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine</b> .... | 1054 |
| <b>Magnetic properties of dinuclear Cu(II) compound with two phase<br/>transitions</b> .....                                              | 1056 |
| <b>Magnetic properties of methoxy bridged dinuclear copper(II)<br/>complex with hexaaza macrocyclic ligand</b> .....                      | 1059 |

|                                                                                                                                                                               |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| <b>Magnetic properties of dinuclear copper(II) complex with thiosemicarbazone and formate ligands</b> .....                                                                   | 1062 |
| <b>Magnetic properties of dinuclear copper(II) complex with thiosemicarbazone and propionate ligands</b> .....                                                                | 1065 |
| <b>Magnetic properties of dinuclear copper(II) complex with thiosemicarbazone and nitrito ligands</b> .....                                                                   | 1068 |
| <b>Magnetic properties of dinuclear copper(II) complex with 3-methyl-5-oxo-N,1-diphenyl-4,5-dihydro-1-H-pyrazol-4-carbothioamide</b> .....                                    | 1071 |
| <b>Magnetic properties of 1D copper(II) coordination polymer containing fumarate and 2,2'-bipyridine</b> .....                                                                | 1074 |
| <b>Magnetic properties of polymeric copper(II) azido complex with <i>trans</i>-4-cotininecarboxylic acid</b> .....                                                            | 1077 |
| <b>Magnetic properties of heterobimetallic cyano-bridged copper(II)-platinum(II) coordination polymer</b> .....                                                               | 1079 |
| <b>Magnetic properties of trinuclear copper(II) complex with bridging ligand 2,4,6-triacetylphlorogucinol and capping ligand 2-[bis(2-hydroxyethyl)amino]ethylamine</b> ..... | 1082 |
| <b>Magnetic properties of trinuclear copper(II) complex with triplesalacen ligand</b> .....                                                                                   | 1085 |
| <b>Magnetic properties of tricopper(II) polyoxomolybdate cluster</b> .....                                                                                                    | 1088 |
| <b>Magnetic properties of trinuclear copper(II) complex with bithiocarbohydrazone</b> .....                                                                                   | 1090 |
| <b>Magnetic properties of linear mixed <math>\mu</math>-acetato, <math>\mu</math>-hydroxo bridged trinuclear copper(II) complex with N-sulfonamide derivative</b> .....       | 1093 |
| <b>Magnetic properties of linear mixed <math>\mu</math>-acetato, <math>\mu</math>-hydroxo bridged trinuclear copper(II) complex with N-sulfonamide derivative</b> .....       | 1096 |
| <b>Magnetic properties of trimetallic, Ag<sub>2</sub>-Cu complex with pyridin-2,6-dicarboxylic acid-N-oxide</b> .....                                                         | 1100 |
| <b>Magnetic properties of tetranuclear copper(II) complex with macrocyclic ligand</b> .....                                                                                   | 1102 |
| <b>Magnetic properties of tetranuclear copper(II) complex with sulfinylcalix[4]arene</b> .....                                                                                | 1105 |
| <b>Magnetic properties of Tetranuclear copper(II) complex with ligand bearing phthalocyanine acid Schiff-base coordination</b> .....                                          | 1108 |

|                                                                                                                                                                                                                                       |             |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Magnetic properties of tetranuclear copper(II) complex derived from (2-carboxymethoxyphenylamino)-acetic acid</b> .....                                                                                                            | 1111        |
| <b>Magnetic properties of <math>\mu</math>-hydroxo tetranuclear copper(II) complex with tetrazole-1-acetate</b> .....                                                                                                                 | 1113        |
| <b>Magnetic properties of tetranuclear copper(II) acetato complex with 2-(aminomethyl)pyridine</b> .....                                                                                                                              | 1115        |
| <b>Magnetic properties of tetranuclear copper(II) complex with 1,2-bis(1,2,4-triazol-4-yl)ethane and benzene-1,3,5-tricarboxylate</b> .....                                                                                           | 1117        |
| <b>Magnetic properties of tetranuclear copper(II) complex with bisthiocarbo-hydrazone</b> .....                                                                                                                                       | 1120        |
| <b>Magnetic properties of heterometallic chromato bridged tetranuclear copper(II) complex with 2,2'-bipyridine</b> .....                                                                                                              | 1123        |
| <b>Magnetic properties of supramolecular assembly with calix[6]arene and copper ions</b> .....                                                                                                                                        | 1126        |
| <b>Magnetic properties of bimetallic copper-molybdophosphonate constructed from [Mo<sub>4</sub>FO<sub>12</sub>] clusters and copper-ligand chains</b> .....                                                                           | 1129        |
| <b>Magnetic properties of tetranuclear mixed ligand copper(II)-terbium(III) Cu<sub>2</sub>-Tb<sub>2</sub> complex with 1-(2-hydroxybenzamido)-2-(2-hydroxy-3-methoxybenzylideneamino)ethane and 3-methoxysalicylaldehyde</b> .....    | 1132        |
| <b>Magnetic properties of tetranuclear mixed ligand copper(II)-dysprosium(III) Cu<sub>2</sub>-Dy<sub>2</sub> complex with 1-(2-hydroxybenzamido)-2-(2-hydroxy-3-methoxybenzylideneamino)ethane and 3-methoxysalicylaldehyde</b> ..... | 1135        |
| <b>Magnetic properties of tetranuclear mixed ligand copper(II)-gadolinium(III) Cu<sub>2</sub>-Gd<sub>2</sub> complex with 1-(2-hydroxybenzamido)-2-(2-hydroxy-3-methoxybenzylideneamino)ethane and 3-methoxysalicylaldehyde</b> ..... | 1138        |
| <b>Magnetic properties of <i>meso</i>-tris(4-nitrophenyl)corroalato-copper complex</b> .....                                                                                                                                          | 1141        |
| <b>Part XII Zn</b> .....                                                                                                                                                                                                              | <b>1143</b> |
| <b>Magnetic properties of zinc(II) complex with iminonitroxides diradical</b> .....                                                                                                                                                   | 1145        |

|                                                                                                                                         |             |
|-----------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Part XIII Ce</b> .....                                                                                                               | <b>1149</b> |
| Magnetic properties of cerium(III) complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine .....                  | 1151        |
| Magnetic properties of cerium complex with hydrogenated Schiff-base ligand .....                                                        | 1153        |
| Magnetic properties of binuclear cerium(III) complex with Schiff-base derived from 4,6-diacetylresorcinol and ethanolamine ....         | 1155        |
| <b>Part XIV Nd</b> .....                                                                                                                | <b>1157</b> |
| Magnetic properties of $\mu$ -cyano bridged heterobimetallic Nd(III)-Mo(V) complex with pyrazine-2-carboxamide .....                    | 1159        |
| <b>Part XV Sm</b> .....                                                                                                                 | <b>1161</b> |
| Magnetic properties of samarium(III) complex with bioactive carboxamide ligand .....                                                    | 1163        |
| <b>Part XVI Eu</b> .....                                                                                                                | <b>1167</b> |
| Magnetic properties of europium(III) complex with a ligand obtained by the condensation of benzildihydrazone and 2-acetylpyridine ..... | 1169        |
| Magnetic properties of europium(III) complex with bioactive carboxamide ligand .....                                                    | 1171        |
| <b>Part XVII Gd</b> .....                                                                                                               | <b>1175</b> |
| Magnetic properties of gadolinium(III) nitronyl nitroxide complex ....                                                                  | 1177        |
| Magnetic properties of gadolinium(III) complex with 4- <i>n</i> -propoxybenzoate .....                                                  | 1180        |
| Magnetic properties of gadolinium(III) complex with 4-trifluoromethoxybenzoate and methanol .....                                       | 1182        |
| Magnetic properties of gadolinium(III) complex with 4-trifluoromethylbenzoate and methanol .....                                        | 1184        |
| Magnetic properties of gadolinium(III) complex with benzoate and dimethylformamide .....                                                | 1186        |
| Magnetic properties of gadolinium(III) complex with 4-ethoxybenzoate .....                                                              | 1188        |
| Magnetic properties of gadolinium(III) complex with benzoate and methanol .....                                                         | 1191        |

---

|                                                                                                                                            |             |
|--------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Magnetic properties of heterobimetallic Gd(III)-Cu(II) complex using copper-based <i>o</i>-phenylenebisoxamato derivative as ligand</b> | <b>1193</b> |
| <b>Part XVIII Tb</b>                                                                                                                       | <b>1195</b> |
| <b>Magnetic properties of terbium(III) complex with benzoate and dimethylformamide</b>                                                     | <b>1197</b> |
| <b>Magnetic properties of terbium(III) complex with 4-trifluoromethyl-benzoate and methanol</b>                                            | <b>1199</b> |
| <b>Magnetic properties of terbium(III) complex with 4-trifluoromethoxybenzoate and methanol</b>                                            | <b>1201</b> |
| <b>Magnetic properties of terbium(III) complex with 4-ethoxybenzoate</b>                                                                   | <b>1203</b> |
| <b>Magnetic properties of terbium(III) complex with 4-<i>n</i>-propoxybenzoate</b>                                                         | <b>1205</b> |
| <b>Magnetic properties of terbium(III) complex with bioactive carboxamide ligand</b>                                                       | <b>1207</b> |
| <b>Magnetic properties of terbium(III) complex with benzoate and methanol</b>                                                              | <b>1209</b> |
| <b>Part XIX Dy</b>                                                                                                                         | <b>1211</b> |
| <b>Magnetic properties of heterobimetallic Dy(III)-Cu(II) complex using copper-based <i>o</i>-phenylenebisoxamato derivative as ligand</b> | <b>1213</b> |