CONTENTS

PREFACE .. .. (v)

CHAPTER 1: EXISTENCE AND FIRST PROPERTIES OF TRANSCENDENTAL NUMBERS 1

1. The distinction between algebraic and transcendental numbers. The algebraic numbers are dense on the real line and in the complex plane 1
2. Height, length, and degree of a polynomial 1
3. Conjugates and minimal polynomial of an algebraic number .. 2
4. Cantor's proof that the algebraic numbers are countable .. 3
5. Cantor's proof that the real numbers are not countable, hence that there exist transcendental numbers 4
6. The measure of a polynomial and its multiplicative property 5
7. The measure of a polynomial expressed in terms of the roots 5
8. A basic inequality for the length of a product of polynomials 7
9. Güting's lower bound for a polynomial at an algebraic point 8
10. Application to the rational approximations of an algebraic number. Liouville's sufficient condition for transcendency. Liouville numbers 10
11. More recent theorems on the rational approximations of algebraic numbers 12
12. Another class of transcendental numbers 12
13. A first remark on the algebraic approximations of transcendental numbers 16
14. An upper bound for positive definite quadratic forms at integral points 16
15. Evaluation of the discriminant of a special quadratic form 18
16. Application to the algebraic approximations of numbers, real or complex 20
17. The general theorem 22
18. A necessary and sufficient condition for the transcendency of a number 25

CHAPTER 2: CONVERGENT LAURENT SERIES AND FORMAL LAURENT SERIES 29

19. Numbers as values of analytic functions at algebraic points. Convergent Laurent series 29
20. Notations on fields, rings of polynomials, and the ring of formal Laurent series 29
21. The ring of formal Laurent series is a field 31
22. Properties of the order of a formal Laurent series 31
23. Differentiation of a formal Laurent series 32
24. The subfield of convergent Laurent series 32

25. Series with coefficients in a field K_0 which satisfy an algebraic differential equation with coefficients in a larger field, also satisfy an algebraic differential equation with coefficients in K_0 33

26. The distinction between formal Laurent series that are algebraic or transcendental over the field of rational functions 35

27. Notations 36

28. A necessary condition for a formal Laurent series to be algebraic 37

29. A strongly lacunary formal Laurent series is transcendental 39

30. Algebraic approximations of transcendental formal Laurent series 40

31. A necessary and sufficient condition for transcendency of formal Laurent series. A stronger result by Osgood 41

32. Eisenstein's theorem on algebraic functions quoted 44

CHAPTER 3: FIRST RESULTS ON THE VALUES OF ANALYTIC FUNCTIONS AT ALGEBRAIC POINTS 45

33. The general problem on function values 45

34. The algebraic points at which an algebraic function is algebraic 45

35. Historical remarks 47

36. Examples of transcendental functions which with all their derivatives are algebraic at all algebraic points 47

37. Both a transcendental function and its inverse may with all their derivatives be algebraic in all algebraic points of a certain disk 50

38. Sequences which may be the zeros of an entire function with rational coefficients 53

39. The analogue for series with finite radius of convergence .. 55

40. Lekkerkerker's theorem 56

41. An unsolved problem 58

CHAPTER 4: LINEAR DIFFERENTIAL EQUATIONS: THE LEMMAS OF SHIDLOVSKI 60

42. Historical remarks 60

43. Notations on linear differential equations and their solutions in formal Laurent series 60

44. Shidlovski's lemma on the order of linear forms in finitely many series 62

45. The vector space V_Q of the solutions $\mathbf{w}$ of a system Q of homogeneous linear differential equations with coefficients in a field $K(z)$ 63

46. Linear forms $\lambda(\mathbf{w})$ with rational coefficients in these solutions. The derived form $D\lambda(\mathbf{w})$ 64

47. Linear spaces of forms λ which are closed under derivation and their bases 66

48. Regular systems Q and their vector spaces V_Q 68

49. When Q is regular, then V_Q has the dimension m over K 69

50. The rank ρ of a special solution $\mathbf{f}$ of Q. A basis of the linear forms λ satisfying $\lambda(\mathbf{f}) = 0$ when $1 \leq \rho \leq m-1$ 70

51. Properties of this basis 71

52. Proof of Shidlovski's Reduction Theorem 72

53. The sequence of linear forms $\lambda_1 = \lambda$ and $\lambda_{h+1} = D\lambda_h$. The linear vector space spanned by these forms is closed under derivation. Its dimension μ 75

54. The corresponding basis $\mathbf{w}_1, \ldots, \mathbf{w}_m$ of V_Q and the rational functions e_{ij} 76

55. Proof that the matrix $\underline{\underline{w}}^{(0)}$ is regular 77

56. An upper bound for the degrees of the rational functions e_{ij} 78

57. The case when the linear forms λ_h have polynomial coefficients 80

58. Conclusion of the proof of Shidlovski's Determinant Theorem 81

59. Specialisation of the polynomial coefficients of λ 84

60. Proof of Shidlovski's generalisation of Siegel's Determinant Theorem 86

CHAPTER 5: LINEAR DIFFERENTIAL EQUATIONS: A LOWER BOUND FOR THE RANK OF THE VALUES OF FINITELY MANY SIEGEL E-FUNCTIONS AT ALGEBRAIC POINTS .. 90

61. Introductory remarks 90

62. Algebraic numbers and their conjugates. Algebraic integers 90

63. A theorem on linear diophantine equations 92

64. E-functions relative to an algebraic number field. Siegel E-functions 94

65. General properties of (Siegel) E-functions 96

66. Solution vectors $\mathbf{f}(z)$ of systems Q of linear differential equations the components of which are Siegel E-functions. The ranks ρ and $\rho(\alpha)$ 98

67. First properties of $\mathbf{f}(z)$ 98

68. Construction of a linear form $\lambda\{\mathbf{f}(z)\}$ which vanishes to a high order at $z = 0$, but has small coefficients 99

69. Estimates for the derived linear forms $\lambda_h\{\mathbf{f}(z)\}$ and their coefficients 102

70. Estimate for $\lambda_h\{\mathbf{f}(\alpha)\}$ 104

71. Exclusion of a trivial case. The general case 106

72. The determinant S 108

73. Estimates for S and its conjugates, and lower bounds for $\rho(\alpha)$ when $\rho = m$ 109

74. The general lower bound for $\rho(\alpha)/\rho$ for systems of homogeneous linear differential equations 112

75. Extension to inhomogeneous systems of linear differential equations 113

CHAPTER 6: LINEAR DIFFERENTIAL EQUATIONS: SHIDLOVSKI'S THEOREMS ON THE TRANSCENDENCY AND ALGEBRAIC INDEPENDENCE OF VALUES OF SIEGEL E-FUNCTIONS 115

76. At algebraic points general E-functions can be algebraic or transcendental 115

77. Finite (algebraic or transcendental) extensions of fields. Notations 116

78. The extension fields U and V 118

79. The products $Z_{h_1 \dots h_{n-1}}$ and their basis representations .. 120

80. An upper estimate for the degrees of their coefficients $E_{hh_1 \dots h_{n-1}}$ 121

81. The relation to Hilbert's function $h(t)$ 122

82. Elementary proof of the lower and upper estimates for $h(t)$ 123

83. Application to the solutions of a homogeneous system Q of linear differential equations 127

84. The derived set $Q(t)$ of differential equations 128

85. Proof of a weak form of Shidlovski's First Main Theorem .. 129

86. Remarks on finite algebraic extensions of the algebraic number field K 131

87. First reduction 132

88. Second reduction 133

89. Conclusion of the proofs of the two Main Results of Shidlovski 135

90. Further study of the components of $\mathbf{f}(z)$ assumed to be Siegel E-functions 136

91. Proof that every transcendental Siegel E-function is algebraic at no more than finitely many algebraic points .. 138

92. Application to the E-functions which satisfy a linear differential equation of order m 140

93. The zeros of such an E-function 141

94. The values at algebraic points of a system of finitely many algebraically independent Siegel E-functions are in general algebraically independent 141

CHAPTER 7: APPLICATIONS OF SHIDLOVSKI'S MAIN THEOREMS TO SPECIAL FUNCTIONS .. 145

95. Introductory remarks 145

96. An arithmetic lemma 145

97. Two further lemmas 147

98. The generalised hypergeometric function $f(z)$ 148

99. It is a Siegel E-function for rational parameters α_i, β_j distinct from the integers 0, -1, -2, ... 149

100. The problem of algebraic independence of Siegel E-functions 150

101. The algebraic independence of finitely many exponential functions, and the general theorem of Hermite-Lindemann .. 151

102. The function $f(z;\boldsymbol{\beta})$ 153

103. A general independence theorem on the solutions of linear differential equations 154

104. The Siegel E-functions $f_k(z;\beta)$ 157

105. The algebraic independence of the solutions of a special system of homogeneous linear differential equations 158

106. Application to the functions $f_k(z;\beta)$ 162

107. Arithmetical consequences 165

108. Oleinikov's method for the study of the algebraic independence of solutions of linear differential equations .. 167

109. The Siegel E-function $f(z;\beta_1,\beta_2)$, and the derived function $K(\lambda,\mu)$ 168

110. Cases when it is an algebraic function 169

111. The case when it satisfies a first order algebraic differential equation 171

112. Application to Siegel's theorem on Bessel functions 174

113. Siegel's more general theorem 175

114. His proof 176

115. The proof continued 178

116. The arithmetic consequences 181

117. The functions $K_k(z)$ and statement of some of their properties 182

CHAPTER 8: FORMAL POWER SERIES AS SOLUTIONS OF ALGEBRAIC DIFFERENTIAL EQUATIONS 185

118. Introductory remark 185

119. Notations 185

120. The problem 186

121. The differential operator $F((w))$ 187

122. The higher derivatives of a product: a general formula .. 187

123. Its application to $F^{(h)}((w))$ 188

124. The expression $F^{(h,k)}((w))$ 189

125. Kakeya's formula for $F^{(h,k)}((w))$ 189

126. Application of the preceding results to $F^{(h)}((f))$ when f is a formal power series satisfying $F((f)) = 0$ 191

127. Recursive formulae for the Taylor coefficients f_h of f .. 193

128. An identity and an inequality 194

129. A recursive inequality for f_k 195

130. The numbers u_k 197

131. The numbers v_k 198

132. Maillet's inequality for f_h 199

133. The case when the coefficients f_h are algebraic numbers .. 200

134. Study of their denominators 201

135. Simplification of this problem 202

136. Estimate of the denominators 204
137. The Theorem of Popken 206
138. Weierstrass's σ-function 207
139. The function $s(u)$ 208
140. The function $f(z)$ 209
141. An application of Popken's theorem to transcendency 210

APPENDIX: CLASSICAL PROOFS OF THE TRANSCENDENCY OF e AND π 213
1. Historical remarks 213
2. Statement of the General Lindemann Theorem 213
3. Reduction to the Special Lindemann Theorem 214

I. HERMITE'S APPROXIMATION FUNCTIONS 215
4. The functions A_k and R_{kl} 215
5. Properties of the A_k 215
6. Properties of the R_{kl}. Integral-free proofs by Hurwitz and Gordan 216

II. APPLICATION OF THE LAST FORMULAE TO PROOFS OF TRANSCENDENCY .. 218
7. The sums S and T 218
8. Hypothesis 1 and its consequence 219
9. Hypothesis 2 and its consequence 221
10. Hypothesis 3 and its consequence 223
11. The three kinds of proof 226

III. PROOFS DEPENDING ON THE NON-VANISHING OF A DETERMINANT 227
12. The determinant D 227
13. A transformation of D 227
14. The asymptotic evaluation of D, and Hermite's first proof 228
15. Hermite's second proof: Recursive evaluation of D 229
16. A simpler proof of Hermite's formula for D 232
17. Venske's proof that D does not vanish 233
18. Weierstrass's proof 235
19. A remark 237

IV. PROOFS DEPENDING ON DIVISIBILITY PROPERTIES 237
20. Arithmetic properties of the A_k 237
21. Hilbert's proof of the transcendency of e 238
22. Hilbert's proof of the transcendency of π 239
23. A remark 241
24. An arithmetic proof using determinants 241
25. Weber's method 242
26. The disadvantages of choosing a prime for the parameter .. 243

V. PROOFS DEPENDING ON ANALYTIC ESTIMATES 244
27. Remarks 244
28. The method by Stieltjes 244
29. Späth's proof of the transcendency of e 245
30. Späth's proof of the General Theorem of Lindemann 246
31. Final remarks 248

BIBLIOGRAPHY 249