

Kind (15 kg; ca. 3–4 J.)

Normwerte	HF (Herzschläge/min)	110	Analgetika	Fentanyl	30–45 µg
	RR _{syst.} (mmHg)	100		Piritramid	1,5 mg
	Blutvolumen (ml)	1200		Nalbuphin i. v.	1,5–3 mg
	AF (Atemzüge/min)	25		Paracetamol rektal/p. o.	225–375 mg
	AZV (ml)	150		Esketamin i. v. (Analgesie)	7,5–15 mg
	AMV (ml)	3750		Esketamin i. v. (Anästhesie)	15–30 mg
Intubation	Spatel, Maske, Guedel-Tubus	# 2	Sedativa	Midazolam i. v.	1,5 mg
	Tubus-ID (mm)	4,5		Midazolam rektal	7,5–15 mg
	Tubus-Größe (Charr)	20	Relaxanzien	Cisatracurium i. v.	2,25 mg
	Intubationstiefe oral (cm)	14		Rocuronium i. v.	9–15 mg
	Intubationstiefe nasal (cm)	17		Succinylcholin i. v.	15 mg
CPR	Defibrillation (J)	60	Gase	Desfluran in 100 % O ₂	MAC: 8–9,2 %
	Adrenalin i. v.	0,15 mg		Isofluran in 100 % O ₂	MAC: 1,28–1,8 %
	Atropin i. v.	0,15–0,3 mg		Sevofluran in 100 % O ₂	MAC: 2,5 %
	Amiodaron i. v.	75 mg		Propofol 1 % i. v. (Narkose-Induktion)	37,5–75 mg
Eigene Notizen			Hypnotika	Propofol 1 % i. v. (Aufrechterhaltung)	15 ml/h
				Etomidat i. v.	2,25–4,5 mg
				Thiopental i. v. (Induktion)	30–90(–120) mg