Essentials in Ophthalmology *Series Editor:* Arun D. Singh

Ahmad A. Aref Rohit Varma *Editors*

Advanced Glaucoma Surgery





Essentials in Ophthalmology

Series editor Arun D. Singh

More information about this series at http://www.springer.com/series/5332

Ahmad A. Aref • Rohit Varma Editors

Advanced Glaucoma Surgery



Editors Ahmad A. Aref, MD Illinois Eye & Ear Infirmary University of Illinois at Chicago School of Medicine Chicago, IL, USA

Rohit Varma, MD, MPH USC Eye Institute Department of Ophthalmology University of Southern California Los Angeles, CA, USA

Videos to this book can be accessed at http://link.springer.com/book/10.1007/978-3-319-18060-1.

 ISSN 1612-3212
 ISSN 2196-890X
 (electronic)

 Essentials in Ophthalmology
 ISBN 978-3-319-18059-5
 ISBN 978-3-319-18060-1
 (eBook)

 DOI 10.1007/978-3-319-18060-1

Library of Congress Control Number: 2015946238

Springer Cham Heidelberg New York Dordrecht London

© Springer International Publishing Switzerland 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer International Publishing AG Switzerland is part of Springer Science+Business Media (www.springer.com)

Contents

1	Advanced Glaucoma Surgery: An Overview Ahmad A. Aref and Rohit Varma	1
2	Trabecular Micro-Bypass Stent for the Treatment of Coexistent Cataract and Mild Open-Angle Glaucomas Wanda D. Hu and Marlene R. Moster	5
3	Trabectome-Mediated Ab Interno Trabeculectomy for Secondary Glaucoma or as a Secondary Procedure Kevin Kaplowitz and Nils A. Loewen	15
4	Lens Extraction in Angle Closure Glaucoma Sameer Trikha, Shamira A. Perera, Rahat Husain, and Tin Aung	31
5	Gonioscopy Assisted Transluminal Trabeculotomy (GATT) for the Treatment of Adult and Developmental Glaucomas Davinder S. Grover and Ronald L. Fellman	41
6	Ab-Interno Subconjunctival Glaucoma Implant for Advanced Open-Angle Glaucomas Alena Reznik and Rohit Varma	51
7	Nonpenetrating Glaucoma Surgery Ziad Khoueir and Tarek M. Shaarawy	57
8	Endoscopic Cyclophotocoagulation in the Treatment of Plateau Iris and Angle-Closure Glaucomas Robert Noecker and Shiyoung Roh	73
9	Ex-PRESS Miniature Glaucoma Shunt for the Treatment of Uveitic Glaucomas Evan Allan and Steven R. Sarkisian Jr.	77
10	Surgical Management of Normal Tension Glaucoma Sarah S. Khodadadeh and James C. Tsai	87
11	Glaucoma Drainage Implant Surgery Combined with Pars Plana Vitrectomy and KPro Implantation Thasarat S. Vajaranant and Ahmad A. Aref	99

12	The Modified Schocket Procedure Luis E. Vazquez and Steven J. Gedde	107
13	Ab-interno Bleb Needling Paul F. Palmberg	113
14	Drainage of Choroidal Effusions Kyla M. Teramoto and Troy M. Tanji	117
15	Surgical Management of Malignant Glaucoma Devesh K. Varma and Iqbal Ike K. Ahmed	125
Index		135

Contributors

Iqbal Ike K. Ahmed, M.D., F.R.C.S.C. Department of Ophthalmology and Vision Sciences, The University of Toronto, Toronto, ON, Canada Trillium Health Partners, Mississauga, ON, Canada

Credit Valley Eyecare, Mississauga, ON, Canada

Evan Allan, M.D. Dean McGee Eye Institute, Oklahoma City, OK, USA

Ahmad A. Aref, M.D. Illinois Eye and Ear Infirmary, University of Illinois at Chicago School of Medicine, Chicago, IL, USA

Tin Aung, F.R.C.Ophth., Ph.D. Singapore Eye Research Institute and Singapore National Eye Centre, Singapore, Singapore

Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore

Ronald L. Fellman, M.D. Glaucoma Associates of Texas, Dallas, TX, USA

Steven J. Gedde, M.D. Bascom Palmer Eye Institute, University of Miami, Miami, FL, USA

Davinder S. Grover, M.D., M.P.H. Glaucoma Associates of Texas, Dallas, TX, USA

Wanda D. Hu, M.D. Miramar Eye Specialists Medical Group, Ventura, CA, USA

Rahat Husain, M.D. Res., F.R.C.Ophth. Singapore Eye Research Institute and Singapore National Eye Centre, Singapore, Singapore

Kevin Kaplowitz, M.D. Department of Ophthalmology, Stony Brook University Hospital Medical Center, Stony Brook, NY, USA

Sarah S. Khodadadeh, M.D. Center for Advanced Eye Care, Vero Beach, FL, USA

Ziad Khoueir, M.D., M.S.Ed. Department of Glaucoma, Beirut Eye Specialist Hospital, Beirut, Lebanon

Nils A. Loewen, M.D., Ph.D. Department of Ophthalmology, University of Pittsburgh, Pittsburgh, PA, USA

Marlene R. Moster, M.D. Glaucoma Division, Wills Eye Hospital, Philadelphia, PA, USA

Robert Noecker, M.D., M.B.A. Ophthalmic Consultants of Connecticut, Fairfield, CT, USA

Paul F. Palmberg, M.D., Ph.D. Bascom Palmer Eye Institute, University of Miami, Miami, FL, USA

Shamira A. Perera, B.Sc. (Hons). F.R.C.Ophth. Singapore Eye Research Institute and Singapore National Eye Centre, Singapore, Singapore

Alena Reznik, M.D. Department of Ophthalmology, USC Eye Institute, University of Southern California, Los Angeles, CA, USA

Shiyoung Roh, M.D. Lahey Clinic, Peabody, MA

Steven R. Sarkisian Jr., M.D. Dean McGee Eye Institute, Oklahoma City, OK, USA

Tarek M. Shaarawy, M.D., M.Sc. Department of Clinical Neurosciences, Ophthalmology Service, University of Geneva, Geneva, Switzerland

Troy M. Tanji, M.D. University of Hawaii John A. Burns School of Medicine, Honolulu, HI, USA

Kyla M. Teramoto, M.D. University of Hawaii John A. Burns School of Medicine, Honolulu, HI, USA

Sameer Trikha, B.Sc. (Hons), F.R.C.Ophth. Singapore Eye Research Institute and Singapore National Eye Centre, Singapore, Singapore

James C. Tsai, M.D., M.B.A. New York Eye and Ear Infirmary of Mount Sinai, New York, NY, USA

Thasarat S. Vajaranant, M.D. Illinois Eye and Ear Infirmary, University of Illinois at Chicago School of Medicine, Chicago, IL, USA

Devesh K. Varma, M.D., F.R.C.S.C. Department of Ophthalmology and Vision Sciences, The University of Toronto, Toronto, ON, Canada

Trillium Health Partners, Mississauga, ON, Canada

Credit Valley Eyecare, Mississauga, ON, Canada

Rohit Varma, M.D., M.P.H. Department of Ophthalmology, USC Eye Institute, University of Southern California, Los Angeles, CA, USA

Luis E. Vazquez, M.D. Bascom Palmer Eye Institute, University of Miami, Miami, FL, USA

About the Authors

Ahmad A. Aref, M.D. graduated from the Feinberg School of Medicine, Northwestern University, Chicago, IL, in 2006 and subsequently completed a residency at the Penn State Hershey Eye Center, Penn State University College of Medicine, and a fellowship in Glaucoma at Bascom Palmer Eye Institute, Miller School of Medicine, University of Miami. Since 2011, he has been Assistant Professor of Ophthalmology and Chief Quality Officer at Illinois Eye and Ear Infirmary, University of Illinois at Chicago School of Medicine. He is board certified by the American Board of Ophthalmology and the Royal College of Physicians and Surgeons of Canada. In 2011, Dr. Aref received an American Glaucoma Society MAPS Award, and he currently sits on the society's Multimedia Committee. He has published 25 articles in peer-reviewed journals and is a reviewer for a number of journals, including *Ophthalmology*, *JAMA Ophthalmology*, and *American Journal of Ophthalmology*.

Rohit Varma, M.D., M.P.H. graduated from Maulana Azad Medical College, University of Delhi, India, and completed fellowships in the Glaucoma Services of Wills Eye Hospital, Philadelphia, and the Doheny Eye Institute, University of Southern California School of Medicine. In 2005, he was appointed Professor (tenured) at the Department of Ophthalmology and Preventive Medicine, University of Southern California, Keck School of Medicine, Los Angeles. Since 2014, Dr. Varma has been Professor and Chairman, Department of Ophthalmology, USC Eye Institute, Keck School of Medicine, Los Angeles. During his distinguished career he has been the recipient of various honors, including the American Academy of Ophthalmology Honor Award (1998) and Senior Achievement Award (2006) and the President's Award from the Glaucoma Research Foundation (2009). Dr. Varma served on the Editorial Advisory Board of Ophthalmology from 1999 until 2013 and since 2010 has been an Examiner for the American Board of Ophthalmology Oral Examinations. He has published approximately 220 articles in peer-reviewed journals and edited/co-edited two books: Essentials of Eye Care: The Johns Hopkins Wilmer Handbook (Lippincott Raven, 1997) and The Optic Nerve in Glaucoma (Lippincott 1992).

Advanced Glaucoma Surgery: An Overview

Ahmad A. Aref and Rohit Varma

The current era has been described as a "Renaissance Period" in glaucoma surgical care. Our care has been revolutionized with innovative techniques and devices designed to enhance physiologic aqueous outflow pathways in patients with mild stages of glaucomatous optic neuropathy. Although these techniques in this specific patient population have deservedly taken the spotlight with regard to advancing glaucoma treatment paradigms, it is important to note similarly transformative treatment strategies for patients with late stages of the disease as well as those experiencing persistent complications in the postoperative period. This textbook aims to shed light on these strategies for a host of complex entities that may present to the glaucoma surgical consultant.

Drs. Hu and Moster describe use of a novel trabecular micro-bypass stent to access and enhance physiologic aqueous outflow pathways for the treatment of mild glaucomatous optic neuropathy (Chap. 2). In its current iteration, a

R. Varma, M.D., M.P.H.

Department of Ophthalmology, USC Eye Institute, University of Southern California, Los Angeles, CA, USA e-mail: rvarma@usc.edu single micro-bypass stent (iStent, Glaukos Corp.) is attached to a disposable inserter. The device is currently approved for insertion in conjunction with cataract surgery. Upgrades in stent design are currently under study and will allow for multiple stents to be placed with a single inserter. Current evidence supports multiple stent insertion for enhanced aqueous outflow [1]. In addition, targeted stenting may increase rates of success by increasing chances of accessing functioning trabecular collector systems.

Drs. Kaplowitz and Loewen describe the use of the trabectome procedure (Neomedix, Inc.) for the treatment of complex angle-closure glaucomas (Chap. 3). Although typically categorized as a microinvasive procedure with typical indications, the authors describe a technique with encouraging results for the reversal of peripheral anterior synechiae. In addition, the authors describe the use of the device after failed conventional glaucoma filtering procedures [2]. Supplemental video of the technique as performed by Dr. Nils Loewen provides invaluable instruction.

The conventional treatment paradigm for angle-closure glaucomas is currently under investigation, with lens extraction becoming a prime consideration in earlier stages of the disease. Drs. Trikha, Perera, Husain, and Aung provide convincing evidence from anterior segment imaging studies to support this approach (Chap. 4) [3]. Indeed, the high-quality pre- and postoperative figures in this chapter reveal a marked improvement

A.A. Aref, M.D. (🖂)

Illinois Eye and Ear Infirmary, University of Illinois at Chicago School of Medicine, Chicago, IL, USA e-mail: ahmadaref@gmail.com; aaref@uic.edu

[©] Springer International Publishing Switzerland 2015 A.A. Aref, R. Varma (eds.), *Advanced Glaucoma Surgery*, Essentials in Ophthalmology, DOI 10.1007/978-3-319-18060-1_1

in angle crowding with presumed enhancement in trabecular outflow. The authors provide guidance for the performance safe cataract removal under challenging preexisting conditions of a shallow anterior chamber with angle closure.

Drs. Grover and Fellman describe a truly innovative technique, Gonioscopy-Assisted Transluminal Trabeculotomy (GATT) (Chap. 5). The technique involves use of the iTrack microcatheter for performance of ab-interno trabeculotomy for the treatment of adult and developmental glaucomas. Advantages of this technique include sparing of the conjunctiva and the ability to visualize safe cannulation of Schlemm's canal. The group recently reported a 39.8 and 52.7 % decrease in IOP with this technique in primary and secondary open angle glaucomas, respectively, after 12 months of follow-up [4].

Drs. Reznik and Varma describe a novel microinvasive procedure for bypass of physiologic outflow pathways (Chap. 6). The procedure is performed via an ab-interno approach, utilizing a porcine gelatin tube (XEN Gel Stent) to access the subconjunctival space. Three lumen sizes are available, with the degree of aqueous flow determined by the Hagen–Poiseuille equation. A multicentered randomized controlled clinical trial investigating the safety and efficacy of the device is currently underway.

Drs. Khoueir and Shaarawy describe nonpenetrating glaucoma surgical procedures (Chap. 7). The authors note a relatively long learning curve with these techniques when compared to conventional glaucoma filtering surgeries. This learning curve may have slowed the adoption of these techniques which offer a superior safety profile when compared to trabeculectomy surgery [5]. Step-by-step instruction and supplemental video provided by the authors help to soften this learning curve for the novice while providing invaluable pearls for the experienced surgeon.

Dr. Noecker describes the use of endoscopic cyclophotocoagulation for the treatment of plateau iris and angle-closure glaucomas (Chap. 8). In a modification of the technique typically employed for open angle glaucomas, Dr. Noecker describes

treatment of the posterior aspect of the ciliary processes. This allows for pulling of the anterior processes away from the iris with subsequent angle opening. This technique, termed endocycloplasty, represents an important advance as angle-closure associated with plateau iris may persist after laser iridoplasty and/or lens extraction [6, 7].

Drs. Allan and Sarkisian describe surgical pearls for Ex-Press Mini-Glaucoma Device (Alcon Laboratories, Inc.) implantation for the treatment of complex glaucomas, including uveitic glaucoma (Chap. 9). The device may offer an advantage over conventional trabeculectomy in uveitic diseases as it is typically associated with less postoperative inflammation and anterior chamber shallowing. Supplemental surgical video by Dr. Sarkisian demonstrates key steps of the procedure.

Normal-tension glaucoma represents a challenging entity as it is often difficult to achieve significant intraocular pressure lowering from apparently moderate baseline levels. Drs. Khodadadeh and Tsai describe trabeculectomy pearls for safe achievement of these target levels (Chap. 10). The surgeons also describe their preferred topical anesthetic technique so as to reduce the risks for retrobulbar anesthesia.

The Boston Keratoprosthesis represents a landmark achievement in the treatment of corneal blindness. Unfortunately, postoperative glaucoma represents an important and visionthreatening complication after KPro implantation. Drs. Vajaranant and Aref describe a technique for combined glaucoma, pars plana vitrectomy, and KPro surgery (Chap. 11). The technique allows for posterior placement of the glaucoma implant tube in order to allow for optimal contact lens fitting and decreased risk of tube erosion. Addressing the preexisting (and likely future) glaucoma at the time of KPro implantation minimizes the risk of irreversible glaucomatous visual field loss. A recent report by the authors supports this strategy with a relatively low risk of postoperative complications [8].

Postoperative glaucoma may also develop after retinal detachment surgery. The presence of a scleral buckle makes surgical management of the condition somewhat challenging. Drs. Vazquez and Gedde describe the Modified Schocket Procedure to manage this condition (Chap. 12). A supplemental surgical video by Dr. Aref demonstrates the procedure in an infant suffering traumatic glaucoma after prior scleral buckling surgery.

Dr. Palmberg describes an ab-interno technique for trabeculectomy revision once episcleral fibrosis has limited aqueous outflow (Chap. 13). The technique offers logistical, financial, and psychological advantages over conventional transconjunctival bleb needling. Preliminary results of the technique, as performed by Dr. Palmberg, are encouraging.

Ms. Teramoto and Dr. Tanji describe complex pathophysiology of choroidal effusions and hemorrhage (Chap. 14). The development of a postoperative choroidal hemorrhage may represent a devastating complication of ophthalmic surgery. However, appropriate management, as outlined by the authors, may allow for a successful outcome.

Drs. Varma and Ahmed describe the pathophysiology and management of malignant glaucoma (Chap. 15). Surgical techniques offer definitive therapy and include anterior or posterior iridozonulohyaloidotomy and anterior chamber reformation with intraocular lens pushback. It is important to note that core vitrectomy is not an effective stand-alone treatment. Exceptional figures in this chapter guide the reader through the appropriate surgical techniques.

Each section of the current text may serve as a stand-alone chapter for rapid reference by the practitioner during stages of surgical planning. In that regard, the text is meant to guide and assist the clinician in important surgical decision-making for complex ophthalmic conditions. At the same time, the text offers concise, clear explanations of these conditions while addressing advantages and disadvantages to various treatment strategies. In this regard, the text serves as a valuable reference for experienced and novice ophthalmic surgeons alike.

Acknowledgements Conflicts of Interest: Dr. Aref— Consultant for New World Medical, Inc., Speaker for Alcon Laboratories, Speaker for Carl Zeiss Meditec.

Animal Studies: No animal or human studies were carried out by the authors for this article.

Informed Consent: Informed consent was not applicable for this article.

References

- Belovay GW, Naqi A, Chan BJ, Rateb M, Ahmed IIK. Using multiple trabecular micro-bypass stents in cataract patients to treat open-angle glaucoma. J Cataract Refract Surg. 2012;38(11):1911–7.
- Bussel II, Kaplowitz K, Schuman JS, Loewen NA, Trabectome Study Group. Outcomes of ab interno trabeculectomy with the trabectome after failed trabeculectomy. Br J Ophthalmol. 2015;99(2):258–62. doi:10.1136/bjophthalmol-2013-304717.
- Tham CC, Leung DY, Kwong YY, Li FC, Lai JS, Lam DS. Effects of phacoemulsification versus combined phaco-trabeculectomy on drainage angle status in primary angle closure glaucoma (PACG). J Glaucoma. 2010;19(2):119–23.
- Grover DS, Godfrey DG, Smith O, Feuer WJ. Montes de Oca I, Fellman RL. Gonioscopy-assisted transluminal trabeculotomy, ab interno trabeculotomy: technique report and preliminary results. Ophthalmology. 2014;121(4):855–61.
- El Sayyad F, Helal M, El-Kholify H, Khalil M, El-Maghraby A. Nonpenetrating deep sclerectomy versus trabeculectomy in bilateral primary open-angle glaucoma. Ophthalmology. 2000;107(9):1671–4.
- Ritch R, Tham CC, Lam DS. Long-term success of argon laser peripheral iridoplasty in the management of plateau iris syndrome. Ophthalmology. 2004; 111(1):104–8.
- Tran HV, Liebmann JM, Ritch R. Iridoplasty apposition in plateau iris syndrome persists after cataract extraction. Am J Ophthalmol. 2003;135(1):40–3.
- Huh ES, Aref AA, Vajaranant TS, de la Cruz J, Chau FY, Cortina MS. Outcomes of pars plana glaucoma drainage implant in Boston type 1 keratoprosthesis surgery. J Glaucoma. 2014;23:e39–44.

Trabecular Micro-Bypass Stent for the Treatment of Coexistent Cataract and Mild Open-Angle Glaucomas

Wanda D. Hu and Marlene R. Moster

2.1 Introduction

In the past few years, there has been an increasing interest in microinvasive glaucoma surgery (MIGS) procedures as they may play a novel role in the glaucoma treatment algorithm. More traditional ab externo glaucoma procedures such as trabeculectomies and glaucoma drainage devices are known to very effectively reduce intraocular pressure (IOP), but are also associated with numerous sight-threatening complications that may occur even years after surgery (e.g., bleb leak, bleb-related endophthalmitis, hypotony) [1,2]. Because of the relatively high complication rates associated with filtration procedures, practitioners often turn to medications and/or laser trabeculoplasty as first-line therapies to reduce IOP [3]. However, many patients struggle with compliance and ocular surface issues with medications [4,5], and laser trabeculoplasty may only have a limited and modest effect [6,7]. The hope is that MIGS procedures

Miramar Eye Specialists Medical Group, Ventura, CA 93003, USA e-mail: wandahu@gmail.com

M.R. Moster, M.D. Glaucoma Division, Wills Eye Hospital, Philadelphia, PA 19107, USA e-mail: marlenemoster@gmail.com; marlenemoster@aol.com will bridge the gap between medications and more invasive procedures for patients with mild to moderate glaucoma.

MIGS procedures utilize an ab interno approach, sparing the conjunctiva to allow for future glaucoma surgical procedures if necessary, and offer a much higher safety profile with smaller incisions and less distortion of the normal anatomy [8]. Compared to traditional filtering procedures, these procedures are often easier to adopt with fewer postoperative complications and therefore, can be utilized by both glaucoma specialists and comprehensive ophthalmologists in their armamentarium against this potentially blinding disease. Currently, there are several MIGS devices/procedures that may decrease IOP through either (1) improving trabecular outflow to Schlemms' canal (trabectome, Neomedix, Inc.; iStent, Glaukos, Corp.; Hydrus microstent, Ivantis, Inc.), (2) improving outflow through the suprachoroidal space (Cypass, Transcend Medical, Inc.; iStent Supra, Glaukos, Corp.), or (3) creating an alternative outflow pathway through the subconjunctival space (Xen Gel Stent; Aquesys, Inc.).

It is thought that the primary site of outflow resistance in primary open angle glaucoma (POAG) occurs at the juxtacanalicular component of the trabecular meshwork and the endothelial lining of the inner wall of Schlemm's canal [9–11]. A trabecular bypass stent bypasses the region of highest outflow resistance and creates a

© Springer International Publishing Switzerland 2015 A.A. Aref, R. Varma (eds.), *Advanced Glaucoma Surgery*, Essentials in Ophthalmology, DOI 10.1007/978-3-319-18060-1_2 2

W.D. Hu, M.D. (🖂)