

# Index

Note: Italicised *b*, *f* and *t* refer to numbered boxes, figures and tables

5:10, 169  
36 + 5, 169

abbreviations, 167

ABCDE approach, 41

ABCDEFG assessment, 41, 54, 71, 75, 76, 79–82, 86, 88, 94, 96, 112, 115, 121, 123, 124, 126, 129–31, 143

abdominal area, checking, 44b

abdominal examination, 51

abdominal pain, 2, 75, 94

accoucher, 169

acidosis, 155, 157

active second stage, 108

active third stage, 169

acute fatty liver of pregnancy (AFLP), 78

- definition of, 78, 169
- diagnosis, 78
- maternal/fetal mortality rate, 78
- pre-hospital management, 78
- risk factors, 78

acute uterine inversion, 138–40

- complete, 138
- definition of, 138
- diagnosis, 138
- in-house management, 139–40
- partial, 138
- pre-hospital management, 138–9
- repositioning uterus, 139, 140f
- risk factors, 138

admissions procedures, 3

adrenaline, 16, 60f, 66, 93, 155

adult respiratory stress syndrome, 92

airway, 33, 43

albumin, 155

alcohol abuse, 93, 95, 96

all fours position, 127f

allergen, 63t

alpha fetoprotein (AFP), 169

ambulance patient care record, 10

ambulance services

- litigation claims against, 7

midwifery assistance, 112

amenorrhoea, 70

amniotic fluid embolism (AFE), 19, 83

- definition of, 83, 169
- diagnosis, 83
- differential diagnosis of shock in, 63t
- mortality, 15
- pre-hospital care, 16
- pre-hospital management, 83
- rapid management and, 46

amphetamines, 95

ampullary pregnancy, 169

anaemia, 34, 64, 96, 123, 124, 159

anaesthesia, 19

- mortality, 15

analgesics, 8, 64, 71, 80, 84, 109

anaphylactic shock, 20, 30, 62, 63, 63t, 64, 66

antenatal, 169

antenatal care, 1, 52, 124

antepartum, 169

antepartum haemorrhage (APH), 78–80; *see also* postpartum haemorrhage (PPH)

- concealed, 80
- definition of, 78–9, 169
- diagnosis, 79
- diagnosis of shock in, 63t
- external, 81
- interventions for, 80
- pre-hospital management, 79–80
- risk factors, 79
- treatment of, 45

anterior fontanelle, 103

antibiotics, 16, 141, 143, 162, 163, 164

aortocaval compression, 20, 56, 59

- patient position during transfer, 11

Apgar score, 169

apnoea

- primary, 146, 149
- terminal, 146, 147f, 148f

arrhythmias, 97

asphyxia, fetal, 131

aspiration pneumonia, 37

assessment of newborn, 150–1

- outcome, 151

assisted conception, estimated date of delivery, 48

asthma, 30, 92

- diagnosis, 92
- mortality, 15
- pre-hospital management, 92–3

atropine, 69, 139b

back pains, 44, 136, 95

bacterial endocarditis, 96

bag-valve–mask (BVM) technique, 56

benzodiazepines, 77, 88, 95

beta-hCG, 71

bicarbonate, 155

bicornuate uterus, 115, 169

bilirubin, 37t

bimanual uterine compression, 135b, 135f

birth canal, trauma to, 136–7

bleeding; *see also* blood loss; haemorrhage

- checking for, 44
- internal, 44
- in miscarriage, 68
- obstetric history, 49
- postnatal, 2

blood clots, 75, 88

blood glucose, 51, 71, 75, 76, 78, 80, 84, 86, 88, 91, 95, 155

blood loss, 36

- assessing for, 44b, 44f
- in cervical shock, 69
- history of, 49
- obstetric, estimation, 44f
- in placenta praevia, 82
- in placental abruption, 81
- in primary postpartum haemorrhage, 133, 134
- rapid management and, 46
- in shock, 63t

'blood on the floor', 43, 44b

blood pressure, 34–5  
  checking, 52  
  in ectopic pregnancy, 71  
  measurement of, 52  
  primary survey, 43  
blood viscosity, 34  
blood volume, 34  
body, delivery of, 106  
body mass index (BMI), 51  
*Bolam v. Friern Hospital Management Committee* (1957), 10  
born before arrival (BBA), 169  
bougie, 58b  
bowel perforation, 162, 164  
bradycardia, 69, 138, 146, 151, 152  
breast feeding, 143  
  adrenaline in, 16  
breathing, 34, 43; *see also* respiration  
breech birth, algorithm for, 119f  
breech presentation, 114–19; *see also* face presentation  
  algorithm, 119f  
  classification, 114t  
  definition of, 114, 169  
  diagnosis, 115  
  pre-hospital management, 115–18  
  risk factors, 115  
bregma, 103  
broad ligament, 134, 137  
brow, 102, 102f  
brow presentation, 113, 113f, 122  
  definition of, 122  
  diagnosis, 122  
  pre-hospital management, 122  
  risk factors, 122

caesarean section, 1  
  definition of, 169  
perimortem (PMCS/RH), 9, 16, 61–2, 62b  
postpartum infection, 141  
and uterine rupture mortality, 83  
rate, 2

cannulation, 36  
capacity, 8  
capillary refill time (CRT), 71, 76, 80, 84, 143  
carbon monoxide poisoning, 96–8  
  definition of, 96  
  diagnosis, 97  
  incidence of, 96  
  pre-hospital management, 97  
  risk factors, 96  
  symptoms, 97  
carboxyhaemoglobin, 96, 97  
cardiac arrest in pregnancy, 46, 55–61  
  advanced life support algorithm for, 60f  
  incidence, 55  
  management, 59–61  
  mortality rate, 55  
physiological changes  
  airway, 56  
  breathing, 56  
  circulation, 56  
pre-hospital management, 56–9

cricothyroidotomy, 59  
obstetric intubation, 58b  
patient positioning, 56–9, 57f  
supraglottic airway devices, 59  
resuscitation, 55–6

cardiac death, 18  
cardiac disease in pregnancy, 85–6  
  definition of, 85  
  diagnosis, 86  
  maternal deaths, 20  
  mortality rate, 85  
  pre-hospital management, 86  
  risk factors, 85–6

cardiac output, 9, 19, 20, 34–5, 59, 153, 155, 156, 165  
cardiogenic shock, 62, 64, 66  
cardiopulmonary resuscitation (CPR), 16, 60, 61, 62, 66, 112, 126, 155  
Centre for Maternal and Child Health (CMACH), 14  
cephalic, 169  
cephalic presentation, 52, 113f, 115, 120, 125, 128  
cerebral hypoperfusion, 34  
cerebral vein thrombosis, 87  
cervical shock, 63t, 68  
  rapid management and, 46  
cervical trauma, 136  
cervix, 67, 81f, 82f, 103, 104, 120, 134, 161, 169  
  trauma to, 136, 163  
chest compressions, 35, 56, 59–61, 148f, 149, 153–5  
chickenpox, 92, 143  
chorioamnionitis, 133, 142  
circulation, 34–6  
  blood loss, 36  
  and cardiac arrest, 56  
  cardiac output, 34  
  maternal syncope, 35  
  primary survey, 42, 43  
  in third trimester of pregnancy, 35  
  varicose veins, 35  
claims, litigation, 7–8  
Clinical Negligence Scheme for Trusts (CNST), 7  
*Clostridium perfringens*, 142  
clotting factors, 38t  
cocaine, 79, 80, 95, 96  
coccyx, 102  
cognitive aides, 30–1  
coincidental death, definition, 14  
colour of newborn, 150  
colposcopy, 162  
communication, 26–7  
  elements of, 26  
  effective, with feedback loop, 26–7  
competent adults, 8, 10  
complete miscarriage, 67  
compound presentation, 122–3; *see also* presentation, abnormal  
  definition of, 122  
  diagnosis, 122

pre-hospital management, 123  
risk factors, 122  
concealed haemorrhage, 137  
cone biopsy, 162  
Confidential Enquiry into Maternal and Child Health (CEMACH), 14  
Confidential Enquiry into Maternal Deaths (CEMD), 14  
confidentiality, 9, 14  
confirmation bias, 30  
confirmation of death, 9  
consent, 8  
  competent adults, 8, 10  
  implied, 8  
  minors, 8  
  presumed, 8  
continuous positive airways pressure (CPAP), 158  
contractions, 103, 50, 135b  
convulsions, 87–8; *see also* epilepsy  
cord  
  clamping, 149  
  cutting, 106, 107–8, 107f, 131  
  prolapsed, 46, 128–30, 170  
  rupture, 130  
  short, 131  
counselling, 20, 99, 100  
C-reactive protein (CRP), 170  
crack cocaine, 79, 80  
creatinine, 37t, 38t  
cricothyroidotomy, 58b, 59  
  needle, 59  
crowning, 105  
crystalloids, 61, 64, 66b, 75, 80, 84, 93, 95, 143, 155  
cultural issues, 10  
Cushing's syndrome, 73, 74  
cyanosis, 28, 43, 89, 159  
cystic fibrosis, 92  
cystic hygroma, 120, 122

de novo gestational diabetes, 90  
debriefing, 31  
deep vein thrombosis (DVT), 88–9, 164–5, 170  
delivery  
  in back of ambulance, 45  
  of body, 106  
  cutting cord in, 107–8, 108f  
  emergencies after, 133–44  
    acute uterine inversion, 138–40  
    postpartum infection, 141–3  
    primary postpartum haemorrhage, 133–5  
    puerperal sepsis, 141–3  
    secondary postpartum haemorrhage, 136  
    trauma to birth canal, 136–7  
    wound dehiscence, 143–4  
  emergency care in absence of midwife, 109–10  
membranes, 103, 104  
mode of, 2

normal, 101–10  
place of, 1–2  
of placenta, 108  
stages of, 103  
    first stage of, 103–4  
    second stage of, 34, 51, 73, 104  
    third stage of, 108–9  
descent, 104, 106, 123, 124, 128  
dextrose, 75, 155  
diabetes in pregnancy, 6, 89–91  
    de novo diabetes, 90  
    definition of, 89–90  
    diabetic ketoacidosis, 90–1  
    diagnosis, 90–1  
    hypoglycaemia, 90, 91  
    maternal deaths from, 89  
    pre-eclampsia, eclampsia and, 74  
    pregnancy complications, 91  
    pre-hospital management, 91  
    risk factors, 90  
diabetic ketoacidosis (DKA), 90  
    diagnosis, 90–1  
    pre-hospital management, 91  
diastolic blood pressure, 19, 35, 52, 74, 75  
diazemuls, 76  
diazepam, 76, 77f, 88  
direct death, definition, 14  
disability, assessment of, 45  
discharge, 49  
disseminated intravascular coagulation (DIC), 75, 134, 170  
dizziness, 19, 35, 97  
documentation, 10–11  
domestic violence, 17, 82, 93  
Down's syndrome, 169  
dyspnoea, 86, 89, 92, 93, 97  
dysrhythmias, 64, 65f, 80, 84, 87, 90  
dystocia, 46, 48, 91, 109, 118, 125–8, 137, 170  
early pregnancy, complications of, 67–72;  
    *see also* pregnancy  
    assessment unit, 170  
    cervical shock, 69  
    definition of, 170  
    ectopic pregnancy, 69–71  
    miscarriage, 67–9  
    pre-hospital management of, 71  
eclampsia, 2, 77; *see also* pre-eclampsia  
    definition of, 76  
    diagnosis, 76  
    and epilepsy, 87–8  
    hypoxia in, 76  
    mortality rate, 76  
    pre-hospital management, 76  
    rapid management and, 46  
ectopic pregnancy, 69–71; *see also*  
    pregnancy  
    and amenorrhoea, 70  
    biochemical markers, 71  
    clinical history, 70  
    definition of, 69, 170  
    diagnosis, 70–1  
    examination findings, 70  
    laparoscopy and, 162  
    pre-hospital management, 71  
    risk factors, 69  
    ruptured, 19  
    electrocardiogram (ECG), 18, 26, 60f, 86, 89, 91, 160f  
    embolism  
        amniotic fluid (AFE), 15, 16, 19, 62, 63t, 83  
        pulmonary, 63t, 88–9, 92, 100, 164–5  
    emergencies, 2  
        after delivery, 133–44  
        in early pregnancy, 67–72  
        in late pregnancy, 73–84  
    emergency care, 109–10  
    endometritis, 142  
    endotracheal tube, 20, 56, 58b  
    engagement, 104, 170  
    Entonox, 71, 109, 112, 115, 119f  
    environment, evaluation of, 45  
    epilepsy, 19, 87–8  
        definition of, 87  
        diagnosis, 87–8  
        and eclampsia, 76  
        pre-hospital management, 8, 16  
        risk factors, 87  
    epinephrine, 155  
    episiotomy, 143, 170  
    ergometrine, 68, 76, 84  
    error traps, 30  
    errors, medical, 9–10  
    erythrocyte sedimentation rate (ESR), 170  
    *Escherichia coli*, 142  
    estimated date of delivery (EDD), 48, 170  
    ethnic minority women, 55  
    expiratory braking, 146  
        circulatory adaptation, 146  
    express consent, 8  
    extension, 106, 106f  
    external cephalic version (ECV), 114  
    external haemorrhage, 42, 79, 81, 84  
    external rotation, 106  
    face, 102, 102f  
    face presentation, 113, 113f, 120–2; *see also*  
        breech presentation  
        definition of, 120  
        diagnosis, 121  
        mentoanterior, 121f  
        mentoposterior, 121f  
        pre-hospital management, 121  
        risk factors, 120  
    face to pubes delivery, 120  
    fallopian tube, 69, 138  
    fetal alcohol syndrome, 96  
    fetal asphyxia, 131  
    fetal assessment, 52–3  
    fetal head, 104–6  
    fetal heart heard and regular (FHR), 170  
    fetal heart rate, 129  
    fetal heart sounds, 52  
    fetal hypoxia, 35, 128, 130, 145  
    fetal macrosomia, 120, 122, 125  
    fetal movements, 49, 52, 94  
    fetal movements felt (FMF), 170  
    fetal movements not felt (FMNF), 170  
    fetal skull, 102–3, 102f  
    fetus  
        abnormal presentations, 112, 113–14, 114f  
        assessment of, 52–3  
        blood supply, 37  
        breech presentation, 114–19  
        cerebral vasoconstriction, 59  
        in concealed abruption, 81f  
        death in utero, 82  
        definition of, 170  
        in ectopic pregnancy, 69f  
        effect of alcohol misuse on, 96  
        fundal height, 45, 45f, 46, 112  
        intrauterine death, 81  
        legal rights, 8  
        lies, 112, 113–14, 114f  
        miscarriage, 67  
        in placenta praevia, 82f  
        in placental abruption, 81f  
        and pre-eclampsia, 74  
    first degree perineal trauma, 136  
    flexion, 45, 104, 116, 118  
    fluid in airways, expulsion of, 145–6  
    fluids, 45  
    focal convulsion, 87  
    footling breech, 113f, 128  
    fourth degree perineal trauma, 136  
    free birth, 1  
    full blood count, 37t  
    fundal height, 45, 45f, 46, 112  
    fundus, 42, 45, 112, 114, 125, 134, 135, 138, 139  
    gas gangrene, 142, 144, 163  
    gastric acid, 37  
    gastric reflux, 37  
    gastrointestinal system, 33, 37  
    gastro-oesophageal influx, 86  
    general practitioners (GPs), 1  
        out-of-hours record, 10  
    generalised convulsion, 87  
    genital tract, 37  
    gestation, 48  
        definition of, 170  
        and neonatal mortality rate, 111  
        and shock, 63t  
    gestational age, 170  
    gestational diabetes, *see* diabetes in  
        pregnancy  
    gestational epilepsy, 87  
    global overview, 41–2  
    glucagon, 91  
    glucose, 38t  
        blood, 51, 71, 75, 76, 78, 80, 84, 86, 88, 91, 95, 155  
    glyceryl trinitrate (GTN), 86  
    glycosuria, 90  
    Gram-negative organism, 142  
    Gram-positive organisms, 142  
    grand mal seizures, 76  
    grand multiparity, 115, 123

gravid, 170  
 gravidity, 33, 50, 56, 170  
 grazes, 136  
 gurgling sound, 43

haematomas, 137  
 broad ligament, 134, 137  
 paravaginal, 134  
 in primary postpartum haemorrhage, 134  
 vaginal, 137  
 vault, 164  
 vulval, 137

haemodilution, 34, 37t  
 haemoglobin (Hb), 34, 37t, 38t, 96, 97, 170  
 haemolysis-elevated liver enzymes and low platelets (HELLP), 78  
 definition of, 78, 170  
 diagnosis, 78  
 pre-hospital management, 78

haemorrhage, 19, 20, 42  
 antepartum, 45, 78–80  
 assessing bleeding in, 54  
 catastrophic/massive, 42  
 concealed, 137  
 external, 42  
 internal, 36  
 mortality, 15  
 in placental abruption, 81  
 post-gynaecological surgery, 164  
 postpartum, primary, 133–6  
 postpartum, secondary, 136  
 pre-hospital care, 16  
 in torn umbilical cord, 131

haemostatic agents, 137  
 HALT acronym, 30  
 hand-held notes, 3–6, 4–5f, 10, 171  
 handover of patient, 53, 54  
 headaches, new-onset, 16  
 Health and Care Professions Council: *Standards of Conduct, Performance and Ethics*, 10  
 healthcare error  
 causes of, 24  
 extent of, 23–4  
 human error, 24–5  
 improving team and individual performance, 26  
 learning from error, 25  
 types of errors, 24  
 healthcare staff, roles of, 3  
 heart rate of newborn, 150–1  
 hepatitis, 96, 99  
 heroin, 95, 96  
 histamine, 63, 66  
 history, obstetric, 47–50  
 history evaluation, 49  
 history of current pregnancy, 48  
 history of current problem, 49  
 medications and allergies, 48  
 past medical and surgical history, 48  
 past obstetric history, 48

HIV infection, 92, 96, 99  
 home birth, 1–2  
 homicide, 17  
 Human Factors Clinical Working Group and the National Quality Board: concordat statement on human factors, 24  
 hydrocortisone, 66, 93  
 hydrostatic replacement, 139  
 hypercholesterolaemia, 86  
 hyperemesis, 87  
 hyperglycaemia, 90, 91, 155  
 hyperkalaemia, 60f  
 hyperstimulation, 169, 170  
 hypertension, 73–6  
 eclampsia, 76–7  
 in HELLP syndrome, 78  
 pre-eclampsia, 74–6  
 pre-existing, 73  
 pregnancy-induced, 74  
 rapid management and, 46  
 hypertensive disorders of pregnancy  
 maternal mortality, 17  
 pre-hospital care, 19  
 hyperventilation, 34, 92, 97  
 hypocalcaemia, 87  
 hypoglycaemia, 150, 155, 157  
 diagnosis, 90  
 and epilepsy, 87  
 in pregnant women, 90, 91  
 pre-hospital management, 91  
 hypotensive resuscitation, 80  
 hypothermia, 60f, 95, 112, 113, 132, 142, 150, 157, 159, 164  
 hypovolaemia, 60f  
 hypovolaemic shock, 36, 62–4, 80, 130, 131, 134  
 hypoxia, 60f, 63f, 76, 83, 96, 146  
 fetal, 35, 36, 55, 63, 96, 128, 130, 145, 146, 150, 155

hysterectomy, 161–2  
 abdominal, 161  
 laparoscopic, 161  
 radical, 161  
 subtotal, 161  
 total, 161  
 total with bilateral salpingo-oophorectomy, 161  
 vaginal, 161

i-gel, 56, 57  
 ileofemoral thrombi, 88, 89  
 implied consent, 8  
 IMSAFE, 39  
 incomplete miscarriage, 67  
 indirect death, definition, 14  
 induction of labour (IOL), 78, 170  
 inevitable miscarriage, 67  
 infections; *see also* sepsis  
 HIV, 92, 96, 99  
 post-gynaecological surgery, 163–4  
 postpartum, 141–3

urinary tract, 37, 48, 69, 90, 142, 162, 163  
 wound, 142, 163

influenza, 20  
 maternal mortality, 15, 18  
 inpatient antenatal care, 1  
 insulin, 89, 90  
 internal bleeding, 44  
 internal rotation, 105, 105f  
 internal vaginal examination, 52  
 intestinal obstruction, 164  
 intrapartum, 170  
 intrauterine, 170  
 intrauterine death (IUD), 81, 170  
 intravenous fluids, 36, 45, 69, 71, 72, 75, 76, 164

introitus, 170  
 intubation  
 crash, 57, 58b  
 early, 34  
 laryngeal mask airway, 57  
 in late pregnancy, 59  
 in obtunded patients, 97, 152  
 strategy, 58b  
 tracheal, 59, 66f  
 of newborn, 156, 158  
 of preterm babies, 158  
 tube size, 57

ipratropium bromide, 93  
 ischial bone, 102  
 ischial tuberosities, 102  
 isthmus, 69f, 170

kangaroo mother care, 150  
 Korotkoff sound, 74  
 Kussmauls respirations, 91

labial tears, 136  
 laboratory tests, 37–8t  
 labour  
 definition of, 170  
 delivery of placenta, 108–9  
 history, 49  
 membranes, 103  
 normal mechanism of, 104–8  
 crowning, 105  
 cutting the cord, 107–8, 108f  
 delivery of body, 106  
 descent, 104  
 extension, 106, 106f  
 external rotation, 106  
 flexion, 104  
 internal rotation, 105, 105f, 106f  
 restitution, 106  
 obstetric history, 48  
 pain relief, 109  
 preterm, 111–13  
 stages of, 103  
 first stage of, 103–4  
 second stage of, 34, 51, 73, 104  
 third stage of, 108–9

lacerations, 136  
 lambda, 103

laparoscopy, 162  
 hysterectomy via, 161

large loop excision of transformation zone (LLETZ), 162

laryngeal mask airway (LMA), 57

laryngoscopy, 57, 58*b*

laryngospasm, 152

last menstrual period (LMP), 48, 170

late deaths, 17  
 definition, 14

late pregnancy, emergencies in, 73–84  
 acute fatty liver of pregnancy, 78  
 algorithm for managing shock, 65*f*  
 amniotic fluid embolus, 83  
 antepartum haemorrhage, 78–80  
 breech presentation, 114–19  
 brow presentation, 122  
 compound presentation, 122–3  
 eclampsia, 76–7  
 face presentation, 120–2  
 HELLP syndrome, 78  
 hypertension, 73–6  
 multiple pregnancy, 123–5  
 oblique lie, 113, 123  
 occipitoposterior position, 120  
 placenta praevia, 81–2  
 placental abruption, 80–1  
 pre-eclampsia, 74–6  
 pre-existing hypertension, 73  
 pregnancy-induced hypertension, 74  
 prolapsed umbilical cord, 128–30  
 shock, pre-hospital management  
 in, 83–4  
 short cords, 131  
 shoulder dystocia, 125–8  
 transverse lie, 113, 123  
 umbilical cord rupture, 130  
 uterine rupture, 82–3

leadership, 27

legal and ethical issues, 7–11  
 confidentiality, 9  
 consent and capacity, 8  
 cultural issues, 10  
 documentation, 10–11  
 litigation claims, impact of, 7–8  
 medical errors and negligence, 9–10  
 professional accountability, 10  
 recognition of death, 9

lies, 112, 113–14, 113*f*; *see also*  
 positions  
 definition of, 113, 171  
 longitudinal, 113*f*  
 oblique, 113, 123  
 transverse, 113, 113*f*, 123, 128  
 unstable, 113

liquor, 51, 53, 103, 109, 115, 133,  
 152, 171

lithotomy, 88, 115, 119*f*, 171

litigation claims, impact of, 7–8

live birth, 171

liver function tests, 38*t*

lochia, 142, 171

longitudinal lie, 113, 113*f*

Lovset's manoeuvre, 116, 117*f*

lower segment caesarean section (LSCS), 169

lung capacity, 34, 93

magnesium sulphate, 2, 76, 77*f*, 93

malpositions, 113–14

malpresentations, 113–14

Marfan's syndrome, 86

massive external haemorrhage, 42

maternal age, 74, 79, 80, 86, 90, 125, 133

maternal death, definitions, 14

maternal mortality rate, 13  
 MBRRACE-UK (2014), 15  
 MBRRACE-UK (2015), 16  
 MBRRACE-UK (2016), 17  
 MBRRACE-UK (2017), 19, 20  
 in resource-poor countries, 14  
 statistics 2009–14, 14, 15*t*

maternal syncope, 35

Mauriceau-Smellie-Veit manoeuvre,  
 116, 117*f*

MBRRACE-UK and Ireland Maternity  
 Mortality reports 2014–17, 13–21  
 background, 14–15  
 2014, 15–16  
 2015, 16–17  
 2016, 17–19  
 2017, 19–20

McRobert's manoeuvre, 118, 126

mean corpuscular volume (MCV), 171

mechanism of injury, 94

meconium, 51, 53, 103, 109, 115  
 aspiration, 152, 153  
 definition, 171

medical errors, 9–10

medical history, 48

membranes, 103  
 bleeding under, 81*f*  
 and colour of liquor, 53  
 and onset of labour, 111  
 and prolapsed cord, 128  
 ruptured, 51, 53, 112, 128, 142

Mendelson's syndrome, 33

meningitis, 20, 87

mental health, pre-hospital care, 17

Mental Health Act, 100

mentoanterior face presentation, 121*f*

mentoposterior face presentation, 121*f*

mentum, 103

midazolam, 88

midwife  
 definition, 171  
 request for, 109, 112, 115, 124, 126  
 roles of, 3*t*

minors  
 consent, 8  
 parental responsibility for, 8

miscarriage, 67–8  
 bleeding, 68  
 complete, 67  
 definition of, 67, 171  
 diagnosis, 68  
 incomplete, 67

inevitable, 67

missed, 67

pre-hospital management, 68

risk factors, 68

threatened, 67

misoprostol, 68, 134

missed miscarriage, 67

mode of delivery, 2

morphine, 63*t*, 64, 65*f*, 66, 71, 80, 84, 86,  
 95, 109

multigravida, 78, 79, 80, 104, 171

multip, *see* multigravida

multiparity, 1–2, 78, 115, 123, 124, 128, 133

multiple pregnancy, 123–5  
 definition of, 123, 171  
 diagnosis, 124  
 pre-hospital management, 124–5  
 risk factors, 124

muscle twitching, 75

myocardial infarction (MI), 86

naloxone, 155

National Perinatal Epidemiology Unit  
 (NPEU), Oxford, 14

nebuliser, 66, 93

neck stiffness, 16

necrotising fascitis, 144, 163

negligence, 9–10

neonatal, 171

neonatal death (NN), 171

neurogenic shock, 62, 64, 138

neuroses, 99

non-rebreathing mask, 34, 64, 66, 86, 97, 138

normal labour and delivery, 101–10  
 emergency care in absence of midwife,  
 109–10  
 female pelvis, 101, 101*f*  
 fetal skull, 102–3  
 first stage of labour, 103–4  
 normal mechanism of, 104–8  
 crowning, 105  
 cutting the cord, 107–8, 108*f*  
 delivery of body, 106  
 descent, 104  
 extension, 106, 106*f*  
 external rotation, 106  
 flexion, 104  
 internal rotation, 105, 105*f*, 106*f*  
 restitution, 106

stages of, 103  
 first stage of, 103–4  
 second stage of, 34, 51, 73, 104  
 third stage of, 108–9

normoglycaemia, 90

notes, hand-held pregnancy, 3–6, 4–5*f*,  
 10, 171

obesity, 33, 62, 68, 74, 78, 85, 88, 89, 90, 125,  
 133, 164

maternal mortality, 15

oblique lie, 113, 123

obstetric abbreviations, 167

obstetric haemorrhage, treatment of, 36

obstetric history, 47–50  
 bleeding, 49  
 discharge, 49  
 evaluation of, 49  
 fetal movements, 49  
 global overview, 41–2, 42*f*  
 history of current pregnancy, 48  
 history of current problem, 49  
 key findings from, 50*b*  
 labour, 49  
 medications and allergies, 48  
 pain, 49  
 past, 48  
 previous medical and surgical history, 48  
 seizures, 49  
 taking, 47–50  
 obstetric primary survey, 41–6  
 airway, 43  
 breathing, 43  
 circulation, 42, 43  
 circulation/massive external haemorrhage, 42  
 disability, 45  
 expose/environment/evaluate, 45  
 fundus, 45, 45*f*  
 getting to point quickly, 46  
 global overview, 41–2, 42*f*  
 time-critical problems, 46  
 obstetric secondary survey, 50–3  
 fetal assessment, 52–3  
 internal vaginal examination, 52  
 measuring blood pressure, 52  
 performing examination in, 50–1  
 obstetric services, 1–6  
 admissions procedures, 3  
 common pre-hospital emergencies, 2  
 mode of delivery, 2  
 organisation, 1  
 place of delivery, 1–2  
 pregnancy notes, using, 3–6, 4–5*f*  
 roles of healthcare staff, 3*t*  
 transfer, 2  
 obstetricians, roles of, 3*t*  
 obstetrics, 1, 171  
 obturator fascia, 102  
 occipitoanterior position, 114, 114*f*  
 occipitoposterior position, 114*f*, 120  
 definition of, 114, 120  
 diagnosis, 120  
 pre-hospital management, 120  
 risk factors, 120  
 occiput, 102, 102*f*, 103, 171  
 occult prolapse, 128, 129  
 oedema  
 and face presentation, 121  
 peripheral, 35  
 in pre-eclampsia, 75, 76  
 pulmonary, 64, 66, 75, 76, 84, 86  
 varicose veins and, 35  
 oligohydramnios, 115  
 opiates, 63*t*, 96, 155, 159  
 opioids, 66, 155  
 oropharyngeal airway device, 57

ovaries, 69*f*  
 overt prolapse, 128–9  
 oxygen, 66, 71, 75, 76, 79, 83, 92, 95, 97, 98, 112, 127, 138, 143  
 oxygen demand, 34  
 oxygen saturation, 34  
 oxyhaemoglobin, 97  
 PACE structure, 28  
 pain, 49  
 abdominal, 2, 75, 94  
 back, 44, 136, 95  
 in miscarriage, 68  
 pain relief, 109  
 paralytic ileus, 164  
 paramedics, roles of, 3*t*  
 paravaginal haematoma, 134  
 parental responsibility, 8  
 parity, 48, 50, 88, 109, 171  
 partial convulsion, 87  
 past obstetric history, 48  
 pathway of care, 6, 171  
 patient hand-held records, 3–6, 4–5*f*, 10, 171  
 patient positioning, 46–7, 47*f*  
 during transfer, 11  
 lateral tilt, 56  
 peak exploratory flow rate (PEFR), 93  
 pelvic cavity, 102  
 pelvic floor repair, 163  
 pelvic inlet, 102  
 pelvic outlet, 102  
 pelvis, 101–2  
 anatomy of, 101–2, 102*f*  
 checking for injury, 44*b*  
 perimortem caesarean section, *see* resuscitative hysterotomy (RH)/PMCS  
 perinatal, 171  
 perinatal psychiatric illness, 99–100  
 definition of, 99  
 diagnosis, 99  
 pre-hospital management, 100  
 perineal infection, 143  
 perineal trauma, 136  
 peripartum cardiomyopathy, 85  
 peripheral vascular resistance, 34  
 peritonism, 63*t*  
 peritonitis, 164  
 physiological second stage, 108, 171  
 place of delivery, 1–2  
 placenta  
 delivery of, 108, 110  
 and increase in cardiac output, 34  
 partial separation of, 133  
 and pre-eclampsia, 74  
 placenta praevia, 81–2  
 and antepartum haemorrhage, 80  
 blood loss in, 82  
 concealed abruption, 81*f*  
 definition of, 81  
 diagnosis, 82  
 liquor in, 103  
 low lying placenta, 82*f*  
 pre-hospital management, 82  
 rapid management and, 46  
 and shock, 63*t*  
 placental abruption, 80–1  
 concealed, 81*f*  
 definition of, 80  
 diagnosis, 80–1  
 haemorrhage in, 81  
 pre-hospital management, 81  
 rapid management and, 46  
 revealed bleeding, 81*f*  
 risk factors, 80  
 placental location, 171  
 plasma, 34, 36, 63  
 plastic bags, use for preterm babies, 157*b*  
 platelets, 37*t*, 38*t*, 78  
 pneumonia, 20  
 polyhydramnios, 115, 133  
 polyuria, 90  
 positions, 114; *see also* lies  
 definition of, 171  
 occipitoanterior, 114, 114*f*  
 occipitoposterior, 114, 114*f*, 120  
 posterior fontanelle, 103  
 post-gynaecological surgery patient, 161–5  
 postnatal, 171  
 postpartum, 171  
 postpartum haemorrhage (PPH), 61, 63, 63*t*, 64, 79; *see also* antepartum haemorrhage (APH)  
 definition of, 171  
 diagnosis of shock, 63  
 primary, 133–5  
 secondary, 136  
 postpartum infection, 141–3  
 definition of, 141  
 diagnosis, 142  
 mortality rate, 141  
 pre-hospital management, 143  
 risk factors, 142  
 sites of, 142–3  
 potassium, 37*t*  
 precipitate labour, 171  
 prednisolone, 93  
 pre-eclampsia, 74–6; *see also* eclampsia; hypertension  
 in acute fatty liver of pregnancy, 78  
 and asthma, 92  
 definition of, 74  
 diagnosis, 74  
 and hypertension, 74  
 mortality rate, 74  
 pre-hospital management, 74  
 risk factors, 74  
 severe, 75  
 pre-existing hypertension, 73  
 pregnancy, 33–7  
 anatomical/physiological changes in, 33–7  
 airway, 33  
 breathing, 34  
 circulation, 34–6  
 gastrointestinal system, 33, 37  
 genital tract, 37  
 early, emergencies in, 67–72

ectopic, 69–71  
 laboratory tests, 37–8  
 late, emergencies in, 73–84  
 management of convulsions in, 77f  
 miscarriage, 67–8  
 multiple, 123–5  
 renal system in, 37  
 third trimester of pregnancy, 33, 34, 35, 37, 42, 56, 59, 74, 94  
 trauma in, 93–5  
 uterus in, 37  
 pregnancy-induced hypertension, 74  
 pregnancy notes, using, 3–6, 4–5f  
 pre-hospital emergencies, 2  
 presentation, abnormal, 113–23  
   breech, 114–19  
   brow, 122  
   compound, 122–3  
   definition of, 113–14, 171  
   face, 120–2  
   occipitoposterior, 120  
   transverse and oblique lies, 123  
 presumed consent, 8  
 preterm babies, 124  
   resuscitation of, 157–9  
 preterm labour, 2, 111–13  
   definition of, 111  
   diagnosis, 112  
   pre-hospital management, 112  
   risk factors, 111  
 primary apnoea, 146, 149  
 primary obstetric survey, 41–6  
   airway, 43  
   breathing, 43  
   circulation, 42, 43  
   disability, 45  
   environment, 45  
   fundus, 45  
   getting to point quickly, 46  
   quick scan assessment, 41  
 primary postpartum haemorrhage, 133–5  
   bimanual uterine compression for, 135b  
   definition of, 133  
   diagnosis, 134  
   diagnosis of shock in, 63t  
   pre-hospital management, 134–5  
   risk factors, 133–4  
   rubbing up uterus, 135, 135f  
 primigravida, 171  
 professional accountability, 10  
 progesterone, 35, 37  
 prolapsed umbilical cord, 128–30  
   definition of, 128  
   diagnosis, 129  
   occult, 128, 129  
   occurrence of, 128  
   overt, 128–9  
   perinatal death from, 128  
   pre-hospital management, 129–30  
   rapid management and, 46  
   risk factors, 128  
 pronouncing death, 11  
 proteinuria, 74, 75, 88

psychiatric illness, perinatal, 99–100  
   definition of, 99  
   diagnosis, 99  
   pre-hospital management, 100  
 psychoses, 19, 20, 99  
 pubic arch, 102  
 pubic bone, 102  
 puerperal sepsis, 141–3  
   definition of, 141  
   diagnosis, 142  
   mortality rate, 141  
   pre-hospital management, 143  
   risk factors, 142  
   sites of infection, 142–3  
 puerperium, 14, 17, 88, 99, 172  
 pulmonary embolism (PE), 63t, 88–9, 92, 100, 164–5, 171  
 pulse oximetry, 97, 158  
 pulse rate, 34, 43, 50, 51, 63t, 71, 75, 76  
 pulseless electrical activity (PEA), 60  
 pupil size, 45  
 pyelonephritis, 37, 142, 163  
  
 Q wave, 89  
 quick scan assessment *see global overview*  
  
 rape in pregnancy, 98–9  
   definition of, 98  
   diagnosis, 98  
   injuries, 99  
   pre-hospital management, 98–9  
   preserving evidence, 99  
   risk factors, 98  
   trauma, 98  
 recognition of death, 9  
 recognition of life extinct, 9  
 red cells, 36, 63, 66  
 renal system, 37  
 respiration  
   first breath, 145  
   initiation of, 145–6  
   newborn, 145–6, 150  
 respiratory disease in pregnancy, 91–3  
   definition of, 91–2  
   diagnosis, 92  
   peak exploratory flow rate, 93  
   pre-hospital management, 92–3  
   risk factors, 92  
 respiratory rate, 34, 43, 50, 51, 71  
 restitution, 106  
 resuscitation, maternal, 55–6  
   cardiopulmonary, 9  
   fluid, 16  
   hypotensive, 80  
   for obstetric cardiac arrest, 60f  
   and trauma, 94  
 Resuscitation Council, 55, 64  
 resuscitation of newborn, 145–60  
   airway, 152  
   airway opening, 153f  
   algorithm, 160f  
   birth outside delivery suite, 159  
   breathing, 153

circulation, 153–4, 154f  
 communication with parents, 159  
 discontinuation of, 156  
 drugs for, 154–5  
 equipment, 148  
 neutral position, 152f  
 normal physiology, 145–6  
 pathophysiology, 146, 147–8f  
 poor response to, 158–9  
 preterm babies, 157–9  
 response to, 156  
 supraglottic airway device (SAG), 156  
 tracheal intubation, 156  
 resuscitative hysterotomy (RH)/  
   perimortem caesarean section  
   (PMCS), 9, 16, 61–2, 62b  
   definition of, 61  
   diagnosis, 62  
   pre-hospital management, 62  
   risk factors, 62  
   steps in, 62b  
 retained placenta, rapid management and, 46  
 retained products of conception (RPOC), 172  
 rheumatic fever, 86  
 road traffic collisions, 93  
  
 S wave, 89  
 sacrum, 101, 102f  
 salbutamol, 66, 93  
 salpingectomy, 162  
 SBAR (situation, background, assessment  
   and recommendation), 26  
 second degree perineal trauma, 136  
 secondary obstetric survey, 50–3  
   fetal assessment, 52–3  
   internal vaginal examination, 52  
   measuring blood pressure, 52  
   performing examination in, 50–1  
 secondary postpartum haemorrhage, 136  
 seizures, 49; *see also* epilepsy  
   in eclampsia, 76  
   management of, 77f  
 Sellick's manoeuvre, 58b  
 sepsis, 20, 141–3; *see also* infections  
   abdominal, 164  
   definition, 142  
   diagnosis, 142  
   maternal death and, 15  
   mortality, 15  
   pre-hospital management, 16, 143  
   in primary postpartum haemorrhage, 63t  
   puerperal, 141–3  
   risk factors, 142  
   sites of infection, 142–3  
 septic shock, 62, 64, 164  
 sexual assault in pregnancy, 98–9  
   definition of, 98  
   diagnosis, 98  
   injuries, 99  
   pre-hospital management, 98–9  
   preserving evidence, 99  
   risk factors, 98  
   trauma, 98

shock in pregnancy, 62–6  
algorithm for, 65f  
definition of, 62  
diagnosis, 63  
differential diagnosis of, 63t  
late pregnancy, pre-hospital  
management in, 83–4  
pre-hospital management, 64–6  
risk factors, 62–3  
specific management, 64–5  
short cords, 131  
shoulder dystocia, 125–8  
definition of, 125  
diagnosis, 125  
pre-hospital management, 125–8  
rapid management and, 46  
risk factors, 125  
shoulder presentation, 113, 113f  
sinciput, 103  
situation awareness, 28–30, 158  
distraction, 29  
team, 30  
skin colour, 43  
smoking, 68, 69, 85, 92, 96, 111, 164  
snoring sound, 43  
sodium, 37t  
sodium bicarbonate, 155  
spine board, 66, 94  
spontaneous rupture of the membranes  
(SRM), 51, 172  
status epilepticus, 87  
stillbirth, 1, 48, 50, 79, 91, 172  
*Streptococcus*, 142  
stroke, 19  
pre-hospital care, 16  
structured approach, 41  
subarachnoid haemorrhage (SAH), 16  
substance misuse in pregnancy, 95–6  
definition of, 95  
diagnosis, 96  
pre-hospital management, 96  
risk factors, 95  
sudden adult death syndrome (SADS),  
18, 85  
suicide, 17, 18, 20  
supraglottic airway device (SGA), 57, 58b,  
59, 156  
surgical uterine evacuation, 163  
sutures, 103  
Swiss cheese model, 24–5, 25f  
syncope, 35, 56  
syntocinon, 139  
syntometrine, 68, 76, 134  
systolic blood pressure, 19, 34, 35, 66,  
69, 71, 74, 75, 80, 84, 95, 134, 141t,  
143, 164

T wave, 89  
tachycardia, 19, 20, 70, 71, 80, 86, 89, 97,  
134, 142, 164  
tachypnoea, 70, 71, 89, 92, 100, 134, 142  
taxi technique, 139

team debriefing, 31  
team performance, 30–1  
team resources, 31  
team working, 27–8  
followership, 27  
hierarchy, 27–8  
leader, 27  
roles, 27–8  
summoning help, 31  
team situation awareness, 30  
temperature of newborn, 149–50  
tension pneumothorax, 60f  
term, 2, 172  
Term Breech Trial, 114  
termination of pregnancy, 19  
third degree perineal trauma, 136  
third trimester of pregnancy, 33, 34, 35, 37,  
42, 56, 59, 74, 94  
threatened miscarriage, 67  
'Three Ps in a Pod' initiative, 18  
thromboembolism, 17, 62, 163, 164  
venous (VTE), 15, 35, 88–9, 163, 165  
thrombosis, 60f  
thrombus, 88  
tidal volume, 34, 58b, 59, 95  
time of birth, recording, 149  
tonic–clonic episode, 49, 76, 87  
tracheal intubation  
in late pregnancy, 57  
of newborn, 156  
of preterm babies, 158  
traffic accidents, trauma from, 93  
transfer of patients, 2  
patient position during, 35f, 36f  
roles of medical staff, 3t  
time-critical problems, 46  
transverse lie, 113, 113f, 123, 128  
trauma in pregnancy, 42, 93–5  
bleeding, 44  
definition of, 93  
diagnosis, 94  
domestic violence, 93  
mechanism of injury, 94  
mortality rate, 93  
pre-hospital management, 94–5  
risk factors, 93  
road traffic collisions, 93  
trauma to birth canal, 136–7  
cervical, 136  
diagnosis, 137  
diagnosis of shock, 63  
haematomas, 137  
hypovolaemic shock, 63  
perineal, 136  
pre-hospital management, 137  
risk factors, 137  
uterine, 136  
tremor, 75, 87  
Trendelenburg position, 129f  
trimester, 33, 63t, 172  
tubal abortion, 172  
tuberculosis, 92

type I diabetes, 89  
type II diabetes, 89

umbilical cord  
clamping, 149  
cutting, 106, 107–8, 107f, 131  
prolapsed, 46, 128–30  
rupture, 130  
short, 131  
unstable lie, 113  
urea, 37t, 38t  
uric acid, 37t  
urinary tract infection, 37, 48, 69, 90, 142,  
162, 163  
uterine atony, 134  
uterine displacement, manual, 18, 46, 56,  
57f, 59  
cupping technique, 57f  
push technique, 57f  
uterine fibroids, 133  
uterine fundus, inversion of, 138–40  
rapid management and, 46  
uterine rupture, 82–3  
definition of, 82  
diagnosis, 83  
diagnosis of shock in, 63t  
pre-hospital management, 83  
rapid management and, 46  
uterus, 35  
bicornuate, 115, 169  
diagram of, 69f  
inversion of, 138–40  
postpartum haemorrhage, 133–5  
repositioning, 140f  
trauma, 136

vaginal examination, 52  
vaginal haematoma, 137  
varicella, 92, 143  
varicose veins, 35, 88, 137  
vasoconstriction, 35, 59  
vasodilatation, 34  
vasovagal attacks, 87  
vault haemotoma, 164  
venous thromboembolism (VTE), 15, 17,  
88–9  
definition of, 88  
diagnosis, 89  
pre-hospital management, 89  
risk factors, 8  
vertex, 102, 102f, 103, 104, 169  
presentation, 102, 102f, 113f, 122  
viability, 67, 71, 91, 95, 109, 112, 172  
viral infection, 143  
vital signs, rapid management and, 46  
vulva varix, 137  
vulval haematoma, 137  
vulval inspection, 51

white blood cells (WBC), 37t, 172  
white cell count, 37t, 38t  
wound infection, 142, 163