

Springer Theses

Recognizing Outstanding Ph.D. Research

Katrin Sattler

Periglacial Preconditioning of Debris Flows in the Southern Alps, New Zealand

EXTRAS ONLINE

 Springer

Springer Theses

Recognizing Outstanding Ph.D. Research

Aims and Scope

The series “Springer Theses” brings together a selection of the very best Ph.D. theses from around the world and across the physical sciences. Nominated and endorsed by two recognized specialists, each published volume has been selected for its scientific excellence and the high impact of its contents for the pertinent field of research. For greater accessibility to non-specialists, the published versions include an extended introduction, as well as a foreword by the student’s supervisor explaining the special relevance of the work for the field. As a whole, the series will provide a valuable resource both for newcomers to the research fields described, and for other scientists seeking detailed background information on special questions. Finally, it provides an accredited documentation of the valuable contributions made by today’s younger generation of scientists.

Theses are accepted into the series by invited nomination only and must fulfill all of the following criteria

- They must be written in good English.
- The topic should fall within the confines of Chemistry, Physics, Earth Sciences, Engineering and related interdisciplinary fields such as Materials, Nanoscience, Chemical Engineering, Complex Systems and Biophysics.
- The work reported in the thesis must represent a significant scientific advance.
- If the thesis includes previously published material, permission to reproduce this must be gained from the respective copyright holder.
- They must have been examined and passed during the 12 months prior to nomination.
- Each thesis should include a foreword by the supervisor outlining the significance of its content.
- The theses should have a clearly defined structure including an introduction accessible to scientists not expert in that particular field.

More information about this series at <http://www.springer.com/series/8790>

Katrin Sattler

Periglacial Preconditioning of Debris Flows in the Southern Alps, New Zealand

Doctoral Thesis accepted by
Victoria University of Wellington, Wellington,
New Zealand

Author

Dr. Katrin Sattler
School of Geography, Environment
and Earth Sciences
Victoria University of Wellington
Wellington
New Zealand

Supervisors

AProf. Andrew Mackintosh
Antarctic Research Centre
Victoria University of Wellington
Wellington
New Zealand

Dr. Mairéad de Róiste
School of Geography, Environment
and Earth Sciences
Victoria University of Wellington
Wellington
New Zealand

Dr. Brian Anderson
Antarctic Research Centre
Victoria University of Wellington
Wellington
New Zealand

Dr. Kevin Norton
School of Geography, Environment
and Earth Sciences
Victoria University of Wellington
Wellington
New Zealand

Additional material to this book can be downloaded from <http://extras.springer.com>.

ISSN 2190-5053

Springer Theses

ISBN 978-3-319-35073-8

DOI 10.1007/978-3-319-35074-5

ISSN 2190-5061 (electronic)

ISBN 978-3-319-35074-5 (eBook)

Library of Congress Control Number: 2016938670

© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Disclaimer: Every effort has been made to contact the copyright holders of the figures and tables which have been reproduced from other sources. Anyone who has not been properly credited is requested to contact the publisher, so that due acknowledgement may be made in subsequent editions.

Printed on acid-free paper

This Springer imprint is published by Springer Nature

The registered company is Springer International Publishing AG Switzerland

Parts of this thesis have been published in the following journal article:

Material from Chap. 4 has been published in: Sattler K, Anderson B, Mackintosh A, Norton K, de Róiste M (2016) Estimating permafrost distribution in the maritime Southern Alps, New Zealand, based on climatic conditions at rock glacier sites. *Frontiers in Earth Science* 4. doi:[10.3389/feart.2016.00004](https://doi.org/10.3389/feart.2016.00004)

Supervisor's Foreword

Debris flows share an intimate relationship with mountain permafrost; climate warming is leading to greater melting of permafrost, activation of previously stable mountain slopes, and an increase in associated hazards. The New Zealand Southern Alps contain both mountain permafrost and debris flow activity; however, the relationship between these two phenomena has not been studied.

In this thesis, Katrin Sattler investigates both. First, she develops a spatial and temporal inventory of debris flow activity for the last few decades. Second, she develops a comprehensive rock glacier inventory for New Zealand and uses it to create a first regional estimate of the spatial distribution of mountain permafrost. Finally, Katrin uses statistics to assess the impact of permafrost, intensive frost weathering as well as other non-periglacial environmental factors on debris flow activity in the Southern Alps.

What does she discover? Unlike in the European Alps, permafrost within debris slopes or its degradation do not appear to be playing a key role in debris flow activation, as most debris flows originate below the permafrost zone. In the New Zealand Southern Alps, the availability of readily mobilised sediment, promoted in high-alpine areas by intense frost-weathering activity, and the spatial distribution of heavy rainfall events are more important preconditions and triggers of debris flow activity. This negative result is informative—pointing to subjects (such as understanding bedrock-weathering rates and mountain rainfall patterns) that need more attention. However, arguably the largest contribution of Katrin's work is her estimate of the mountain permafrost distribution in the Southern Alps, a valuable benchmark in today's rapidly changing environment.

Wellington, New Zealand
February 2016

AProf. Andrew Mackintosh

Abstract

The lower boundary of alpine permafrost extent is considered to be especially sensitive to climate change. Ice loss within permanently frozen debris and bedrock as a consequence of rising temperature is expected to increase the magnitude and frequency of potentially hazardous mass wasting processes such as debris flows. Previous research in this field has been generally limited by an insufficient understanding of the controls on debris flow formation. A particular area of uncertainty is the role of environmental preconditioning factors in the spatial and temporal distribution of debris flow initiation in high-alpine areas. This thesis aims to contribute by investigating the influence of permafrost and intensive frost weathering on debris flow activity in the New Zealand Southern Alps. By analysing a range of potential factors, this study explores whether debris flow systems subjected to periglacial influence are more active than systems outside of the periglacial domain.

A comprehensive debris flow inventory was established for thirteen study areas in the Southern Alps. The inventory comprises 1534 debris flow systems and 404 regolith-supplying contribution areas. Analysis of historical aerial photographs, spanning six decades, identified 240 debris flow events. Frequency ratios and logistic regression models were used to explore the influence of preconditioning factors on the distribution of debris flows as well as their effect on sediment reaccumulation in supply-limited systems. The preconditioning factors considered included slope, aspect, altitude, lithology, Quaternary sediment presence, neo-tectonic uplift rates (as a proxy for bedrock fracturing), permafrost occurrence, and frost-weathering intensity. Topographic and geologic information was available in the form of published data sets or was derived from digital elevation models. The potential extent of contemporary permafrost in the Southern Alps was estimated based on the statistical evaluation of 280 rock glaciers in the Canterbury region. Statistical relationships between permafrost presence, mean annual air temperature, and potential incoming solar radiation were used to calculate the spatially distributed probability of permafrost occurrence. Spatially distributed frost-weathering intensities were estimated by calculating the number of annual freeze–thaw cycles

as well as frost-cracking intensities, considering the competing frost-weathering hypotheses of volumetric ice expansion and segregation ice growth.

Results suggest that the periglacial influence on debris flow activity is present at high altitudes where intense frost weathering enhances regolith production. Frost-induced debris production appears to be more efficient in sun-avert than sun-facing locations, supporting segregation ice growth as the dominant bedrock-weathering mechanism in alpine environments. No indication was found that permafrost within sediment reservoirs increases slope instability. Similarly, the presence of permanently frozen bedrock within the debris flow contribution areas does not appear to increase regolith production rates and hence debris flow activity. Catchment topography and the availability of unconsolidated Quaternary deposits appeared to be the cardinal non-periglacial controls on debris flow distribution.

This thesis contributes towards a better understanding of the controls on debris flow formation by providing empirical evidence in support of the promoting effect of intense frost weathering on debris flow development. It further demonstrates the potential and limitations of debris flow inventories for identifying preconditioning debris flow controls. The informative value of regional-scale data sets was identified as a limitation in this research. Improvement in the spatial parameterisation of potential controls is needed in order to advance understanding of debris flow preconditioning factors.

Acknowledgements

Numerous people have accompanied me on this Ph.D. journey and have in some way contributed to the completion of this thesis.

I especially would like to thank my supervisors Andrew Mackintosh, Kevin Norton, Brian Anderson, and Mairéad de Róiste for their support, advice, and feedback on endless drafts. Throughout my journey, all have had an open door for questions and constructive discussions, helping me to regain order when ideas and thoughts were all over the place. A special thank you goes to Brian for sharing his knowledge of and experience in the Southern Alps, which was vital in planning and accomplishing my permafrost fieldwork in the Ben Ohau Range. I also greatly appreciate Brian's support and patience in all things modelling, including provision of the temperature interpolations (mean annual temperature, number of frost cycles, and frost-cracking intensities), which represent a vital part of this research. A big thanks to Dalice Sim who patiently talked me through the intricacies of interpreting logistic regression outputs. A special thank you to Sam McColl for his support and boundless enthusiasm in planning and executing fieldwork, but also for opening his Canterbury home for me and other field assistants to wait when the weather did not play along. I would also like to thank Michael Guggenmos, Robert Williams, Leonhard Gerd, Angelika Riegler, Rory Hart, Brian Anderson, and Andrew Mackintosh for their help in the field. A big thank you to Malcolm Ingham and Gillian Turner who offered their geotechnical experience in a last attempt to make the geoelectric surveys work. I would further like to acknowledge the Department of Conservation, Environment Canterbury, and Andrew Mackintosh for financial support towards fieldwork expenses. Thank you also to my examiners Michael Crozier, Stefan Winkler, and Martin Hoelzle for their constructive comments, suggestions, and effort, which are greatly appreciated. I would particularly like to acknowledge Nick Preston, who secured the Victoria Vice Chancellor's Strategic Research Ph.D. Scholarship for this project, supervised and mentored the first stages of my research, and welcomed me into his home for the first weeks settling in New Zealand. I hope he would enjoy reading through these pages and seeing what the original project has developed into.

I would also like to thank the Spatial, Forestry and Land Operations Team at the Ministry for Primary Industries, who facilitated part-time employment over the last two years to finance the last stretches of my Ph.D. journey. I truly enjoyed showing up to work every Monday morning and applying GIS in areas completely contrary to the barren high-alpine landscape of my research. To be surrounded by a great team of people who are fun to work with, kept me sane and provided a much appreciated distraction from my thesis.

Heartfelt thanks to all my friends inside and outside university, who celebrated with me the highs and supported me through the lows of this journey, but more importantly, were part of all the amazing non-thesis adventures and activities that have made the last five years in New Zealand the fantastic experience they have been.

Lastly, I would like to thank Will and my family for their boundless support over these years. Will, thank you for being at my side, celebrating with me the milestones and steering me through the times when everything just got a bit too much. My love goes to my family, who have been incredibly understanding and put up with my decision to study and live on the other side of the world. Mama and Papa, thank you for making it possible for me to follow my dreams; this thesis is dedicated to you.

Contents

1	Introduction	1
1.1	Background and Rationale	1
1.2	Study Aims and Objectives	4
1.3	Thesis Structure	5
	References	6
2	The Southern Alps	9
2.1	Geography	9
2.2	Geology	11
2.3	Climate	12
2.4	Geomorphology	14
	References	15
3	Debris Flow Inventory	17
3.1	Background—Debris Flows	18
3.1.1	Definition and Characteristics	18
3.1.2	Factors Controlling Debris Flow Initiation	22
3.1.3	Debris Flow Research in New Zealand	32
3.2	Methodology	33
3.2.1	Study Areas	33
3.2.2	Assessment of Debris Flow Activity in the Selected Study Areas	39
3.2.3	Delineation of Debris Flow Contribution Areas	45
3.2.4	Delineation of Steep Rock Surfaces	49
3.3	Results	50
3.3.1	General Debris Flow Activity in Study Areas	50
3.3.2	Event Catalogue	55
3.3.3	Multiple Event Systems	60

3.3.4	Debris Flow Contribution Areas	62
3.4	Discussion	63
3.4.1	Debris Flow Activity in the Southern Alps	63
3.4.2	Debris Flow Activity According to Debris Flow Type	64
3.5	Summary	66
	References	67
4	An Estimate of Alpine Permafrost Distribution	
	in the Southern Alps	77
4.1	Background—Alpine Permafrost	77
4.1.1	Definition and Characteristics	77
4.1.2	Indicators, Prospecting Methods, and Distribution Modelling of Alpine Permafrost	82
4.1.3	Permafrost Research in the Southern Alps	87
4.2	Methodology	89
4.2.1	Rock Glacier Inventory	89
4.2.2	Present-Day Permafrost Distribution Estimate	94
4.3	Results	100
4.3.1	Rock Glacier Inventory	100
4.3.2	Regression Models	109
4.3.3	Permafrost Distribution Estimate for the Southern Alps	119
4.3.4	Local Evaluation of the Permafrost Distribution Estimate	123
4.4	Discussion	133
4.4.1	Rock Glacier Distribution and Derived Permafrost Extent in the Southern Alps	133
4.4.2	Applicability of the BTS Method for Permafrost Prospecting in New Zealand	141
4.5	Summary	143
	References	144
5	Statistical Analysis of Debris Flow Preconditioning Factors	157
5.1	Statistical Methods	158
5.1.1	Frequency Ratios	158
5.1.2	Logistic Regression	159
5.2	Analysis of Preconditioning Factors	160
5.2.1	Slope	160
5.2.2	Aspect	162
5.2.3	Altitude	164
5.2.4	Lithology and Quaternary Deposits	165
5.2.5	Neo-tectonic Uplift Rates	170
5.2.6	Permafrost	173
5.2.7	Regolith Recharge	176
5.3	Summary	202
	References	204

6 Conclusions	209
6.1 Main Findings.	209
6.2 Methodological Aspects of the Investigation of Debris Flow Preconditioning Factors	211
6.2.1 Informative Value of the Datasets Describing Predisposing Factors	211
6.2.2 Informative Value of the Debris Flow Inventory	214
6.2.3 Choice of Statistical Methods	215
6.3 Implications of Findings.	216
6.4 Recommendations for Future Research.	218
6.5 Concluding Remarks	220
References	221
Appendix 1: Debris Flow Inventory in Maps (Sect. 3.3)	223
Appendix 2: List of Aerial Photographs Used for Debris Flow Event Mapping (Sect. 3.2.2)	237
Appendix 3: Regional Characteristics of Debris Flow Activity in the Study Areas (Sect. 3.3.1)	239
Appendix 4: Debris Flow Activity in Study Areas Through Time (Sect. 3.3.2)	247
Appendix 5: Regional Characteristics of Debris Flow Events in the Study Areas (Sect. 3.3.2)	249
Appendix 6: Climate Stations Used for Temperature Interpolations (Sects. 4.2.2, 5.2.6, and 5.2.7)	255
Appendix 7: Preliminary Permafrost Distribution Estimate for the Irishman Stream Valley, Central Ben Ohau Range (Sect. 4.3.4)	259
Appendix 8: Modelled Permafrost Extent and Frost-Weathering Intensities in the Study Areas (Sects. 5.2.6 and 5.2.7).	261
Appendix 9: Variance Inflation Factors for Regression Analyses	275

Abbreviations

Study Regions

AP	Arthur's Pass region
KR	Kaikoura ranges
MA	Mount Aspiring region
MC	Mount Cook region
TT	Two Thumb Range

Study Areas

BBS	Black Birch Stream valley
CC	Camp Creek
CHFS	Castle Hill and Foggy Stream valley
DC	Denas Creek
ES	Enys Stream valley
FC	Forest Creek
GB	Glacier Burn valley
KCET	Kay Creek eastern tributary
LDS	Lower Dart Stream valley
MSV	Middlehead Stream valley
SSV	Stony Stream valley
TSV	Trolove Stream valley
USCT	Upper Stony Creek tributary

Debris Flow Types

COMB	Combined-type debris flow (slide-related)
RIT	Run-off-generated debris flow
SIT	Slide-initiated debris flow