

Contents

Volume 8A

Introduction	1
Part I V	13
Magnetic properties of dimeric vanadium(II) complex with bulky terphenyl thiolate	15
Magnetic properties of vanadium(III) complex with anthranilate	18
Magnetic properties of vanadium(III) complex with imidazol-4-carboxylate	20
Magnetic properties of vanadium(III) chloro complex with <i>trans</i> -urocanic acid	22
Magnetic properties of vanadium(III) complex with bulky terphenyl thiolate	24
Magnetic properties of vanadium(III) 2-dimensional network (vanadyl chains linked by aryldioxide)	26
Magnetic properties of oxovanadium(IV) complex of 2-acetylpyridine semicarbazone	28
Magnetic properties of oxovanadium(IV) complex of 3-acetylpyridine semicarbazone	30
Magnetic properties of oxovanadium(IV) complex of 4-acetylpyridine semicarbazone	32
Magnetic properties of oxovanadium(IV) complex with imidazol-4-carboxylate	34
Magnetic properties of oxovanadium(IV) complex with anthranilate ..	36
Magnetic properties of oxovanadium(IV) complex with dipyrido[3,2-a:2',3'-c]phenazine	38

vii

Magnetic properties of mixed ligand complex of oxovanadium(IV) with <i>N</i>-salicylidene-<i>S</i>-methyldithiocarbazate and 1,10-phenanthroline	40
Magnetic properties of mixed ligand complex of oxovanadium(IV) with <i>N</i>-salicylidene-<i>S</i>-methyldithiocarbazate and dipyrido[3,2-<i>d</i>:2',3'-<i>f</i>]quinoxaline	42
Magnetic properties of mixed ligand complex of oxovanadium(IV) with <i>N</i>-salicylidene-<i>S</i>-methyldithiocarbazate and dipyrido[3,2-<i>a</i>:2',3'-<i>C</i>]phenazine	44
Magnetic properties of oxovanadium(IV) complex with 1,8-<i>N</i>-bis(3-carboxy)disalicylidene-3,6-diazaocatane-1,8-diamine	46
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-dimethylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	48
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-isopropylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	50
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-trimethylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	52
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-tetra-methylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	54
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-penta-methylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	56
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-hexa-methylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	58
Magnetic properties of oxovanadium(IV) complex with bulky <i>N,N'</i>-isopentylenebis(3,5-<i>t</i>Bu₂salicylaldimine) ligand	60
Magnetic properties of dinuclear oxovanadium(IV) complex with Schiff-base derived from 4,6-diacetylresorcinol and diethylenetriamine	62
Magnetic properties of hetero-trimetallic, VO₂Mn complex with macrocyclic oxamido vanadium(IV)-oxo ligand	64
Magnetic properties of hetero-trimetallic, VO₂Cu complex with macrocyclic oxamido vanadium(IV)-oxo ligand	67
Part II Cr	71
Magnetic properties of chromium(III) chloro complex with 4'-(2-pyridyl)-2,2'-6',2''-terpyridine	73

Magnetic properties of chromium(III) azido complex with 4'-(2-pyridyl)-2,2'-6',2''-terpyridine	75
Magnetic properties of chromium(III) thiocyanato complex with 4'-(2-pyridyl)-2,2'-6',2''-terpyridine	77
Magnetic properties of chromium(III) complex with 1,8-<i>N</i>-bis(3-carboxy)- disalicylidene-3,6-diazaocatane-1,8-diamine as acetate salt	79
Magnetic properties of chromium(III) complex with 1,8-<i>N</i>-bis(3-carboxy)- disalicylidene-3,6-diazaocatane-1,8-diamine as chloride salt	81
Magnetic properties of chromium(III) complex with oxalohydrazide	83
Magnetic properties of dinuclear chromium(III) complex with Schiff-base derived from 4,6-diacetylresorcinol and diethylenetriamine	85
Magnetic properties of heterometallic ion-pair complex having copper(II) Schiff-base cation and diaminotetrathiocyanatochromate(III) anion	87
Magnetic properties of heterometallic Cu(II)/Cr(III) complex with an open-chain Schiff-base ligand	90
Part III Mo	93
Magnetic properties of three-dimensional cyano-bridged heterobimetallic Ni(II)-Mo(IV) complex with macrocyclic ligand	95
Magnetic properties of bimetallic oxide of Cu-Mo₆ constructed from molybdenum oxo cluster having bis-<i>N,N</i>(methylphosphonic acid)amine	97
Part IV Mn	101
Magnetic properties of manganese(II) complex with sulpiride drug	103
Magnetic properties of manganese(II) complex with 6-acetyl-1,3,7-trimethylumazine	105
Magnetic properties of imidazole complex of manganese(II) benzenesulfonate	107
Magnetic properties of manganese(II) complex with triazine ligand derived from 3-benzyl-7-hydrazinyl-4H-[1,3,4]thiadiazolo [2,3-<i>c</i>][1,2,4]triazin-4-one	109

Magnetic properties of manganese(II) complex with pyrimidine derived Schiff-base	111
Magnetic properties of manganese(II) complex with pyrimidine derived Schiff-base	113
Magnetic properties of manganese(II) complex with 17-membered pentaaza macrocyclic Schiff-base ligand containing piperazine moiety	115
Magnetic properties of manganese(II) complex with 16-membered pentaaza macrocyclic Schiff-base ligand containing piperazine moiety	117
Magnetic properties of manganese(II) complex with 16-membered pentaaza macrocyclic Schiff-base ligand containing piperazine moiety	119
Magnetic properties of manganese(II) complex with 17-membered pentaaza macrocyclic Schiff-base ligand containing piperazine moiety	121
Magnetic properties of mixed ligand manganese(II) complex with <i>N'</i>-(pyridine-3-carboxyl)-hydrazinecarbodithioic acid ethyl ester and 1,10-phenanthroline	123
Magnetic properties of manganese(II) complex with hexadentate Schiff-base ligand	125
Magnetic properties of manganese(II) complex with mesogenic Schiff-base	127
Magnetic properties of aqua-bridged binuclear manganese(II)	129
Magnetic properties of dinuclear manganese(II) complex with thiophenedicarboxylate	132
Magnetic properties of bis-μ-η'-carboxylate-bridged dinuclear manganese(II) complex containing a tetradentate tripodal ligand	134
Magnetic properties of trinuclear manganese(II) complex with thiophenedicarboxylate	137
Magnetic properties of metallamacrocyclic of manganese(II,III) with (z)-3-(salicylhydrazinocarbonyl)propenoic acid	139
Magnetic properties of manganese(II) complex with 1-3-phenyl-2-(3-phenyl-ureido)propionic acid	142
Magnetic properties of manganese(II) diphosphonate	144
Magnetic properties of manganese(III) complex with sulphonato-substituted Schiff-base	146

Magnetic properties of manganese(III) complex with 1,8-<i>N</i>-bis(3-carboxy)-disalicylidene-3,6-diazaocane-1,8-diamine	149
Magnetic properties of 18-membered hexanuclear manganese(III) macrocycle	152
Magnetic properties of polymeric manganese(III) carboxylate complex with salicylaldehyde	155
Magnetic properties of hexanuclear manganese(III) carboxylate complex with salicylaldehyde	158
Magnetic properties of manganese(III) complex with 2-(2-pyridylmethyleneamino)pyridine	161
Magnetic properties of manganese(II) coordination polymer with mixed pyrimidine-2-carboxylate and oxalate bridges	164
Part V Re	167
Magnetic properties of rhenium(III) complex of 1-(2-pyridylazo)-2-naphthol	169
Magnetic properties of rhenium(III) complex with 1-isoquinoline phenyl ketone	172
Magnetic properties of hetero-trimetallic Gd-Re-Te three dimensional coordination polymer	175
Magnetic properties of hetero-trimetallic Ho-Re-Te three dimensional coordination polymer	177
Part VI Fe	181
Magnetic properties of iron(II) complex with 1,3,5-triazine containing ligand (as perchlorate salt)	183
Magnetic properties of iron(II) complex with 1,3,5-triazine containing ligand as perchlorate salt	186
Magnetic properties of iron(II) complex with 1,3,5-triazine containing ligand as perchlorate salt	189
Magnetic properties of iron(II) complex with 1,3,5-triazine containing ligand as perchlorate salt	191
Magnetic properties of iron(II) complex with 1,3,5-triazine containing ligand as hexafluorophosphate salt	194
Magnetic properties of iron(II) complex with thiocyanate, tris-pyrazolylmethane and pyridine	196

Magnetic properties of iron(II) complex with selenocyanate, tris-pyrazolymethane and pyridine	198
Magnetic properties of iron(II) complex with selenocyanate, tris-pyrazolymethane and imidazole	200
Magnetic properties of dinuclear iron(II) complex with thiocyanate, tris-pyrazolymethane and 4,4'-bipyridine	202
Magnetic properties of dinuclear iron(II) complex with selenocyanate, tris-pyrazolymethane and 4,4'-bipyridine	205
Magnetic properties of ion-pair complex of bis(4-methylpyridine)iron(II) cation and dicyanoaurate(I) anion	208
Magnetic properties of cyanide-bridged Fe₂Cu₂ molecular square having bipyridine ligands	211
Magnetic properties of iron(III) porphyrin radical diperchlorate complex	214
Magnetic properties of tosyl iron(III)-corrole radical cation	217
Magnetic properties of chlorate iron(III)-corrole radical cation	219
Magnetic properties of iron(III) complex with sulphur drug	221
Magnetic properties of iron(III) complex with triazine ligand derived from 3-benzyl-7-hydrazinyl-4H-[1,3,4]thiadiazolo[2,3-c][1,2,4]triazin-4-one	223
Magnetic properties of iron(III) complex with [1,8-<i>N</i>-bis(3-carboxy)-disalicylidene-3,6-diazaocatane-1,8-diamine] as chloride salt	225
Magnetic properties of iron(III) complex with [1,8-<i>N</i>-bis(3-carboxy)-disalicylidene-3,6-diazaocatane-1,8-diamine] as nitrate salt	227
Magnetic properties of iron(III) diperchlorate porphyrin radical complex	229
Magnetic properties of iron(III) diperchlorate porphyrin radical complex	232
Magnetic properties of iron(III) complex with 2-pyridinoformamide derived thiosemicarbazone	234
Magnetic properties of iron(III) complex with 2-pyridinoformamide derived thiosemicarbazone	236
Magnetic properties of iron(III) complex with <i>N</i>-salicylidenearginine	238
Magnetic properties of iron(III) complex with hydroxynaphthylidene-arginine	241

Magnetic properties of iron(III) complex with <i>N</i>-salicylidenelysine . . .	244
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	246
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	249
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	251
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	253
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	256
Magnetic properties of iron(III) complex with 2,6,9-trisubstituted purine	258
Magnetic properties of iron(III) complex with 2-pyridinoformamide derived thiosemicarbazone	260
Magnetic properties of dinuclear iron(III) complex with Schiff-base derived from 4,6-diacetylresorcinol and diethylenetriamine	262
Magnetic properties of dinuclear iron(III) complex incorporating N_4O_3 coordinating helptadentate ligand	264
Magnetic properties of bimetallic cyanide-bridged Ni-Fe complex based on nitroprusside anion and paramagnetic metal complex	267
Magnetic properties of bimetallic cyanide-bridged Mn-Fe complex based on nitroprusside anion and paramagnetic metal complex	269
Magnetic properties of bimetallic cyanide-bridged Mn-Fe complex based on nitroprusside anion and paramagnetic metal complex	272
Magnetic properties of chiral heptanuclear iron(III) cluster with <i>N</i>-(2-hydroxybenzyl)-L-histidine	275
Magnetic properties of bimetallic cyanide-bridged Ni-Fe complex based on nitroprusside anion and paramagnetic metal complex	278
Magnetic properties of metastable porphyrin-iron(IV) diperchlorate complex	280
Magnetic properties of metastable porphyrin-iron(IV) diperchlorate complex	282
Magnetic properties of corrole-iron(IV) chloro complex	284
Magnetic properties of corrole-iron(IV) nitrate complex	287

Magnetic properties of corrole-iron(IV) tosyl complex	289
Magnetic properties of corrole-iron(IV) chlorato complex	292
Part VII Ru	295
Magnetic properties of ruthenium(III) chloro complex with triphenylphosphine and methylamine-bis(<i>p</i> -crisolate) tripodal ligand	297
Magnetic properties of ruthenium(III) chloro complex with triphenylarsine and methylamine-bis(<i>p</i> -crisolate) tripodal ligand	299
Magnetic properties of ruthenium(III) bromo complex with triphenylarsine and methylamine-bis(<i>p</i> -crisolate) tripodal ligand	301
Magnetic properties of ruthenium(III) bromo complex with triphenylphosphine and methylamine-bis(<i>p</i> -crisolate) tripodal ligand	303
Magnetic properties of ruthenium(III) chloro complex with triphenylphosphine and cyclohexylamine-bis(<i>p</i> -cresolate) tripodal ligand	305
Magnetic properties of ruthenium(III) chloro complex with triphenylarsine and cyclohexylamine-bis(<i>p</i> -cresolate) tripodal ligand	307
Magnetic properties of ruthenium(III) bromo complex with triphenylarsine and cyclohexylamine-bis(<i>p</i> -cresolate) tripodal ligand	309
Magnetic properties of ruthenium(III) bromo complex with triphenylphosphine and cyclohexylamine-bis(<i>p</i> -cresolate) tripodal ligand	311
Magnetic properties of ruthenium(III) chloro complex with triphenylphosphine and 2-aminopyridine-bis(<i>p</i> -cresolate) tripodal ligand	313
Magnetic properties of ruthenium(III) chloro complex with triphenylarsine and 2-aminopyridine-bis(<i>p</i> -cresolate) tripodal ligand	315
Magnetic properties of ruthenium(III) bromo complex with triphenylarsine and 2-aminopyridine-bis(<i>p</i> -cresolate) tripodal ligand	317
Magnetic properties of ruthenium(III) bromo complex with triphenylphosphine and 2-aminopyridine-bis(<i>p</i> -cresolate) tripodal ligand	319

Magnetic properties of ruthenium(III) Schiff-base complex containing triphenylphosphine co-ligand	321
Magnetic properties of ruthenium(III) Schiff-base complex containing triphenylarsine co-ligand	323
Magnetic properties of mixed valent diruthenium(II, III) complex with ibuprofen	325
Magnetic properties of mixed valent diruthenium(II, III) complex with aspirin	327
Magnetic properties of mixed valent diruthenium(II, III) complex with naproxen	329
Magnetic properties of mixed valent diruthenium(II, III) complex with indomethacin	332
Part VIII Co	335
Magnetic properties of cobalt(II) nitrate complex with methyl-2-benzimidazolecarbamate	337
Magnetic properties of cobalt(II) complex with furanyl-derived sulfonamide	339
Magnetic properties of cobalt(II) complex with furanyl-derived sulfonamide	342
Magnetic properties of cobalt(II) complex with mesogenic Schiff-base	345
Magnetic properties of cobalt(II) complex with [18]-membered tetra-oxodithiatetraaza macrocyclic ligand	347
Magnetic properties of cobalt(II) complex with [20]-membered tetra-oxodithiatetraaza macrocyclic ligand	349
Magnetic properties of cobalt(II) complex with [18]-membered tetra-oxodithiatetraaza macrocyclic ligand	351
Magnetic properties of cobalt(II) complex with [20]-membered tetra-oxodithiatetraaza macrocyclic ligand	353
Magnetic properties of cobalt(II) complex with quinolone-2-carboxylate	355
Magnetic properties of cobalt(II) complex with sterically hindered <i>o</i>-aminophenol derivative	358
Magnetic properties of cobalt(II) complex with sterically hindered <i>o</i>-aminophenol derivative	360

Magnetic properties of cobalt(II) chloro complex with 2-acetyl-2-thiazoline hydrazone	362
Magnetic properties of cobalt(II) complex with spirocyclophosphazene	364
Magnetic properties of cobalt(II) chloro complex with 2-chlorobenzimidazole	367
Magnetic properties of cobalt(II) bromo complex with 2-chlorobenzimidazole	369
Magnetic properties of cobalt(II) chloro complex with methyl-2-benzimidazolecarbamate	371
Magnetic properties of cobalt(II) bromo complex with methyl-2-benzimidazolecarbamate	373
Magnetic properties of cobalt(II) chloro complex with 2-guanidinobenzimidazole	375
Magnetic properties of cobalt(II) bromo complex with 2-guanidinobenzimidazole	377
Magnetic properties of cobalt(II) nitrato complex with 2-guanidinobenzimidazole	379
Magnetic properties of cobalt(II) chloro complex with 2-phenylbenzimidazole	381
Magnetic properties of cobalt(II) bromo complex with 2-phenylbenzimidazole	383
Magnetic properties of cobalt(II) chloro complex with 2-methylbenzimidazole	385
Magnetic properties of cobalt(II) bromo complex with 2-methylbenzimidazole	387
Magnetic properties of cobalt(II) complex with 2-acetyl-γ-butyrolactone thiosemicarbazone	389
Magnetic properties of cobalt(II) chloro complex with 2-furancarbaldehyde thiosemicarbazone	391
Magnetic properties of cobalt(II) chloro complex with 6-acetyl-1,3,7- trimethylumazine	393
Magnetic properties of cobalt(II) perchlorato complex with 6-acetyl- 1,3,7-trimethylumazine	395
Magnetic properties of cobalt(II) bromo complex with benzimidazole	397

Magnetic properties of cobalt(II) nitrate complex with benzimidazole	399
Magnetic properties of cobalt(II) chloro complex with 2-aminobenzimidazole	401
Magnetic properties of cobalt(II) bromo complex with 2-aminobenzimidazole	403
Magnetic properties of cobalt(II) nitrate complex with 2-aminobenzimidazole	405
Magnetic properties of cobalt(II) chloro complex with albandazole	407
Magnetic properties of cobalt(II) bromo complex with albandazole	409
Magnetic properties of cobalt(II) nitrate complex with albandazole	411
Magnetic properties of cobalt(II) chloro complex with tris(2-benzimidazolylmethyl)amine	413
Magnetic properties of cobalt(II) bromo complex with tris(2-benzimidazolylmethyl)amine	415
Magnetic properties of cobalt(II) nitrate complex with tris(2-benzimidazolylmethyl)amine	417
Magnetic properties of cobalt(II) chloro complex with benzimidazole	419
Magnetic properties of cobalt(II) complex with Schiff-base derived from 2,6-diformyl-4-methylpheol and 5,6-diamino-1,3-dimethyluracil	421
Magnetic properties of cobalt(II) complex with 3-acetylcoumarin-isonicotynoylhydrazone	423
Magnetic properties of imidazole complex of cobalt(II) benzenesulfonate	425
Magnetic properties of cobalt(II) complex with triazine ligand derived from 3-benzyl-7-hydrazinyl-4H-[1,3,4]thiadiazolo[2,3-c][1,2,4]triazin-4-one	427
Magnetic properties of cobalt(II) complex with ligand derived from reduced N,N'-<i>o</i>-phenylenebis(salicylideneimine)	429
Magnetic properties of cobalt(II) complex with ligand derived from reduced N,N'-<i>o</i>-phenylenebis(salicylideneimine)	431

Magnetic properties of cobalt(II) complex with ligand derived from reduced N,N'-<i>o</i>-phenylenebis(salicylideneimine)	433
Magnetic properties of cobalt(II) complex with 1,8-N-bis(3-carboxy)-disalicylidene-3,6-diazaocatane-1,8-diamine	435
Magnetic properties of cobalt(II) complex with sulpiride drug	437
Magnetic properties of dinuclear cobalt(II) complex with 28-membered [2+2] macrocyclic ligand derived from 1,4-bis(2'-formylphenyl)-1,4-dioxabutane and <i>cis</i>-1,2-diaminocyclohexane (as perchlorate salt)	439
Magnetic properties of dinuclear cobalt(II) complex with 28-membered [2+2] macrocyclic ligand derived from 1,4-bis(2'-formylphenyl)-1,4-dioxabutane and <i>cis</i>-1,2-diaminocyclohexane (as nitrate salt)	441
Magnetic properties of binuclear cobalt(II) complex with binucleating Schiff-base ligand with a phenylene spacer	443
Magnetic properties of dinuclear cobalt(II) complex with 'end-off' compartmental Schiff-base ligand	445
Magnetic properties of dinuclear cobalt(II) complex with Schiff-base derived from 4,6-diacetylresorcinol and diethylenetriamine	447
Magnetic properties of diacetato and monohydroxamato bridged dinuclear cobalt(II) complex containing tetramethylethylenediamine	449
Magnetic properties of triacetato bridged dinuclear cobalt(II) urease complex containing tetramethylethylenediamine	452
Magnetic properties of diacetato-bridged, dinuclear cobalt(II) hydrolase complex containing tetramethylethylenediamine	455
Magnetic properties of heterometallic cobalt(II)-copper(II) complex formed by one pot reaction of 2,6-pyridinedicarboxylic acid and ethylenediamine with metal ion	458
Magnetic properties of ion-pair complex having chloro-tris {1-(3,5-dimethylpyrazolyl)methyl}aminecobalt(II) cation	460
Magnetic properties of ion-pair complex having {tris[(2-benzimidazolyl-methyl)amine]chlorocobalt(II)} cation and tetrachlorocobaltate(II) anion	462
Magnetic properties of ion-pair complex having {tris[(2-benzimidazolyl-methyl)amine]bromocobalt(II)} cation and tetrabromocobaltate(II) anion	464

Magnetic properties of tetranuclear cobalt(II) cluster with pyridine alcoholate ligand	466
Magnetic properties of oxalato-bridged cobalt(II) complex with 4-aminopyridine	469
Magnetic properties of oxalato-bridged cobalt(II) complex with 3-aminopyridine	471
Magnetic properties of oxalate-bridged cobalt(II) complex with 2-aminopyridine	473
Magnetic properties of polymeric cobalt(II) complex with pyrazine and pivalic acid	475
Magnetic properties of polymeric cobalt(II) complex with pyrimidine and pivalic acid	478
Magnetic properties of cobalt(III) complex with N,N-bis (2-aminothiophenol)-1,4-bis(carboxylidene phenoxy)butane	481
Magnetic properties of cobalt(III)-cobalt(II) complex formed by one pot reaction of 2,6-pyridinedicarboxylic acid and ethylenediamine with metal ion	483
Part IX Ni	485
Magnetic properties of nickel(II) complex with 28-membered [2+2] macrocyclic ligand derived from 1,4-bis(2'-formylphenyl)-1,4-dioxabutane and <i>cis</i>-1,2-diaminocyclohexane (as nitrate salt)	487
Magnetic properties of nickel(II) complex with imino nitroxide diradical	489
Magnetic properties of nickel(II) complex with sulpiride drug	492
Magnetic properties of nickel(II) complex with 3-acetylcoumarin-isonicotynoylhydrazone	494
Magnetic properties of imidazole complex of nickel(II) benzenesulfonate	496
Magnetic properties of nickel(II) complex with triazine ligand derived from 3-benzyl-7-hydrazinyl-4H-[1,3,4]thiadiazolo[2,3-c][1,2,4]triazin-4-one	498
Magnetic properties of nickel(II) complex with ligand derived from reduced N,N'-<i>o</i>-phenylenebis(salicylideneimine)	500
Magnetic properties of nickel(II) complex with ligand derived from reduced N,N'-<i>o</i>-phenylenebis(salicylideneimine)	502

Magnetic properties of nickel(II) chloro complex with 2-acetyl-2-thiazoline hydrazone	504
Magnetic properties of nickel(II) complex with N(4),N(4)-(butane-1,4-diyl)thiosemicarbazone	506
Magnetic properties of nickel(II) complex with N(4),N(4)-(butane-1,4-diyl)thiosemicarbazone	508
Magnetic properties of nickel(II) chloro complex with N(4),N(4)-(butane-1,4-diyl)thiosemicarbazone	510
Magnetic properties of mixed ligand nickel(II) complex with N-benzyl-N-methyldithiocarbamate and 1,10-phenanthroline	512
Magnetic properties of mixed ligand nickel(II) complex with N,N-dipentyl-dithiocarbamate and 1,10-phenanthroline	514
Magnetic properties of mixed ligand nickel(II) complex with 4-benzyl-piperazinedithiocarbamate and 1,10-phenanthroline	516
Magnetic properties of mixed ligand nickel(II) complex with 4-benzyl-piperazinedithiocarbamate and 1,10-phenanthroline having thiocyanate counteranion	518
Magnetic properties of mixed ligand nickel(II) complex with N,N-benzyl-methyldithiocarbamate and 2,2'-bipyridine	520
Magnetic properties of mixed ligand nickel(II) complex with N,N-dipentyl-dithiocarbamate and cyclam	522
Magnetic properties of mixed ligand nickel(II) complex with N-benzyl-N-methyldithiocarbamate and cyclam	524
Magnetic properties of mixed ligand nickel(II) complex with N,N-dibenzyl-dithiocarbamate and cyclam	526
Magnetic properties of mixed ligand nickel(II) complex with N,N-dibenzyl-dithiocarbamate and 1,10-phenanthroline	528
Magnetic properties of nickel(II) aqua complex with antibacterial drug, flumequine	530
Magnetic properties of mixed ligand nickel(II) complex with 4-benzylpyridine and antibacterial drug, flumequine	532
Magnetic properties of mixed ligand nickel(II) complex with pyridine and antibacterial drug, flumequine	534
Magnetic properties of mixed ligand nickel(II) complex with 2,2'-bipyridine and antibacterial drug, flumequine	536

Magnetic properties of mixed ligand nickel(II) complex with 1,10-phenanthroline and antibacterial drug, flumequinato	538
Magnetic properties of nickel(II) complex with Schiff-base obtained from isatin and 5-methylthiocarbazate	540
Magnetic properties of nickel(II) complex with Schiff-base derived from 2,6-diformyl-4-methylpheol and 5,6-diamino-1,3-dimethyluracil	542
Magnetic properties of nickel(II) complex with 2-phenyl-2-(2-pyridyl)imidazolidine	544
Magnetic properties of nickel(II) azido complex with 2-phenyl-2-(2-pyridyl)imidazolidine	546
Magnetic properties of nickel(II) complex with hexadentate Schiff-base ligand	548
Magnetic properties of nickel(II) azido complex with 1-methyl-2-(naphthyl-α-azo)imidazole	550
Magnetic properties of nickel(II) azido complex with 1-ethyl-2-(naphthyl-α-azo)imidazole	552
Magnetic properties of nickel(II) azido complex with 1-methyl-2-(naphthyl-β-azo)imidazole	554
Magnetic properties of nickel(II) azido complex with 1-ethyl-2-(naphthyl-β-azo)imidazole	556
Magnetic properties of nickel(II) thiocyanato complex with 1-methyl-2-(naphthyl-α-azo)imidazole	558
Magnetic properties of nickel(II) thiocyanato complex with 1-methyl-2-(naphthyl-β-azo)imidazole	560
Magnetic properties of nickel(II) thiocyanato complex with 1-ethyl-2-(naphthyl-α-azo)imidazole	562
Magnetic properties of nickel(II) thiocyanato complex with 1-ethyl-2-(naphthyl-β-azo)imidazole	564
Magnetic properties of mixed ligand nickel(II) complex with 5-(3-pyridyl)-1,3,4-oxadiazol-2-thione and ethylenediamine as co-ligand	566
Magnetic properties of nickel(II) complex with (2-methoxycarbonylmethylimino)-5-methyl-thiazol-3-yl)acetic acid	568
Magnetic properties of mixed ligand nickel(II) complex with hydrotris(3-phenyl-5-methylpyrazol-1-yl)borate, 3-phenyl-5-methylpyrazole and sodium benzoate	570

Magnetic properties of mixed ligand nickel(II) complex with hydrotris(3-phenyl-5-methylpyrazol-1-yl)borate, 3-phenyl-5-methylpyrazole and sodium <i>p</i>-flourobenzoate	573
Magnetic properties of mixed ligand nickel(II) complex with hydrotris(3-phenyl-5-methylpyrazol-1-yl)borate, 3-phenyl-5-methylpyrazole and sodium <i>p</i>-nitrobenzoate	575
Magnetic properties of nickel(II) complex with azo-linked Schiff-base	577
Magnetic properties of nickel(II) complex with azo-linked Schiff-base	580
Magnetic properties of nickel(II) complex with azo-linked Schiff-base	583
Magnetic properties of nickel(II) complex formed by one pot reaction of 2,6-pyridinedicarboxylic acid and ethylenediamine with metal ion	586
Magnetic properties of dinuclear nickel(II) complex with tetra[(1-hydroxyiminomethylnaphthalen-2-yloxy)methyl]ethene	588
Magnetic properties of dinuclear nickel(II) complex with ‘end-off’ compartmental Schiff-base ligand	591
Magnetic properties of binuclear nickel(II) complex with binucleating Schiff-base ligand with a phenylene spaer	594
Magnetic properties of dinuclear nickel(II) complex with 28-membered [2+2] macrocyclic ligand derived from 1,4-bis(2'-formylphenyl)-1,4-dioxabutane and <i>cis</i>-1,2-diaminocyclohexane (as perchlorate salt)	596
Magnetic properties of chloro bridged dinuclear nickel(II) complex with pyridylthioazophenol	598
Magnetic properties of chloro bridged dinuclear nickel(II) complex with pyridylthioazophenol	601
Magnetic properties of chloro bridged dinuclear nickel(II) complex with pyridylthioazophenol	603
Magnetic properties of chloro bridged dinuclear nickel(II) complex with pyridylthioazophenol	605
Magnetic properties of chloro dinuclear nickel(II) complex with pyridylthioazophenol	607

Magnetic properties of azido bridged dinuclear nickel(II) complex with pyridylthioazophenol	609
Magnetic properties of azido bridged dinuclear nickel(II) complex with pyridylthioazophenol	611
Magnetic properties of azido bridged dinuclear nickel(II) complex with pyridylthioazophenol	613
Magnetic properties of azido bridged dinuclear nickel(II) complex with pyridylthioazophenol	615
Magnetic properties of azido dinuclear nickel(II) complex with pyridylthioazophenol	617
Magnetic properties of binuclear nickel(II) complex with 1,4,8-triazacycloundecane	619
Magnetic properties of unsymmetrical macrocyclic dicompartmetnal binuclear nickel(II) complex	622
Magnetic properties of unsymmetrical macrocyclic dicompartmetnal binuclear nickel(II) complex	625
Magnetic properties of unsymmetrical macrocyclic dicompartmetnal binuclear nickel(II) complex	628
Magnetic properties of unsymmetrical macrocyclic dicompartmetnal binuclear nickel(II) complex	631
Magnetic properties of unsymmetrical macrocyclic dicompartmetnal binuclear nickel(II) complex	634
Magnetic properties of $[\text{Ni}_4^{\text{II}}]$ distorted cubane assembly	637
Magnetic properties of mixed ligand nickel(II) complex with tetraaza-macrocycle and isophthalic acid	640
Magnetic properties of mixed ligand nickel(II) complex with tetraaza-macrocycle and 1,3,5-cyclohexanetricarboxylic acid	643
Magnetic properties of nickel(II) complex with 1,5,9-triazacyclododecane	646
Magnetic properties of oxalato-bridged nickel(II) complex with 2-aminopyridine	648
Magnetic properties of oxalato-bridged nickel(II) complex with 3-aminopyridine	650
Magnetic properties of oxalato-bridged nickel(II) complex with 4-aminopyridine	653

Volume 8B

Part X Pt	657
Magnetic properties of ion-pair complex having tetramethylethylenediaminecopper(II) cation and tetracyanoplatinate(II) anion	659
Magnetic properties of tetra- <i>n</i> -butylammonium bis(benzene-1,2-dithiolato-(2-) $k^2(S,S')$ platinate(III)	661
Part XI Cu	663
Magnetic properties of copper(II) complex with clotrimazole	665
Magnetic properties of copper(II) complex with ketoconazole	667
Magnetic properties of copper(II) complex with Schiff-base obtained from isatin and 5-methylthiocarbazate	669
Magnetic properties of copper(II) complex with spirocyclophosphazene	672
Magnetic properties of copper(II) complex with salicylideneanthranilic acid	674
Magnetic properties of mixed ligand copper(II) complex with salicylideneanthranilic acid and 2-methylimidazole	676
Magnetic properties of mixed ligand copper(II) complex with salicylideneanthranilic acid and 2-ethylimidazole	678
Magnetic properties of mixed ligand copper(II) complex with salicylideneanthranilic acid and benzimidazole	680
Magnetic properties of mixed ligand copper(II) complex with salicylideneanthranilic acid and 2-methylbenzimidazole	682
Magnetic properties of copper(II) complex with N,N' - <i>o</i> -phenylenebis(3-ethoxysalicylaldimine)	684
Magnetic properties of 2-nitrobenzoatocopper(II) complex containing N,N -diethylnicotinamide(monomeric)	686
Magnetic properties of copper(II)-saccharinato complex with N -(2-aminoethyl)piperazine	689
Magnetic properties of mixed ligand copper(II) complex with azide and 2-phenyl-2-(2-pyridyl)imidazolidine (as perchlorate salt)	691
Magnetic properties of mixed ligand copper(II) complex with aqua and 2-phenyl-2-(2-pyridyl)imidazolidine (as nitrate salt)	693

Magnetic properties of mixed ligand copper(II) complex with azide and 2-phenyl-2-(2-pyridyl)imidazolidine (as nitrate salt)	695
Magnetic properties of mixed ligand copper(II) complex with acetonitrile and 2-phenyl-2-(2-pyridyl)imidazolidine	697
Magnetic properties of copper(II) complex with pyrazine-2,3-dicarboxylic acid and ethylenediamine	699
Magnetic properties of copper(II) complex with pyrazine-2,3-dicarboxylic acid and 2,2-dimethylpropane-1,3-diamine	701
Magnetic properties of copper(II) complex with 1,2,4-triazolo-[1,5-a]-5,7-dimethylpyrimidine having ethylenediamine as auxiliary ligand	703
Magnetic properties of copper(II) complex with 1,2,4-triazolo-[1,5-a]-pyrimidine having ethylenediamine as auxiliary ligand	705
Magnetic properties of copper(II) complex with 1,2,4-triazolo-[1,5-a]-pyrimidine having 1,10-phenanthroline as auxiliary ligand (as nitrate salt)	707
Magnetic properties of copper(II) complex with 1,2,4-triazolo-[1,5-a]-5,7-dimethylpyrimidine having 1,10-phenanthroline as auxiliary ligand (as perchlorate salt)	709
Magnetic properties of copper(II) complex with 1,2,4-triazolo-[1,5-a]-pyrimidine having 1,10-phenanthroline as auxiliary ligand (as perchlorate salt)	711
Magnetic properties of copper(II) complex with 1-(2-aminoethyl) biguanide	714
Magnetic properties of copper(II) complex with (2-methoxycarbonylmethylimino)-5-methyl-thiazol-3-yl)-acetic acid and pyridine	716
Magnetic properties of mixed ligand copper(II) complex with ethylene-diamine as 5-(3-pyridyl)-1,3,4-oxadiazol-2-thione salt	719
Magnetic properties of copper(II) nitrate complex with 2-hydroxymethyl-benzimidazole	721
Magnetic properties of copper(II) complex with 2-hydroxymethylbenzimidazole	723
Magnetic properties of copper(II) complex with mesogenic Schiff-base	725
Magnetic properties of copper(II) complex of Schiff-base derived from fururylamine	727

Magnetic properties of copper(II) complex of Schiff-base derived from fururylamine	729
Magnetic properties of imidazole complex of copper(II) benzenesulfonate	731
Magnetic properties of copper(II) thiocyanato complex with Schiff-base derived from 2,6-diformyl-4-methylpheol and 5,6-diamino-1,3-dimethyluracil	733
Magnetic properties of copper(II) nitrato complex with Schiff-base derived from 2,6-diformyl-4-methylpheol and 5,6-diamino-1,3-dimethyluracil	735
Magnetic properties of copper(II) perchlorato complex with Schiff-base derived from 2,6-diformyl-4-methylpheol and 5,6-diamino-1,3-dimethyluracil	737
Magnetic properties of copper(II) complex with 6-acetyl-1,3,7-trimethylumazine	739
Magnetic properties of hetero-trihalide (ClBr₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-furaldehyde	741
Magnetic properties of homo-trihalide (BrBr₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-furaldehyde	743
Magnetic properties of hetero-trihalide (ClI₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-furaldehyde	745
Magnetic properties of hetero-trihalide (BrI₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-furaldehyde	747
Magnetic properties of hetero-trihalide (ClBr₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-thiophenecarboxaldehyde	749
Magnetic properties of homo-trihalide (ClI₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-thiophenecarboxaldehyde	751
Magnetic properties of hetero-trihalide (BrBr₂) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-thiophenecarboxaldehyde	753

Magnetic properties of hetero-trihalide (BrI_2) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and 2-thiophenecarboxaldehyde	755
Magnetic properties of hetero-trihalide (ClBr_2) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and pyrrole-2-carboxaldehyde	757
Magnetic properties of hetero-trihalide (ClI_2) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and pyrrole-2-carboxaldehyde	759
Magnetic properties of hetero-trihalide (BrI_2) mononuclear copper(II) complex with Schiff-base derived from diethylenetriamine and pyrrole-2-carboxaldehyde	761
Magnetic properties of copper(II) complex with 2-acetyl-γ-butyrolactone thiosemicarbazone	763
Magnetic properties of copper(II) chloro complex with 2-furancarbaldehyde thiosemicarbazone	765
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 2,2'-bipyridine	767
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 4,4'-dimethylbipyridine	769
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 1,10-phenanthroline	771
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 4-methylphenanthroline	773
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 5-methylphenanthroline	775
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 4,7-dimethyl-1,10-phenanthroline	777
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 5,6-dimethyl-1,10-phenanthroline	779
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 3,4,7,8-tetramethyl-1,10-phenanthroline	781
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 5-phenyl-1,10-phenanthroline	783
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 4,7-diphenyl-1,10-phenanthroline	785

Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 5-chloro-1,10-phenanthroline	787
Magnetic properties of mixed ligand copper(II) complex with acetylacetone and 5-nitro-1,10-phenanthroline	789
Magnetic properties of mixed ligand copper(II) complex with glycine and 1,10-phenanthroline	791
Magnetic properties of mixed ligand copper(II) complex with glycine and 4-methyl-1,10-phenanthroline	793
Magnetic properties of mixed ligand copper(II) complex with glycine and 5-methyl-1,10-phenanthroline	795
Magnetic properties of mixed ligand copper(II) complex with glycine and 4,7-dimethyl-1,10-phenanthroline	797
Magnetic properties of mixed ligand copper(II) complex with glycine and 5,6-dimethyl-1,10-phenanthroline	799
Magnetic properties of mixed ligand copper(II) complex with glycine and 3,4,7,8-tetramethyl-1,10-phenanthroline	801
Magnetic properties of mixed ligand copper(II) complex with glycine and 4,7-diphenyl-1,10-phenanthroline	803
Magnetic properties of mixed ligand copper(II) complex with glycine and 5-chloro-1,10-phenanthroline	805
Magnetic properties of mixed ligand copper(II) complex with glycine and 5-nitro-1,10-phenanthroline	807
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	809
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	811
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	813
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	815
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	817
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	820
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	823

Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	826
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	829
Magnetic properties of copper(II) chlorocomplex with methyl-2-benzimidazolecarbamate	832
Magnetic properties of copper(II) bromocomplex with methyl-2-benzimidazolecarbamate	834
Magnetic properties of copper(II) nitrate complex with 2-methylbenzimidazole	836
Magnetic properties of copper(II) bromo complex with 2-phenylbenzimidazole	838
Magnetic properties of copper(II) chloro complex with 2-guanidinobenzimidazole	840
Magnetic properties of copper(II) bromo complex with 2-guanidinobenzimidazole	842
Magnetic properties of copper(II) nitrate complex with 2-guanidinobenzimidazole	844
Magnetic properties of copper(II) chloro complex with 2-phenylbenzimidazole	846
Magnetic properties of copper(II) chloro complex with 2-methylbenzimidazole	848
Magnetic properties of copper(II) bromo complex with 2-methylbenzimidazole	850
Magnetic properties of copper(II) complex with 1,4-dihydroxy-9,10-anthraquinone-2-sulphonate	852
Magnetic properties of copper(II) complex of Schiff-base containing 2,4-disubstituted thiazole	854
Magnetic properties of copper(II) complex of Schiff-base containing 2,4-disubstituted thiazole	856
Magnetic properties of copper(II) nitrate complex with 2-methyl-4-nitropyridine N-oxide	858
Magnetic properties of copper(II) nitrate complex with 2,6-dimethyl-4-nitropyridine N-oxide	860
Magnetic properties of copper(II) nitrate complex with 2,3,6-trimethyl-4-nitropyridine N-oxide	862

Magnetic properties of copper(II) complex with azo-linked Schiff-base	864
Magnetic properties of copper(II) complex with 15-membered diaza dioxo macrocyclic Schiff-base	867
Magnetic properties of copper(II) complex with 16-membered diaza dioxo macrocyclic Schiff-base	869
Magnetic properties of copper(II) complex with 15-membered diaza dioxo macrocyclic Schiff-base	871
Magnetic properties of copper(II) complex with 15-membered diaza dioxo macrocyclic Schiff-base	873
Magnetic properties of copper(II) complex with N-cycloalkyl-3,5-<i>t</i>Bu₂-salicylaldimine ligand	875
Magnetic properties of copper(II) complex with N-cyclohexyl-3,5-<i>t</i>Bu₂-salicylaldimine ligand	877
Magnetic properties of copper(II) complex with N-cycloheptyl-3,5-<i>t</i>Bu₂-salicylaldimine ligand	879
Magnetic properties of copper(II) complex with N-cyclooctyl-3,5-<i>t</i>Bu₂-salicylaldimine ligand	881
Magnetic properties of copper(II) complex with N-(1-adamantyl)-3,5-<i>t</i>Bu₂-salicylaldimine ligand	883
Magnetic properties of copper(II) complex with N-(2-adamantyl)-3,5-<i>t</i>Bu₂-salicylaldimine ligand	885
Magnetic properties of copper(II) complex with heterocyclic Schiff-base ligand derived from pyrazolone and semicarbazide	887
Magnetic properties of copper(II) complex with heterocyclic Schiff-base ligand derived from pyrazolone and thiosemicarbazide ...	890
Magnetic properties of copper(II) complex with heterocyclic Schiff-base ligand derived from pyrazolone and semicarbazide	893
Magnetic properties of copper(II) complex with heterocyclic Schiff-base ligand derived from pyrazolone and thiosemicarbazide ...	896
Magnetic properties of copper(II) complex with sulpiride drug	899
Magnetic properties of copper(II) complex with 1,4,8,11-tetra (4-nitrobenzoyl)-1,4,8,11-tetraazacyclotetradecane	901
Magnetic properties of copper(II) complex with 1,4,8,11-tetra (2-nitrobenzoyl)-1,4,8,11-tetraazacyclotetradecane	903

Magnetic properties of copper(II) complex with N-benzoylated 1,4,8, 11-tetraazacyclotetradecane	905
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	907
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	909
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	911
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	913
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	915
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	917
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	919
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	921
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	923
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	925
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	927
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	929
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	931
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	933
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	935
Magnetic properties of mixed ligand copper(II) complex with diester and dinitrogen base ligands	937
Magnetic properties of copper(II) complex with 3-acetylcoumarin-isonicotynoylhydrazone	939

Magnetic properties of copper(II) complex with triazine ligand derived from 3-benzyl-7-hydrazinyl-4H-[1,3,4]thiadiazolo[2,3-c][1,2,4]triazin-4-one	941
Magnetic properties of copper(II) complex with 1,8-N-bis(3-carboxy)disalicylidene- 3,6-diazaocatane-1,8-diamine	943
Magnetic properties of dimenic copper(II) complex with clotrimazole	945
Magnetic properties of dimeric copper(II) complex with ketoconazole	947
Magnetic properties of dinuclear copper(II) complex with Schiff-base derived from salicylaldehyde and β-alanine	949
Magnetic properties of dinuclear copper(II) complex with pyridine-2-carbaldehyde thiosemicarbazone and trifluoroacetic acid	952
Magnetic properties of dinuclear copper(II) complex with pyridine-2-carbaldehyde thiosemicarbazone and trifluoroacetic acid	955
Magnetic properties of dinuclear copper(II) complex with (1,6)diazecine-based ligand	958
Magnetic properties of μ-pyrazine bridged dinuclear copper(II) complex with pyridine-2-carbaldehyde thiosemicarbazone	961
Magnetic properties of dinuclear copper(II) complex with pyridine-2-carbaldehyde thiosemicarbazone and dicyanamide	964
Magnetic properties of <i>syn-anti</i> acetate-bridged dinuclear copper(II) complex with 1,10-phenanthroline	967
Magnetic properties of mixed ligand dicopper complex with dibenzodioxo-tetraazamacrocyclic and succinate	969
Magnetic properties of mixed ligand dicopper complex with dibenzodioxo-tetraazamacrocyclic and oxalate	972
Magnetic properties of dinuclear copper(II) complex with deprotonated 2-(5-carboxy-1H-pyrazol-3-yl)-4,4,5,5-tetramethyl-4,5-dihydro-1H-imidazole	975
Magnetic properties of dinuclear copper(II) acetato complex with pyrazol-3-yl-substituted imino nitroxide	978
Magnetic properties of dinuclear copper(II) complex with 5-carboxy-pyrazol-3-yl-substituted nitronyl nitroxide	981

Magnetic properties of chloro bridged dinuclear copper(II) complex with 2-acetyl-2-thiazoline hydrazone	984
Magnetic properties of hydroxo-bridged dinuclear copper(II) complex with saccharinate-N	987
Magnetic properties of hydroxo-bridged dinuclear copper(II) complex with saccharinate-O	990
Magnetic properties of 2-nitrobenzoatocopper(II) complex containing N,N-dimethylformamide	993
Magnetic properties of dinuclear copper(II) complex with pyrazine and 2-phenyl-2-(2-pyridyl)imidazolidine (as perchlorate salt)	995
Magnetic properties of dinuclear copper(II) complex with 28-membered [2 + 2] macrocyclic ligand derived from 1,4-bis(2'-formylphenyl)-1,4-dioxabutane and <i>cis</i>-1,2-diaminocyclohexane (as nitrate salt)	997
Magnetic properties of dinuclear copper(II) complex with pyrimidine derived Schiff-base ligand and 4,4'-bipyridine	999
Magnetic properties of bis(μ-phenoxo) bridged dinuclear copper(II) complex with pyrimidine derived Schiff-base ligand	1002
Magnetic properties of binuclear copper(II) complex with macrocyclic dicompartmental ligand	1005
Magnetic properties of binuclear copper(II) complex with macrocyclic dicompartmental ligand	1008
Magnetic properties of binuclear copper(II) complex with macrocyclic dicompartmental ligand	1011
Magnetic properties of binuclear copper(II) complex with macrocyclic dicompartmental ligand	1014
Magnetic properties of binuclear copper(II) complex with macrocyclic dicompartmental ligand	1017
Magnetic properties of dinuclear copper(II) complex with 1,2,4-triazolo-[1,5-a]pyrimidine	1020
Magnetic properties of dinuclear copper(II) hydrazido complex ligated by hydrotris- (pyrazolyl)borate	1022
Magnetic properties of amidocyanamide bridged dinuclear copper(II) complex with picolinic acid	1024
Magnetic properties of imidazolato-bridged homo-dinuclear copper(II) complex with macrocyclic ligand	1027

Magnetic properties of macrocyclic unsummetrical binuclear copper(II) complex with macrocyclic ligand derived from 2,6-diformylphenol-1,2-bis(aminoxy)- ethane and 1,8-diaminonaphthalene	1030
Magnetic properties of unsummetrical binuclear copper(II) complex with macrocyclic ligand derived from 2,6-diformylphenol-1,2-bis(aminoxy)- ethane and 1,2-diaminobenzene	1033
Magnetic properties of unsummetrical binuclear copper(II) complex with macrocyclic ligand derived from 2,6-diformylphenol-1,2-bis(aminoxy)- ethane and 1,4-diaminobutane	1036
Magnetic properties of unsummetrical binuclear copper(II) complex with macrocyclic ligand derived from 2,6-diformyl-4-methylphenol 1,2-bis(aminoxy)ethane and propylenediamine	1039
Magnetic properties of unsummetrical binuclear copper(II) complex with macrocyclic ligand derived from 2,6-diformyl-4-methylphenol 1,2-bis(aminoxy)ethane and ethylenediamine	1042
Magnetic properties of macrocyclic symmetrical binuclear copper(II) complex derived from mononuclear copper(II) complex having macrocyclic ligand	1045
Magnetic properties of dinuclear copper(II) complex containing 6-(benzylamino)purine as bridging ligand	1047
Magnetic properties of dinuclear copper(II) complex containing 6-(benzylamino)purine as bridging ligand	1050
Magnetic properties of dinuclear copper(II) complex containing 6-(benzylamino)purine as bridging ligand	1052
Magnetic properties of dinuclear copper(II) complex containing 6-(benzylamino)purine as bridging ligand	1055
Magnetic properties of dinuclear copper(II) complex having chloro and 6-(benzylamino)purine as bridging ligand	1058
Magnetic properties of dinuclear copper(II) complex having chloro and 6-(benzylamino)purine as bridging ligand	1061
Magnetic properties of dinuclear copper(II) complex having chloro and 6-(benzylamino)purine as bridging ligand	1064
Magnetic properties of copper(II) complex with coumarin-derived Schiff-base	1067

Magnetic properties of dinuclear copper(II) complex with ‘end-off’ compartmental Schiff-base ligand	1070
Magnetic properties of mixed ligand binuclear copper(II) complex with 1,10-phenanthroline and N,N,N',N'-tetrakis (2-pyridylmethyl)benzene-1,3-diamine	1073
Magnetic properties of chloro-bridged binuclear copper(II) complex with N,N,N',N'-tetrakis(2-pyridylmethyl)benzene-1,3-diamine	1076
Magnetic properties of binuclear copper(II) complex with 1,2,4-triazole based ligand	1079
Magnetic properties of dinuclear copper(II) complex with Schiff-base derived from 4,6-diacetylresorcinol and diethylenetriamine	1082
Magnetic properties of dinuclear copper(II) complex with 1,10-phenanthroline as 4,6-dimethyl-1,2,3-triazolo[4,5-d]pyrimidin-5,7-dionato salt	1084
Magnetic properties of dinuclear mixed ligand copper(II) complex with diester and dinitrogen base ligands	1087
Magnetic properties of dinuclear mixed ligand copper(II) complex with diester and dinitrogen base ligands	1089
Magnetic properties of dinuclear copper(II) complex with tetra [(1-hydroxyiminomethylnaphthalen-2-yloxy)methyl]ethene	1091
Magnetic properties of binuclear copper(II) complex with binucleating Schiff-base ligand with a phenylene spacer	1094
Magnetic properties of imidazolate-bridged hetero-dinuclear copper(II)-zinc(II) complex with macrocyclic ligand	1096
Magnetic properties of copper(II) complex with pyrimidine derived Schiff-base ligand	1099
Magnetic properties of μ-hydroxo- μ-propionato bridged trinuclear copper(II) complex with N-(pyrid-2-ylmethyl) benzenesulfonylamide	1102
Magnetic properties of μ-hydroxo- μ-propionato bridged trinuclear copper(II) complex with N-(pyrid-2-ylmethyl)- <i>p</i>-toluenesulfonylamide	1105
Magnetic properties of μ-hydroxo- μ-propionato bridged trinuclear copper(II) complex with N-(pyrid-2-ylmethyl) naphthalenesulfonylamide	1108

Magnetic properties of trinuclear copper(II) complex with 1,5-bis(di-2-pyridylketone)carbohydrazone	1111
Magnetic properties of copper(II) complex formed by one pot reaction of 2,6-pyridinedicarboxylic acid and ethylenediamine with metal ion	1113
Magnetic properties of tetranuclear pyrophosphate bridged copper(II) complex with 2,2'-dipyridylamine	1115
Magnetic properties of tetranuclear copper(II) acetato complex with pyrazol-3-yl-substituted nitronyl nitroxide	1118
Magnetic properties of phenoxo-bridged tetranuclear copper(II) complex having 2-hydroxy-N-(2-pyridylmethyl)benzamide	1121
Magnetic properties of heterotetranuclear Cu₃Mn complex with N,N'-<i>o</i>-phenylene-bis(3-ethoxysalicylaldehyde)	1123
Magnetic properties of heterotetranuclear Cu₃Co complex with N,N'-<i>o</i>-phenylene-bis(3-ethoxysalicylaldehyde)	1126
Magnetic properties of ion-pair complex containing substituted pentadecanecopper(II) cation and hexacyanoferrate anion	1129
Magnetic properties of hexanuclear copper(II) carboxylate complex with 2,2'-bipyridine	1132
Magnetic properties of heptanuclear Cu₇^{II} complex with N,N'-<i>o</i>-phenylene-bis(3-ethoxysalicylaldehyde)	1134
Magnetic properties of carbonato and hydroxide bridged octanuclear copper(II) complex with 1,10-phenanthroline	1137
Magnetic properties of paradodecatungstate derivative of trinuclear copper(II) azido complex with 2-pyrazine carboxylic acid	1140
Magnetic properties of anionic 4-terpyridone based copper(II) azide supramolecule containing dinuclear-mononuclear co-crystals	1143
Magnetic properties of copper(II) azido cluster with Schiff-base derived from 2,6-diformyl-4-methylphenol and tris(hydroxymethyl)aminomethane	1146
Magnetic properties of dicyanamide bridged polymeric copper(II) complex with di(2-pyridylmethyl)amine	1149
Magnetic properties of copper(II) polymer with alternating $\mu_{1,3}$-squarato and binucleating polypyridylamine bridges	1152

Magnetic properties of copper(II) complex with 5,6-diphenyl-3(2-pyridyl)-1,2,4-triazine	1155
Magnetic properties of copper(II) complex with <i>vic</i>-dioxime ligand containing 1,8-diamino-3,6-dioxaoctane group	1158
Magnetic properties of 2-nitrobenzoatocopper(II) complex containing N,N-diethylnicotinamide(polymeric)	1160
Magnetic properties of copper(II)-saccharinato complex with piperazine	1163
Magnetic properties of polymeric copper(II) complex with Schiff-base derived from salicylaldehyde and β-alanine	1165
Magnetic properties of copper(II) complex with <i>vic</i>-dioxime ligand containing 1,8-diamino-3,6-dioxaoctane group	1168
Magnetic properties of oxalato-bridged copper(II) complex with 4-aminopyridine	1170
Magnetic properties of oxalato-bridged copper(II) complex with 3-aminopyridine	1173
Magnetic properties of oxalato-bridged copper(II) complex with 2-aminopyridine	1175
Magnetic properties of bimetallic oxide Cu-Mo₄ containing <i>p</i>-xylene diphosphonic acid	1177
Part XII Gd	1179
Magnetic properties of quaternary selenide of barium gadolinium and copper	1181
Magnetic properties of quaternary selenide of cesium gadolinium and zinc	1183
Part XIII Tb	1185
Magnetic properties of bipyrimidine-bridged dinuclear terbium(III) complex containing bulky-β-diketonate terminal ligand	1187