

Index

NOTE: Page numbers in *italic* refer to figures. Page numbers in **bold** refer to tables.

- A-lines, lungs, 30, 181
- Abbreviated Cognitive Test for Delirium, **245**
- abxiximab, **131**
- ABCs, ABCDs, 307
 - trauma, 340, 342
- abdomen
 - compartment syndrome, 324, 326–329
 - sepsis, 321–325
 - surgery, NPPV, 198
 - trauma, 343
 - ultrasonography, 31–32
 - X-rays, poisonings, 302
- abdominal aorta
 - aneurysm
 - rupture, 332
 - screening, 103
 - ultrasonography, 30, 31
- abscesses
 - abdominal, 324
 - brain, 464, 465, 478
 - diagnosis, 467, 468
 - mortality, 476
 - treatment, 472–473
 - epidural, 465, 468, 473, 476, 479
- absolute neutrophil count (ANC), low, 400
- AC/AHA classification, heart failure, 139
- acalculous cholecystitis, 393, 424
- ACCOAST trial, 137
- ACE inhibitors, **132**
 - hyperkalemia, 514
 - peri-partum cardiomyopathy, 382
- acetaminophen
 - acute hepatic failure, 360
 - N*-acetylcysteine, 363
 - anti-shivering protocol, 164
 - as antipyretic, 551
 - King's College criteria, 361
 - overdose, 300, 302, **303**
- acetylcholine, **84**
- acid-base balance
 - acute hepatic failure, 362
 - disorders, 537–542
 - diagnosis, 538–541
 - mechanical ventilation, 233–234
 - pregnancy, **378**
 - shock, 152
 - total parenteral nutrition, 53
 - Winter's formula, 302
- acquired immune deficiency syndrome *see* HIV infection
- activated charcoal, 303
 - anticoagulant removal, 557
- activated partial thromboplastin time, 561, 564,
566–567, 570
 - abnormal, 567–568
- ACUITY trial, 137
- acute abdomen, 321–325
- acute coronary syndromes, 123–138, 162–163, 219, 578
 - acute mesenteric ischemia, 334
 - aortic dissection *vs*, 103, 126
- acute exacerbation of COPD, 216–230, 453, 578
 - hospitalization, 222
 - NPPV, 223
 - oxygen therapy, excess, 222
- acute interstitial nephritis, 501, 502
- acute interstitial pneumonia, 208
- acute ischemic stroke, 250, 251, 252–253, 263, 264
 - diagnosis, 254–257
 - presentation, 254
 - treatment, 258–260
- acute kidney injury, 500–506
 - diagnosis, 503–504
 - dopamine, **86**
 - HIV infection, 486
 - prevention, 502
 - prognosis, 506
 - treatment, 504–506
- acute mesenteric arterial thrombosis, 330
- acute mesenteric ischemia, 330–336
- acute myeloid leukemia, tumor lysis syndrome, 586
- acute respiratory distress syndrome, 203–215, 453,
see also respiratory failure
 - diagnosis, 207–209
 - hypoxemia, 169, 205, 207
 - pneumonia, 457
 - prognosis, 213
 - rescue strategies, 209

Mount Sinai Expert Guides: Critical Care, First Edition. Edited by Stephan A. Mayer, Janet M. Shapiro, Umesh K. Gidwani, and John M. Oropello.

© 2021 John Wiley & Sons Ltd. Published 2021 by John Wiley & Sons Ltd.

Companion website: www.wiley.com/go/mayer/mountsinai/criticalcare

- treatment, 209–213
- ventilator settings, 185, 186, 190–191
- acute tubular necrosis, 501, 502
 - prognosis, 506
- acyclovir, **447**
 - encephalitis, 471, 472
- adenosine, 114, 115
- adrenal insufficiency, 155
 - relative, HIV infection, 488
- adrenergic receptors, 84–85, *see also* β -2 adrenergic receptors
- Adrogue–Madias formula, 524, 525
- adsorption, hemoperfusion, 509
- adverse events, ICU-related, 62–67
- aerophagia, 201
- aerosolized antibiotics, 421, 456
- afterload, 82
 - intra-aortic balloon pumps, 91
 - mechanical ventilation on, 192
- agitation
 - dexmedetomidine, 247
 - extubation, 239
 - NPPV, 199
- AIDS *see* HIV infection
- AIMS65 score, gastrointestinal bleeding, 314
- air bronchograms, ultrasonography, 31
- air contamination, arterial blood gases, 174
- air trapping, 178
- airflow obstruction, bronchospastic conditions, 217
- airway pressure release ventilation, 188–189
- airway protection, 235
- airway resistance, 179, 183, 186
- airway (upper), 3–9
 - assessment, 4
 - burns, 350, 352
 - difficult, 7–8, 65–66
 - equipment, 5
 - obstruction, 219, 236
 - pregnancy, 377
 - pressures and resistance, 183
 - trauma, 343
- AKIN criteria, acute kidney injury, 500, **501**
- albumin
 - anion gap, 540
 - as colloid, 496, 497
 - corrected calcium, 531, 532
- albumin–furosemide chaser, 497
- albuterol, **225**, 515
- alcohol consumption
 - limits, 254
 - trauma, 338
- alcohol hand rubs, 414, 415
- alcohol withdrawal, 405
 - fever, 395
- alcoholism
 - ARDS, 206
 - pneumonia, 458
- alcohols, toxic, 541
- aldosterone, 494
- aldosterone : renin ratio, 517
- alkalis, burns, 355
- α -1 adrenergic receptors, drug actions, **84**
- α -2 adrenergic receptors, drug actions, **84**
- alpha angle, capnography, 173
- alteplase, **132**
 - pregnancy, 260
- American Spinal Injury Association (ASIA) impairment scale, 265, 268
- American Thoracic Society/IDSA criteria, ICU admission, 454, 455
- amikacin, **439**, **457**
- aminoglycosides, **439–440**, **457**
- aminotransferases, neutropenic fever, 589
- amiodarone, 114, 115, 119
 - cardiac arrest, 161
- ammonia, acute hepatic failure, 362
- amniotic fluid embolism, 379, 383–384, 572
- amoxicillin/clavulanate, **435**
- amphotericin B, **446**
 - liposomal, **446**, 520
- ampicillin, **435**
- ampicillin-sulbactam, **456**
- AMPLE (trauma history), 339–340, 350
- amyl nitrite, **304**
- amyloid angiopathy, strokes, 251
- anaerobic metabolism, 149
- analgesia, 10–16, 62–63
- analgesedation, 10
- anaphylactoid reactions, hemodialysis, **511**
- anaphylaxis
 - blood transfusions, 548
 - hemodialysis, **511**
- anastomoses, kidney transplantation, 369
- andexanet alfa, 558
 - dosage, **558**
- anemia
 - packed red blood cells for, 546
 - pulse oximetry, 172
 - sepsis, 409
- anesthesia
 - delirium, 244
 - epidural, minimum platelet count, **571**
 - laryngoscopy, 6, 8
- anesthetic agents, 226
 - bronchospastic conditions, 222
 - status epilepticus, 279
- aneurysm, *see also* under abdominal aorta
 - cerebral, 251, 253, 258, 267
- angiitis, CNS, 467
- angiocatheters, 21, 24
- angiodysplasia, gastrointestinal bleeding, 311, 312
- angiography, *see also* CT angiography
 - acute mesenteric ischemia, 333
 - brain death, 296

- angiography (*cont'd*)
 - gastrointestinal bleeding, 315, 316
 - magnetic resonance imaging
 - brain death, 296
 - strokes, 257
 - trauma, 341
- angiotensin I, 494
- angiotensin II, 494
- angiotensin-converting enzyme, 494
 - inhibitors *see* ACE inhibitors
- angiotensin receptor blockers, **132**
 - hyperkalemia, 514
 - peri-partum cardiomyopathy, 382
- angiotensinogen, 494
- anidulafungin, **446**
- animals, diseases from, 466
- anion gap, 540
 - serum, 302
 - urine, 541
- anthrax, meningitis, 474
- antibiotics, 414, 429–448
 - antimicrobial stewardship, 443–444
 - brain abscess, 472–473
 - bronchospastic conditions, 226
 - CAUTIs, 419
 - CLABSIs, 416
 - for *Clostridium difficile* infection, 423
 - Clostridium difficile* infection from, 423
 - fever, 399
 - intravenous albumin with, 497
 - meningitis, **430**, 466, 470, 471–472
 - neutropenic fever, 592
 - pneumonia, **432**, 455–456
 - HIV infection, **487**
 - ventilator-associated, 421, **422**, **432**, 456, **457**
 - sepsis, 409, 410
 - spinal abscess, 473
 - topical, burns, **353**
- antibody induction, kidney transplantation, 370
- anticholinergic toxidrome, **301**, **303**
- anticoagulants
 - acute coronary syndromes, **131–132**, 134
 - acute mesenteric ischemia, 331
 - atrial fibrillation, 254
 - hemorrhage, 132–133, 552–560
 - monitoring, 554, 564
 - OASIS-5 trial, 136
 - renal replacement therapy, 510
 - reversal, 556–558, 571
 - traumatic brain injury and, 273
 - venous thromboembolism, **582**
- anticoat silver mesh, **353**
- antidiuretic hormone, 494, 523–524
- antiepileptic drugs
 - status epilepticus, **279–280**, 281
 - traumatic brain injury, 270
- antifactor Xa, monitoring, 554
- antifungals, **445–446**
- antigen testing, 398
- antimicrobial stewardship, 443–444
- antiplatelet therapy, 125, 130, **131**, 254
 - agents, 554
 - hemorrhage, 132–133
 - gastrointestinal, 312
 - pregnancy and elderly patients, 134
 - reversal, 558
 - traumatic brain injury and, 273
- antipyretics, 399, 551
- antiretroviral therapy (ART), 482, 485–486
- antivirals, **447**
 - encephalitis, 472
- anuria, sepsis, 409
- aortic aneurysm, screening, 103
- aortic dissection, 102–107
 - acute coronary syndromes vs, 103, 126
- apical four chamber position, 28, 34, 79
- apixaban, **582**
- apnea test, 295–296
- apnea time, 5, 6
- apnea trial, 227
- Apo B/Apo A1 ratio, risk of acute coronary syndromes, 124
- Aquacel® Ag, **353**
- ARAS (ascending reticular activating system), 294
- arboviruses, CNS infections, 467, 469, 474
- ARDSnet PEEP/FiO₂ table, 210
- arginine vasopressin, 494
 - organ donors, 297
- arrhythmias, 63, 108–122
 - from central venous access, 65
 - tumor lysis syndrome, **590**
 - vasoactive drugs, 155
- ART (antiretroviral therapy), 482, 485–486
- arterial access, 20–22, 24, 25, 75
 - inadvertent, 19
- arterial blood gases (ABG), 174–177, 538–539, 541
 - acute hepatic failure, 362
 - acute mesenteric ischemia, 333
 - ARDS, 208
 - bronchospastic conditions, 220–221
 - hyperglycemia, 57
 - prone positioning, 211
 - shock, 152
- arterial lines, hemodynamic monitoring, 75–77, 80, 85
- arterial oxygen pressure (PaO₂), *see also* fraction of inspired oxygen (FiO₂)
 - ARDS, 207, 208
 - hypoxemia, 192
 - neurotrauma, 270
 - organ donors, 297
 - pregnancy, 227, 378
 - pulse oximetry, 171–172

- respiratory failure, 169
- symptomatic leukostasis, 591
- arterial reconstruction, liver transplantation, 367
- arteriovenous difference in O₂ content (AVDO₂), 287
- arteriovenous malformations, strokes, 251
- artifacts, ultrasonography, 28
- ASCEND trial, 146
- ascending reticular activating system (ARAS), 294
- ASIA impairment scale, 265, 268
- aspiration (pulmonary), 8–9, 452
 - prevention, 206
 - risk factors, 4
 - stroke patients, 259
- aspirin
 - for acute coronary syndrome, 130, **131**
 - coagulation parameters, 567
 - gastrointestinal bleeding, 312
 - low dose, 125, 254
 - time to hemostasis after stopping, 559
- asplenia, sepsis, 400
- assist control, mechanical ventilation, 187
- assisted wave, intra-aortic balloon pumps, 91
- asthma, 191, 216–230, 578
 - hospitalization, 222
 - NPPV, 197, 199
- asystole, 159
- atelectasis
 - ARDS, 204
 - ventilation waveforms, 182
 - ventilator-associated events vs, 421
- atherosclerosis, 123–124, 138
 - acute mesenteric ischemia, 331
 - strokes, 251
- atony, uterus, 381
- ATP, phosphate depletion, 521
- atrial fibrillation, 110
 - acute mesenteric ischemia, 331
 - treatment, 114, 254
- atrial flutter, misdiagnosis, 113
- atrial natriuretic peptide, 494
- atrial tachycardia, 115, 117, 119
- atrioventricular block, 109, 117, 120
 - diagnostic algorithm, 112
 - treatment algorithm, 116
- atrioventricular dissociation, 109, 111
- atrioventricular nodal re-entrant tachycardia, 114–115, 121
- atrioventricular re-entrant tachycardia, 114–115
- atropine, 113, **306**
- attention, assessment of, **245**
- atypical symptoms, acute coronary syndromes, 130
- augmented wave, intra-aortic balloon pumps, 91
- auto-PEEP, 178, 182, 191, 192, 217, 230
 - examination, 220
 - management, 224
 - pressure-regulated volume control, 188
- autocycling, ventilators, 186, 188
- autoimmune diseases
 - encephalitis, 467
 - lungs, 453
- autoregulation, cerebral, 266, 285, 291
- avibactam, **436**
- awake look, difficult airway, 7
- axial flow pump, Impella, 91, 92, 99
- axillary artery, 21, 25, 75
- azithromycin, **438, 456**
- azoles, **445**
- aztreonam, **437**
- B-lines, lungs, 30, 181
- B-mode ultrasonography, 27
- B-type natriuretic peptide, 140, 142, 143, 580
- bacitracin, **353**
- bacterial DNA, meningitis, 465
- bactericidal activity, 433
- bacteriology, 397
- bacteriostatic activity, 433
- bacteriuria, urinary catheter-associated, 393
- bag and mask ventilation, 6
 - difficult airway, 7
- barcode sign, 30, 35
- barotrauma, 192, 205, 211, 227, 230, *see also*
 - pneumothorax
 - examination, 220
 - ventilation waveforms, 182
- basic metabolic profile, trauma, 341
- basiliximab, 370
- basophilia, causes, **391–392**
- Battle's sign, 268
- Baux score, 355
- bedside percutaneous dilational tracheostomy, 42–48
- bedside shivering assessment scale, 165
- bedside ultrasonography, 26–36
 - pulmonary embolism, 580
- behavioral pain scale, 12
- benign postoperative fever, 394
- benzodiazepines, 14–15
 - cholinergic toxidrome, **306**
 - delirium, 246
 - NPPV, 200
 - overdose, **304**
 - pregnancy, 228
 - status epilepticus, 278
- Berlin definition, ARDS, 207
- β-1 adrenergic receptors, drug actions, **84**
- β-2 adrenergic receptors
 - agonists, 225
 - pregnancy, 227
 - drug actions, **84**
- beta-2 transferrin, 269

beta-blockers

acute coronary syndromes, 125, **132**aortic dissection, **105**

atrial fibrillation, 114

gastrointestinal bleeding, 313

overdose, **304**

peri-partum cardiomyopathy, 382

sepsis, 411

 β -D-glucan, 398

beta-human chorionic gonadotropin, 341

beta-lactam/beta-lactamase inhibitors, 442

beta-lactamase, organisms producing, 442, 443

beta-lactams, pneumonia, **456, 457**

bicarbonate (ion), 537, 539, 540

pregnancy, **378**

bilevel positive pressure ventilation, 188, 196

complications, 66

biliary reconstruction, liver transplantation, 367

bilirubin, SOFA score, **407**Biobrane®, **353**

biofilms, 418

biopsy

endomyocardial, 143

minimum platelet count, **571**

status epilepticus, 278

bivalirudin, **131, 137**bladder drainage, pancreas transplantation, **373**

bladder pressure, 503

bladder view, ultrasonography, 31, 32, 36

blast crisis, 586

blast injury, 342

Blatchford score, 314

bleeding see hemorrhage

blisters, burns, 352

blood cultures, 397

CLABSI, 416

CNS infections, 469

neutropenic fever, 591

pneumonia, 454

sepsis, 404

blood gases, see also arterial blood gases

venous, 77, 175–176

blood glucose, 60, 61

blood loss

burns surgery, 354

hypotension, 314

obstetric, 380–381

trauma, 342

blood pressure, see also mean arterial pressure

children, hemorrhage, 344

intra-aortic balloon pumps, 91, 98

intracranial pressure management, 287

monitoring, 75, 80

organ donors, 297

pre-eclampsia, 385

screening, 103

shock, 152, 154

blood product transfusion, 545–547

ARDS, **206**

fever from, 550–551

trauma, 342

blood transfusion, 63–64

acute coronary syndromes, 133

acute hepatic failure, 362

ARDS, 206

delayed complications, 551

gastrointestinal bleeding, 317

lung injury, 207

massive, 547

reactions, 395, **396**, 548–551

blood urea nitrogen, shock, 152

body mass index, pancreas transplantation criteria, 371

body weight

predicted, 185

total parenteral nutrition, 51

bone marrow biopsy, minimum platelet count, **571**

bone marrow suppression, 564

botulinum toxicity, **304**

brachial artery, 75

bradyarrhythmias, 108, 110

treatment, 113

brain, see also central nervous system; intracranial

pressure; traumatic brain injury

abscess, 464, 465, 478

diagnosis, 467, 468

mortality, 476

treatment, 472–473

after cardiac arrest, 165–166

infarction, 252

oxygenation monitoring, 287

brain death, 293, 294, 295–296

psychosocial issues, 297–298

brain natriuretic peptide, 494

brain tissue oxygen partial pressure (PbtO₂), 287

branched chain amino acids, 53

breath sounds, 170

Bristol stool scale, 422

broad complex tachyarrhythmias, 108, 110,

111, 122

diagnostic algorithm, 113

treatment algorithm, 117

bromocriptine, **305, 383**

bronchial tree, 41

bronchiectasis, 219

bronchoalveolar lavage, *P. jirovecii* pneumonia,

483, 490

bronchodilators, 222, 225

bronchoscopes, 37

bronchoscopy, 37–41

bacteriology, 397

bedside percutaneous dilational tracheostomy,

42, 46

bronchospastic conditions, 225

HIV infection, 483

- minimum platelet count, **571**
 - NPPV, 198
 - pneumonia, **457**
- bronchospasm, **175**, *see also* asthma; status asthmaticus
 - auto-PEEP, 224
- Brudzinski's sign, 468
- Brugada's sign, 111
- bundle branch block, 122
- burns, 347–358
 - diagnosis, 349–351
 - evolution of injury, 348
 - prevention, 349
 - prognosis, 355–356
 - severity classification, 351, 358
 - treatment, 351–355
- bupirone, anti-shivering protocol, 164
- C peptide, pancreas transplantation criteria, 371
- C-type natriuretic peptide, 494
- Cairo-Bishop grading, tumor lysis syndrome, **590**
- calcitonin, **531**
- calcitriol, 530
- calcium acetate, **520**
- calcium channel blockers
 - atrial fibrillation, 114
 - complications, 115
 - overdose, **304**
 - sepsis, 411
- calcium chloride, **304**, 515, 533, **534**
- calcium citrate, 534
- calcium gluconate, **304**, 533, 534
- calcium (level)
 - arrhythmias, 111
 - corrected, 531, 532
 - tumor lysis syndrome, 589
- calculated osmolality, 302
- caloric requirements, total parenteral nutrition,
 - 51, **52**
- cancer
 - emergencies, 586–595
 - gastrointestinal bleeding, 311
 - hypercalcemia, 530
 - paraneoplastic encephalitis, 467
- candesartan, 382
- Candida* (spp), **430**
 - CLABSIs, 417
- candiduria, urinary catheter-associated, 393
- cangrelor, **131**, 137
- capillary refill, 406
- capnography, 172–174, **175**, 182
- captopril, 382
- carbapenem-resistant Enterobacteriaceae, 442–443
- carbapenemases, 443
- carbapenems, **436–437**, 442, 443, **457**
- carbohydrates, total parenteral nutrition, **52**
- carbon dioxide, *see also* end-tidal CO₂
 - blood values, 176, 537, 539
 - apnea test, 296
 - extracorporeal removal, 222
 - pregnancy, 378
 - respiratory failure, 170
 - Winter's formula, 302
- carbon monoxide poisoning, **304**
- carboxyhemoglobinemia, **304**
 - arterial blood gases, 177
- cardiac arrest, 159–166
 - advanced life support, 161–162
 - capnography, 174
 - capnography waveform, **175**
 - fever after, 400
 - on mechanical circulatory support, 94
 - pregnancy, 165, 378–380
 - prognosis, 165–166
- cardiac failure *see* heart failure
- cardiac output, 82, 149
 - abdominal compartment syndrome, 326–327
 - arterial line monitoring, 76–77
 - LIDCO system, 76
 - pregnancy, **378**
 - pulmonary artery catheter monitoring, 78
- cardiac surgery, NPPV, 198
- cardiac troponins, 129, 580
- cardiac ultrasonography (TTE), 28–29, 34, 63, 78–79,
 - see also* transesophageal echocardiography
 - aortic dissection, 104
 - ARDS, 209
 - arrhythmias, 111
 - heart failure, 143
 - pregnancy, 379
 - pulmonary embolism, 181, 580
 - shock, 153
- cardiogenic pulmonary edema
 - ARDS vs, 207
 - NPPV, 197
 - ultrasonography, **31**
- cardiogenic shock, 83, 148–149, 150, 158
 - peripartum, 156
 - ultrasonography, 33
- cardiomyopathy
 - peri-partum, 156, 379, 381–383
 - signs, 111
- cardiopulmonary resuscitation, 160–161
 - capnography, 174
- cardiotoxicity, screening for, 140
- cardioversion, 114
- CardioWest™ total artificial heart, 97
- carina, 39
- carotid artery stenosis, prophylaxis, 254
- carotid sinus pressure, 121
- caspofungin, **446**
- catabolic state, 53
- catheter-associated urinary tract infections (CAUTIs), 67,
 - 393, 397, 417–419
- catheter-based axial flow pump, Impella, 91, 92, 99

- catheter tip cultures, 397, 416
- catheters, 22
 - arterial access, 24
 - central venous access, 18
 - external ventricular drainage (EVD), 286
 - infections, 155, 392–393
 - total parenteral nutrition, 53
- caustic burns, 355
- caval replacement, liver transplantation, 367, 375
- cavernous sinus thrombosis, 473
- CD4 cell counts, HIV infection, **481**
- cefazolin, **435**
- cefepime, **436, 457**
- cefotaxime, pneumonia, **456**
- cefoxitin, **436**
- cefpodoxime, **436**
- ceftaroline, **436**
- ceftazidime, **436, 457**
- ceftazidime-avibactam, 442, 443
- ceftolozane, **436**
- ceftolozane-tazobactam, 442
- ceftriaxone, **436, 456**
- cellulitis, 578
- Center to Advance Palliative Care (CAPC), guidelines, 69
- central line-associated bloodstream infections (CLABSIs), 66–67, 415–417, 419, **430**
- central lines, 76–77
- central nervous system, *see also* brain
 - fever, 400
 - infections, 394, 463–479, *see also* encephalitis; meningitis
 - monitoring, 284–292
 - sepsis, **405**
- central pontine myelinolysis, 65, 526
- central transtentorial herniation, 286
- central venous access, 17–20, 23, 24, *see also* intravascular devices
 - complications, 65
 - minimum platelet count, **571**
 - oxygen measurement, 157
 - total parenteral nutrition, 52
 - vasoactive drugs, 85
- central venous blood gases, 175, 176
- central venous pressure, 76, 77, 80
- centrifugal flow pumps, 94
- CentriMag extracorporeal pump, 93, 100
- cephalosporins, **435–436, 457**
- cerebellar herniation, 286
- cerebral blood flow, 266, 285, 288, 291
 - brain death, 296
- cerebral edema, 252, 258, 285
 - hyponatremia, 526
- cerebral metabolic rate of O₂ consumption (CMRO₂), 287
- cerebral microdialysis, 288–289
- cerebral perfusion pressure, 266, 285, 291
 - acute hepatic failure, 362
 - children, 273
 - cerebral venous sinus thrombosis, 252
- cerebrospinal fluid
 - CNS infections, 469
 - HIV infection, 484
 - leaks, 466
 - molecular testing, 398
 - pressure, cryptococcal meningitis, 487–488
 - status epilepticus, 278
- cerebrovascular accidents *see* strokes
- cervical spine
 - bedside percutaneous dilational tracheostomy, 44
 - difficult airway, 8
 - trauma, 342
- Cesarean delivery
 - peri-mortem, 380
 - trauma patients, 344
- CHAMPION-PHOENIX trial, 137
- charcoal *see* activated charcoal
- checklist, lung injury prevention, 206
- chemical burns, 355
- chest compressions, 160–161
 - on mechanical circulatory support, 94
- chest pain, 126–127
 - chest X-ray, 129
- chest-tube drainage, pneumonia, **457**
- chest wall rigidity, fentanyl, 12
- chest X-ray
 - after bronchoscopy, 40
 - ARDS, 209, 214
 - barotrauma, 230
 - bedside percutaneous dilational tracheostomy, 46
 - bronchospastic conditions, 221
 - chest pain, 129
 - diaphragmatic dysfunction, 180
 - heart failure, 147
 - HIV infection, 484
 - pneumonia, 398, 454, 461–462
 - Pneumocystis jirovecii*, 490
 - ventilator-associated, 421
 - respiratory monitoring, 181
 - shock, 153
- Cheyne Stokes respiration, 170
- children
 - acute abdomen, 324
 - brain death, 298
 - burns, 354
 - rule of nines, 357
 - cardiac arrest, 165
 - CNS infections, 475
 - gastrointestinal bleeding, 318
 - neurotrauma, 273
 - pneumonia, 458
 - status epilepticus, 282
 - trauma, 344
- chlorhexidine, 67, 415, 416
- chloride, in therapeutic fluids, **497**
- cholecystitis

- acalculous, 393, 424
- ultrasonography, 398
- cholestasis, 393
 - total parenteral nutrition, 54
- cholinergic toxidrome, **301, 306**
- chronic obstructive pulmonary disease, 191, 216–230,
 - see also acute exacerbation of COPD
 - NPPV, 196–197, 199
- Chvostek sign, 533
- cidofovir, pneumonia, children, 458
- cilostazol, 254
- ciprofloxacin, **437, 457**
- cisatracurium, bronchospasm, **225**
- citrate anticoagulation, regional, 510
- clarithromycin, **456**
- Clark electrode, 287
- claustrophobia, NPPV, 200
- clindamycin, **438**
- clinical tumor lysis syndrome, 589
- clipping, cerebral aneurysm, 258
- clips (hemoclips), 316
- clopidogrel, **131**, 136, 254
 - time to hemostasis after stopping, 559
- Clostridium difficile*, 393, 397, 414, 415, 422–426
 - antibiotics for, **431**
- clot retrieval, strokes, 258
- clotting factors, see also factor Xa
 - blood transfusion on levels, 547
 - deficiency, 562, 565, **572**
 - plasma products, 546
 - vitamin K-dependent, 553
- co-oximetry, 176–177
- coagulopathies, 561–574
 - acute hepatic failure, 362
 - acute mesenteric ischemia, 331, 333
 - CNS infections, 473
 - diagnosis, 565–570
 - gastrointestinal bleeding, 312, 566
 - hypercoagulable state, 576
 - prevention, 564–565
 - renal replacement therapy, **511**
 - strokes, 251
 - treatment, 570–573
- cocaine poisoning, **304**
- coiling, cerebral aneurysm, 258
- coinfections, HIV infection, 484
- colistin, **439**
 - inhaled, 442
- colitis
 - bleeding, 311
 - ischemic, 312, 332, 424
 - pseudomembranous, 423
- colloids, 342, 350, 496–497
- colonization, pathogens, 450
- colonoscopy, gastrointestinal bleeding, 315, 316
- color Doppler ultrasonography, 27
- colorimetric capnography, 172
- Columbia anti-shivering protocol, 164
- coma, 293–298
 - CNS infections, 473
 - EEG, 290, 292
- coma vigil, 293
- common femoral vein, ultrasonography, 32
- community-acquired pneumonia, 449
 - antibiotics, 455, **456**
- compartment syndrome, abdominal, 324, 326–329
- compliance
 - intracranial, 284, 291
 - lungs, 179, 186, 209
- complications, ICU-related, 62–67
- compression : ventilation ratio, 161
- computed tomography, see also CT angiography
 - abdominal compartment syndrome, 328
 - ARDS, 209, 214
 - CNS infections, 469
 - fever, 398
 - hemorrhage, 556
 - pleural effusion, 462
 - Pneumocystis jirovecii* pneumonia, 490
 - pulmonary embolism, pregnancy, 584
 - shock, 153
 - status epilepticus, 278
 - strokes, 257
 - trauma, 341
 - brain injury, 269
 - pregnancy, 344
- concentration dependent activity, antibiotics, 433, 434
- Confusion Assessment Method for the ICU, **245**
- congestion, heart failure, 142
- congestive hepatopathy, 361
- connective tissue diseases, lungs, 453
- consciousness, disorders of, 293
- consent, HIV testing, 482
- consolidation (pulmonary)
 - chest X-ray, 461
 - clinical signs, 453
 - ultrasonography, 31, 36, 181, 461, 462
- consumption coagulopathies, 563, **564**
- context-sensitive half-time, 10
- continuous glucose monitoring, 57
- continuous positive airways pressure, 196, 234
- continuous veno-venous hemodiafiltration, 509
- continuous veno-venous hemodialysis, 509, 512
- continuous veno-venous hemofiltration, 509
 - energy requirements, **51**
- contrast-induced nephropathy, 65
- convulsive status epilepticus, 275
- cooling phase, TTM, 164
- cooling systems, for fever, 400
- copper, liver failure, 53
- core body temperature, 389
- coronary angiography, 129

- coronary artery bypass grafting, **132**, 137
- corrected calcium, 531, 532
- corticosteroids
 - asthma and COPD, 222, 226
 - HIV infection, 488
 - meningitis, 470, 472
 - neurotrauma and, 271
 - pneumonia, **456**, 457
 - pregnancy, 227
 - septic shock, 411
- costs, trauma, 337
- coumadin, **582**
 - coagulation parameters, 567
- coumarins see vitamin K antagonists
- craniotomy, intracerebral hemorrhage, 258
- CRASH-2 trial, tranexamic acid, 346
- creatinine
 - acute kidney injury, 486, 500, **501**
 - Clostridium difficile*, 425, **431**
 - pre-eclampsia, 385
 - pregnancy, 505
 - shock, 152
 - SOFA score, **407**
 - thrombocytopenia, 570
 - tumor lysis syndrome, 589, **590**
- creatinine clearance
 - neutropenic fever, 589
 - tubular reabsorption of phosphate, 520
- cricoid pressure, 7
- cricothyroidotomy, 343
- critical care pain observation tool, 12
- critical illness myopathy, 63
- critical illness polyneuropathy, 63
- cryoprecipitate, 546
 - sepsis from, 549
- cryoprecipitate-reduced plasma, 546
- Cryptococcus neoformans*, 475, 487–488
- cryptogenic organizing pneumonia,
 - 208, 453
- crystalloids, 496–497
- CT angiography
 - acute coronary syndromes, 129
 - acute mesenteric ischemia, 333
 - brain death, 296
 - bronchospastic conditions, 221
 - gastrointestinal bleeding, 315, 316
 - heart failure, 143
 - neck, 269
 - pulmonary embolism, 585
 - strokes, 257
 - venous thromboembolism, 580
- cuff leak tests, 8, 236
- CURB-65 score, 454, 455
- CURE trial, 136
- curvilinear probes, ultrasonography, **27**, 34
- Cushing's disease, 56
- Cushing's syndrome, exclusion, 517
- Cushing's triad, 285, 361
- cyanide poisoning, 302, **304**
- cyanosis, 170
- cycle phase, mechanical ventilation, 187
- cyclic lipopeptides, **441**
- cyproheptidine, **306**
- cystic fibrosis, inhaled antibiotics and, 442
- cysticercosis, 475
- cysts, renal, 32
- cytochrome P450, pregnancy, **378**
- cytokines
 - bronchospastic conditions, 217
 - meningitis, 465
- cytomegalovirus, 397, 400, 474
- cytoreductive therapy, 592
- cytotoxic edema, cerebral, 285
- D-dimer
 - acute mesenteric ischemia, 333
 - aortic dissection and, 103, 104
 - venous thromboembolism and, 580
- dabigatran, reversal, 557, **558**
- dacrocytes, 570
- daily dialysis, 509
- daily weaning protocols, 237
- dalteparin, 384
- damage control, trauma, 343, 344
- daptomycin, **441**
- DC cardioversion, 114, 115
- de-escalation
 - antibiotics, 444
 - treatment for shock, 154
- death see brain death; palliative care
- decerebrate posturing, 267, 294
- decorticate posturing, 267, 294
- deep vein thrombosis see venous thromboembolism
- deferaxamine, **305**
- defibrillators
 - in cardiac arrest, 159, 161
 - withdrawal, 71
- dehydration, 405
- delayed cerebral ischemia (DCI), monitoring, 289
- delayed hemolytic transfusion reaction, 395
- delirium, 243–249
- Delirium Detection Score, **245**
- delta-delta gap, 540, **542**
- delta ratio, 540
- demyelination, central pontine myelinolysis, 65
- depth electroencephalography, 289
- desmopressin
 - antiplatelet agent reversal, **558**
 - hemorrhage on renal replacement therapy, **511**
 - hyponatremia, 526
- dexamethasone, meningitis, 470, 472, 473
- dexamethasone suppression test, 517
- dexmedetomidine, 15, 62, 244
 - anti-shivering protocol, 164

- delirium, 247
- NPPV, 200
- dextrose
 - for hyperkalemia, 515
 - for hyponatremia, 529
 - overdose patients, 307
 - beta-blockers, **304**
 - solutions, 497
- dextrose (50%), **306**
- diabetes insipidus, 528
 - organ donors, 297
- diabetes mellitus, 55, 56, 57
 - acute mesenteric ischemia, 331
 - risk of acute coronary syndromes, 124
- dialysis, *see also* hemodialysis; renal replacement therapy
 - acute kidney injury, 504
 - disequilibrium, **511**
- diaphragm
 - atrophy, 232, 236
 - function, 180
- diarrhea
 - Clostridium difficile*, 393, 422
 - infectious, 424
- diastolic heart failure, 140, 144
- diazepam, 14–15
 - cholinergic toxidrome, **306**
 - status epilepticus, **279**, 281
- Dieulafoy vascular malformations, 312
- diffuse alveolar hemorrhage, 208
- diffuse esophageal spasm, acute coronary
 - syndromes vs, 126
- digibind, **304**
- digifab, **304**
- digoxin, 114, 115
 - poisoning, **304**
- dilational tracheostomy, bedside percutaneous,
 - 42–48
- dilators, tracheostomy, 46
- diltiazem, 114–115
- dilutional thrombocytopenia, coagulation
 - parameters, 567
- dimercaprol, **305**
- dipstick test, urine, 397
- direct oral anticoagulants, 552
 - monitoring, 554
- direct thrombin inhibitors, 553
 - coagulation parameters, 567
 - monitoring, 554
 - reversal, 557
 - time to hemostasis after stopping, 559
- discitis, 467
- dissection, *see also* aortic dissection
 - hepatic artery, liver transplantation, 367
 - intracranial, 251
- disseminated intravascular coagulation (DIC), 555, 562,
 - 566, **571**
- distributive shock, 83, 158
 - ultrasonography, 33
- diuretics, *see also* furosemide
 - acute kidney injury, **505**
 - heart failure, 143, 144, 146, 383
 - potassium-sparing, 518
- diverticulosis, bleeding, 311, 312, 318
- dobutamine, 85, **86–87**
 - heart failure, **87**, 144
 - post-partum cardiomyopathy, 383
 - pulmonary embolism, 581
 - shock, **155**
- donor criteria, pancreas transplantation,
 - 371–372
- dopamine, **84**, 85, **86**
 - bradyarrhythmias, 113
 - heart failure, **144**
 - hypophosphatemia, 521
 - shock, **155**
- DOSE trial, 146
- doxycycline, **438**, **456**
- dressings
 - Aquacel® Ag, **353**
 - central venous access, 19
- Dressler's syndrome, 396
- driving pressure, 209–210
- drug-induced thrombocytopenia, 562, 565, **572**
- drug interactions, antiretroviral therapy, 485
- drug overdose, 299–308
 - ARDS, 206
- drug reactions, 299–308
 - acute hepatic failure, 360
- drug-related fever, 394–395
- drug-related pulmonary toxicity, 453
- dual control pressure ventilation mode, 188
- duodenal ulcers, 312
- duodeno-duodenostomy, pancreas transplantation, **373**
- durotomy, 258
- dynamic compliance, 186
- dynamic hyperinflation, 217, 227
- dynamic intracavitary gradient, phenylephrine for, **87**
- dysoxia, 148

- ear lobe, pulse oximetry, 171, 172
- echinocandins, **446**
- echocardiography *see* cardiac ultrasonography
- eclampsia, 384–385
- eculizumab, **572**
- edema, 495, *see also* cerebral edema; pulmonary edema
 - heart failure, 142
- edetate calcium disodium, **305**
- elderly patients
 - acute abdomen, 324
 - acute coronary syndromes, 134
 - acute kidney injury, 506
 - ARDS, 213
 - CNS infections, 475
 - sepsis, 411

- elderly patients (*cont'd*)
 - shock, 156
 - strokes, 260
 - trauma, 345
- electric cardioversion, 114, 115
- electrical burns, 354–355, 358
- electrical storm, cardiac, 115
- electrocardiography, 63
 - acute coronary syndromes, 129
 - cardiac arrest patients, 163
 - heart failure, 143
 - hyperkalemia, 515
 - pulmonary embolism, 126, 579
 - ventricular tachycardia, 111
- electroencephalography, 276
 - after cardiac arrest, 165
 - brain death, 296
 - coma prognosis, 290, 292
 - continuous monitoring, 289–290
 - intracortical, 289
 - status epilepticus
 - non-convulsive, 277
 - refractory, 281
- electrolytes
 - disorders, 64–65, 513–536
 - total parenteral nutrition, 50, 52
- elimination half-time, 10
- embolectomy, *see also* pulmonary embolectomy
 - acute mesenteric ischemia, 333
- embolism, *see also* pulmonary embolism
 - acute mesenteric arterial, 330, 331
 - amniotic fluid, 379, 383–384, 572
 - cerebral, 251, 264
- embolization (transcatheter), gastrointestinal bleeding, **317**
- empyemas, 457
- enalapril, 382
- enalaprilat, aortic dissection, **105**
- encephalitis, **405**, 463–479
 - cerebrospinal fluid, 398
 - diagnosis, 467
 - imaging tests, 278
 - herpes simplex, 464, 472, 474, 476, 478
 - prognosis, 476
- encephalopathy
 - hepatic, 361
 - enteral protein, 363
 - lead poisoning, **305**
 - metabolic, 467
- end-expiratory positive airway pressure, 196, 199
- end-of-life care *see* palliative care
- end-organ dysfunction, shock, 152, 154
- end-tidal CO₂, 172, 173
 - alveolar vs arterial, 174
- endotracheal intubation, 6
- endocarditis, **405**
- endomyocardial biopsy, 143
- endoscopy
 - biopsy, minimum platelet count, **571**
 - gastrointestinal bleeding, 313, 315, 316, 318
- endotracheal intubation, 6–7
 - bronchospastic conditions, 223–224
 - burns, 352
 - capnography waveform, **175**
 - complications, 65–66
 - end-tidal CO₂, 6
 - gastrointestinal bleeding, 66
 - neurotrauma, 271
 - obesity, 7
 - trauma, 340, 343
 - vomiting, 66
- endotracheal tubes, 5
 - bronchoscopy and, 38, 39
 - leak test, 8, 236
 - migration of bacteria round, 420
 - obstruction, 239, 240
 - patency, 190
 - resistance of, 235
 - subglottic secretion drainage ports, 452
 - trauma from, 8
- energy density, enteral nutrition, 50
- energy requirements, TPN, 51, **52**
- enoxaparin, 64, **131**, 136, 137, 384, 577
- enteral nutrition, 50, 64
- enteric drainage, pancreas transplantation, **373**
- enteric gram-negative bacteria, pneumonia risk factors, 451
- enteroatmospheric fistula, 329
- enteroviruses, CNS infections, 474
- eosinophilia, causes, **391–392**
- eosinophilic pneumonia, 207, 453
- epicardial fat pad, 29
- epidural abscess, 465, 468, 473, 476, 479
- epidural anesthesia, minimum platelet count, **571**
- epidural hematoma, 271
- epiglottis, 3
- epinephrine, **85**, **144**
 - bradyarrhythmias, 113
 - bronchospasm, 225
 - cardiac arrest, 161
 - nebulized treatment, 238
 - shock, **155**
 - transfusion reactions, 548
- epistaxis, LV assist devices, 96–97
- eptifibatide, **131**
- ertapenem, **437**
- erythrocyte studies, gastrointestinal bleeding, 315, 316
- erythrocytosis, 152
- erythromycin, 316
- escape rhythm, 109
- ESCAPE trial, 146
- escharotomy, **353**

- esmolol
 - aortic dissection, **105**
 - atrial fibrillation, 114
- esophageal rupture, aortic dissection vs, 103
- esophageal spasm, acute coronary syndromes vs, 126
- estrogen, hemorrhage on renal replacement therapy, **511**
- ethylene glycol poisoning, **305**
- exercise, 254
- exercise stress testing, 129
- exocrine drainage, pancreas transplantation, **373**
- extended-spectrum beta-lactamase, 443
 - organisms producing, 442
- external ventricular drainage (EVD) catheter, 286
- extracellular fluid, 495
 - fluid therapy on, 495
- extracorporeal carbon dioxide removal, 222
- extracorporeal circulatory support, 93
- extracorporeal membrane oxygenation, 91–92, 99, 209
 - antibiotics, 442
 - bronchospastic conditions, 222
- extubation, 8
 - assessment for readiness, 234–235
 - criteria for, 235–237
 - failure, 237–238
 - high flow oxygen therapy after, 239
 - NPPV before, 198
 - premature, mortality, 231
 - procedure, 237
 - respiratory failure after, NPPV, 197–198
- exudative phase, ARDS, 204–205
- eye response, Glasgow Coma Scale, 294
- eyes, brain death diagnosis, 295

- faces pain scale, 12
- facial sensory and motor response, brain death
 - diagnosis, 295
- factor Xa, target range, 384
- factor Xa inhibitors, 553
 - coagulation parameters, 567
 - monitoring, 554
 - reversal, 557–558
 - time to hemostasis after stopping, 559
- falls, 338, 345
 - transfer to level 1 trauma facilities, 342
- fasciitis, necrotizing, 394
 - antibiotics, **432**
- fasciotomy, **353**
- fatty liver, total parenteral nutrition, 54
- fecal microbiota therapy, 423–424
- femoral artery, 21, 75
- femoral vein, 18
 - ultrasonography, 32
- fentanyl, 12–13
 - anti-shivering protocol, 164
 - bronchoscopy, 38
 - bronchospasm, **225**
- fetal–maternal hemorrhage, 381
- fever, 389–401
 - from blood product transfusion, 550–551
 - bronchoscopy, 40
 - dialysis, **511**
 - differential diagnosis, 391–392
 - examination, 392
 - imaging tests, 398
 - laboratory diagnosis, 396–398
 - measurement, 391
 - mechanism, 390
 - treatment, 399–400
- fever equivalents, 389
- fiberoptic transducers, intracranial pressure, 286
- fibrin degradation products, DIC, 562
- fibrinogen, 566–567
 - blood transfusion on levels, 547
 - DIC, 562
 - transfusion, 546
- fibroproliferative phase, ARDS, 205
- fidaxomicin, 423, 425
- fingerstick glucose, 256, 279
 - organ donors, 297
- fire safety, 349
- firearms, trauma, 338, 339, 342
- fish mouth sign, 39
- fistula
 - bedside percutaneous dilational tracheostomy, 44
 - enteroatmospheric, 329
- ‘5T’ and ‘5H’ rule, 108
- fixed dilated pupil, 270
- FloTrac™ system, 76
- flow cycled ventilation, 187, 188
- flow (parameter), LV assist devices, 95
- flow rate
 - HFNC, 196
 - inspiratory, 186
 - peak expiratory, serial monitoring, 221
- flow trigger, mechanical ventilation, 187
- flow–volume loop, 178
- fluconazole, **445**
- Fluid and Catheter Treatment Trial, 77
- fluid balance, 494
- fluid distribution, 494–495, 498
- fluid restriction, 233
- fluid status, 495–496, *see also* intravascular volume
- fluid therapy, 493–499
 - acute hepatic failure, 362
 - acute kidney injury, 504
 - ARDS, 209
 - burns, 348, 350, 353, 354
 - complications, 497–498
 - on fluid compartments, 495
 - hypercalcemia, 531
 - hypernatremia, 528
 - hyponatremia, 525, 526
 - sepsis, 409, 410

- fluid therapy (*cont'd*)
 - shock, 84, **155**
 - trauma, 342, 346
 - venous thromboembolism, 581
- flumazenil, **304**
- fluoroquinolones, **437–438**
 - pneumonia, **456, 457**
 - Legionella*, 456
- focused assessment using sonography for trauma (FAST)
 - exam, 26, 341
- folate, methanol poisoning, **305**
- fomepizole
 - ethylene glycol poisoning, **305**
 - methanol poisoning, **305**
- fondaparinux, **132**, 136, 553, **582**
- foreign body, upper airway obstruction, 219
- fosphenytoin, 279, **280**, 281
 - children, 282
- 4T scoring system, heparin-induced thrombocytopenia (HIT), 566
- fraction of inspired oxygen (FiO_2), 185, 186, 196
 - ARDS, 210
 - bronchospastic conditions, 224
 - pregnancy, 378
 - ventilator-associated complications, 393
- fractional excretion of phosphate (FePO_4), 522
- fractional excretion of sodium (FeNa), 503
- fractional excretion of urea (FeUrea), 503
- fractures, skull, 268, 271
- free water deficit, 528
- fresh frozen plasma, 496, 546, 564, 571
 - complications, 558
 - vitamin K antagonist reversal, 556, **558**
- frostbite, 349
- frozen plasma (FP24), 546
- fulminant hepatic failure, 359
- functional residual capacity
 - apnea time, 6
 - pregnancy, **378**
- fungal infections, CNS infections, 464, 475
- furosemide, 383
 - albumin–furosemide chaser, 497
 - potassium removal, 515
- GABA-A receptors, status epilepticus, 276
- gain, ultrasonography, 27
- galactomannan, 398
- gallbladder, imaging, 398
- ganciclovir, **447**
- gangrene, 394
- gastric ulcers, 312
- gastroenteritis, 405
- gastroesophageal varices, 312, 313, 316
- gastrointestinal bleeding, 64, 311–320, 405
 - coagulopathy, 312, 566
 - endotracheal intubation, 66
 - left ventricular assist devices, 96–97
 - stress ulcers, 193
- gastrointestinal tract
 - NPPV for procedures, 198
 - shock, 156
- gentamicin, **440, 457**
- gestational thrombocytopenia, **572**
- Glasgow Blatchford score, 314
- Glasgow Coma Scale, 293–294
 - SOFA score, **407**
- Glidescope® laryngoscope, 5
- glomerular filtration rate
 - acute kidney injury, **501**
 - pregnancy, **378**, 505
- glossopharyngeal nerve, 3
- glucagon
 - beta-blocker overdose, **304**
 - calcium channel blocker overdose, **304**
- glucose
 - brain, 289
 - fingerstick, 256, 279
 - organ donors, 297
 - monitoring, 57, 58, 60, 61
- glutamate, brain, 288
- glycemic control, 55–61
- glycerol, brain, 288
- glycolysis, cerebral, 288
- glycopeptides, **440**
- glycopyrrolate, **306**
- gout, 396
- GP IIb/IIIa inhibitors, **131**
- GRACE risk model, 128
- grafting
 - burns, **353**
 - coronary artery bypass, **132**, 137
- Gram-negative bacilli, antibiotics, **422**
- guidewires
 - arterial access, 21
 - central venous access, 18–19, 24
 - tracheostomy, 46
- Guillain–Barré syndrome, 295
 - hyperkalemia vs, 514
- Haemophilus influenzae*, antibiotics, **422**
- half-time, context-sensitive, 10
- haloperidol, 247
- Hamman–Rich syndrome, 208
- hand hygiene, 414, 415
- Hartmann's solution, 497
- HASHTI mnemonic, 558
- head trauma, 342, *see also* traumatic brain injury
- health care-associated meningitis, 463
- health care-associated pneumonia, 449
 - antibiotics, 455, **457**
- heart
 - artificial, 89, 97
 - valvular disease, 207
- heart failure, 139–147, 219, 578

- acute coronary syndromes, 127, 134
- dobutamine, **87**, 144
- management, 139–147
- mechanical ventilation, 192
- NPPV, 143, 199
- peri-partum cardiomyopathy, 381–383
- ventilator-associated events vs, 420
- Heart Failure Survival Scores, 142
- heart rate, 82
- HeartMate™ devices (Thoratec), 94, 101
- HeartWare™ HVAD, 94, 101
- HEAT-PPCI trial, 137
- Helicobacter pylori*, 312
- heliox, 227
- HELLP syndrome, 563, 566, 572
 - acute hepatic failure, 360
 - treatment, **572**
- hematemesis, 313
- hematochezia, 313
- hematocrit, red blood cell preparations, 545
- hematoma
 - subdural, 271, 467
 - traumatic brain injury, 269, 271
- hematopoietic stem cell transplantation, ARDS, 206
- hemicraniectomy, 258
- hemoclips, 316
- hemodialysis, 508, 509
 - antibiotics, 434
 - clotting, **511**
 - continuous veno-venous, 509, 512
 - energy requirements, **51**
 - hypercalcemia, **531**
 - intermittent, 509–510
 - antibiotics, 434
 - lithium poisoning, **305**
- hemodynamic monitoring, 63, 75–81, 85
- hemofiltration, 508, *see also* continuous veno-venous
 - hemofiltration
- hemoglobin levels
 - blood loss, 342
 - for blood transfusion, 63–64
 - organ donors, 297
 - shock, 152
- hemoglobin A1c, 57
- hemolytic transfusion reaction, 395, 548–549
- hemolytic-uremic syndrome, 563, 566, **572**
- hemoperfusion, 509
- hemoptysis, bronchoscopy, 39, 40
- hemorrhage, *see also* blood loss
 - anticoagulants, 132–133, 552–560
 - antiplatelet therapy, 132–133, 312
 - children, 344
 - coagulopathies, 571
 - criteria for major, 552
 - diffuse alveolar, 208
 - gastrointestinal tract *see* gastrointestinal bleeding
 - intraabdominal, 39, 40
 - intracerebral *see* intracerebral hemorrhage
 - parenchymal, strokes, **251**
 - post-partum, 380–381
 - pulmonary, ventilator-associated events vs, 421
 - renal replacement therapy, **511**
 - stopping, 19
 - subarachnoid *see* subarachnoid hemorrhage
 - trauma, 342
- hemorrhagic transformation, strokes, **251**
- Henderson–Hasselbalch equation, 537, 538
- heparin, *see also* low molecular weight heparin; unfractionated heparin
 - acute coronary syndromes, **131**, 137
 - acute mesenteric ischemia, 334
 - arterial blood gases, 174, 176
 - bleeding from, 553
 - monitoring, 554, 564
 - pregnancy, 384
 - reversal, 557, **558**
 - time to hemostasis after stopping, 559
 - venous thromboembolism, 64
- heparin–factor 4 complexes, 562–563
- heparin-induced thrombocytopenia (HIT), 562–563, 566, **572**, 582
 - antibody, 570
- hepatic artery dissection, liver transplantation, 367
- hepatic failure, **572**
 - acute, 359–364
 - diagnosis, 360–361
 - prognosis, 363
 - treatment, 362–363
 - coagulation parameters, 567
 - hemorrhage, 555
 - HIV infection, 482
 - indications for transplantation, 366
 - neutropenic fever, 589
 - poisonings, 308
 - total parenteral nutrition, 53
- hepatic steatosis, total parenteral nutrition, 54
- hepatitis, acute severe, 360
- hepatitis C, HIV infection with, 481
- hepatization *see* consolidation
- hepatobiliary iminodiacetic acid (HIDA) scan, 398
- herniation syndromes, cerebral, 285–286, 294
- herpes simplex encephalitis, 464, 472, 474, 476, 478
- high flow oxygen therapy via nasal cannula, 194, 195, 196
 - after extubation, 239
 - ARDS, 210
 - initiation, 199
 - respiratory failure, 197
- high pressure alarm, ventilators, 190
- histamine 2 receptor antagonists, 64, 313
- HIT *see* heparin-induced thrombocytopenia
- HIV infection, 480–490
 - brain abscess, 473
 - classification, **481**

- HIV infection (*cont'd*)
 - CNS infections, 475
 - diagnosis, 482–484
 - encephalitis, 474
 - pancreas transplantation criteria, 371
 - prognosis in ICU, 488
 - treatment, 484–488
- hospital-acquired pneumonia, 449
 - antibiotics, 455
- human chorionic gonadotropin, 341
- humidification, NPPV, 200
- Hunt and Hess grade, subarachnoid hemorrhage, 256
- hydralazine, 105, 382, **385**
- hydrocephalus
 - subarachnoid hemorrhage, 252, 263
 - treatment, 258
- hydrochlorothiazide, 383
- hydrocortisone, septic shock, 411
- hydrogen ions, 537, 538, 539
- hydromorphone, 13
- hydronephrosis, ultrasonography, 32
- hydrophilic antibiotics, 434
- hydroxycobalamin, **304**
- hydroxyethyl starch, 497, 498
- hydroxyurea, 592
- hyperactive delirium, 243
- hyperbaric oxygen, **304**
- hypercalcemia, 529–532
 - phosphate binder therapy, 520
- hypercapnia, 170, **185**
 - ARDS, 210
 - permissive, 224, 233–234
 - pregnancy, 228
 - traumatic brain injury, 270
- hyperchloremic metabolic acidosis, 498
- hypercoagulable state, 576
- hyperextension, bedside tracheostomy, 44, 45, 47
- hyperfibrinolysis, coagulation parameters, 567
- hyperglycemia, 55–57
 - sepsis, 409
 - total parenteral nutrition, 53
- hyperinflation, dynamic, 217, 227
- hyperinsulinemia–euglycemia therapy, **304**
- hyperkalemia, 65, 513–516, 535, 536
 - arrhythmias, 111
 - from blood transfusion, 550
- hypermagnesemia, 379
- hyponatremia, 527–529
 - arrhythmias, 111
- hyperphosphatemia, 518–520, 535
- hyperpyrexia, 389
 - on oxyhemoglobin curve, 174, 176
- hypersensitivity, drugs causing, 394
- hypersplenism, 564
- hypertension
 - hypokalemia, 518
 - malignant, 563
 - pre-eclampsia, 385
 - risk of acute coronary syndromes, 124, 125
 - shock, 154
 - strokes, 251
- hypertensive emergency, 102–107
- hypertensive urgency, 102–107
- hyperthermia, 389, 390
 - malignant, 395
- hypertonic saline
 - acute hepatic failure, 362
 - cerebral edema, 258
 - hyponatremia, 526
 - intracranial pressure, 271
 - tricyclic antidepressant poisoning, **306**
- hypertriglyceridemia, TPN, 54
- hyperventilation
 - acute hepatic failure, 362
 - intracranial pressure, 271, 286, 287
- hyperventilation syndrome, 219
- hypnotics, toxic syndromes, **301**
- hypoactive delirium, 243
- hypoalbuminemia, anion gap, 540
- hypocalcemia, 532–534
 - from blood transfusion, 550
 - hyperphosphatemia, 519
 - phosphate binder therapy, 520
- hypocapnia, traumatic brain injury, 270
- hypofibrinogenemia, 546
- hypoglycemia, 59, 60
- hypokalemia, 516–518, 535, 536
 - arrhythmias, 111
- hypomagnesemia, 517
- hyponatremia, 64–65, 473, 523–527, 535
 - arrhythmias, 111
- hypoperfusion, 142, 150, 151, 152
- hypophosphatemia, 517, 521–523, 536
- hypotension, 150, 151, 152, 154
 - blood loss, 314
 - blood transfusion reactions, 550
 - brain death vs, 295
 - dynamic hyperinflation, 227
 - for intracranial pressure control, 287
 - organ donors, 297
 - rebound, **88**
 - renal replacement therapy, 510, **511**
 - sepsis, **403**, 406, 409, 410
 - SOFA score, **407**
 - trauma, 342
- hypothalamus, 390
- hypothermia
 - from blood transfusion, 550
 - brain death vs, 295
 - on oxyhemoglobin curve, 174, 176
 - pulse oximetry, 171, 172
 - sepsis, 409
- hypothermia (therapeutic) see therapeutic temperature modulation

- hypovolemia, 495–496
- hypovolemic shock, 83, 149, 150, 158, 405
 - with left ventricular assist devices, 96
 - prognosis, 157
 - ultrasonography, 33
- hypoxemia, 169, **185**
 - ARDS, 169, 205, 207
 - bronchoscopy, 40
 - NPPV for, 197
 - pregnancy, 227
 - pulmonary vascular resistance, 192
 - weaning from ventilation, 239
- hypoxia
 - brain, 287
 - tissues, 149
- ibutilide, 114, 115
- ICU-VTE score, 576, **577**
- idarucizumab, 557, **558**
- ideal body weight, 51
- idiopathic interstitial pneumonia, 453
- idiopathic pulmonary fibrosis, 207
- idiopathic thrombocytopenic purpura (ITP),
 - treatment, **572**
- ileus, *Clostridium difficile*, 393, 424
- imipenem, **457**
- imipenem/cilistatin, **437**
- immune-mediated thrombocytopenia, 563, **564**
- immune reconstitution inflammatory syndrome, 488
- immunocompromised patients
 - CNS infections, 466
 - fever, 400
 - sepsis, 411
- immunosuppression
 - abdominal sepsis, 325
 - fever, 400
 - history, 368
 - pneumonia, 458
 - solid organ transplantation
 - kidney, 370
 - liver, 368
 - pancreas, 372
 - small bowel, 375
- IMPACT (calculator), 273
- impedance monitors, respiratory, 171
- Impella intraluminal catheter-based axial flow pump,
 - 91, 92, 99
- Improving Palliative Care in the ICU (IPAL-ICU) model, 69
- indirect calorimetry, 53
- indirect factor Xa inhibitors, 553
 - time to hemostasis after stopping, 559
- induction agents, laryngoscopy, 6, 7
- infarction, *see also* acute coronary syndromes; myocardial infarction
 - brain, 252
- infection-related ventilator-associated complication (IVAC), 391, 419, 450
- infections, 66–67, **391**, 414–428, *see also* sepsis
 - acute hepatic failure, 360, 362
 - avoidance, 390
 - from blood transfusions, 395
 - bronchospastic conditions, 217
 - central nervous system, 394, 463–479, *see also* encephalitis; meningitis
 - intravascular devices, 392–393
 - bloodstream infections (CLABSIs), 66–67, 415–417, 419, **430**
 - left ventricular assist devices, 97
 - lumbar puncture, 398
 - total parenteral nutrition, 53
 - urinary tract, 67, **405**
 - bacteriology, 397
 - catheter-associated (CAUTIs), 67, 393, 397, 417–419
 - ventilator-associated, 393
- infectious ventriculitis, 286
- inferior vena cava
 - filters, 577, 581
 - ultrasonography, 29–30, 35, 79, 496
- inflammation
 - ARDS, 204
 - bronchospastic conditions, 217
 - meningitis, 465
- influenza, 405
- inhaled antibiotic therapy, 442
- Injury Severity Score, 340
- innervation, airway, 3–4
- innominate artery fistula, percutaneous tracheostomy, 44
- inodilators, 83
- inotropes, 83
 - pulmonary embolism, 581
- inspiratory flow rate, 186
- inspiratory positive airway pressure, 196, 199, 223
- inspiratory rise time, 186
- inspiratory time, 186
- inspiratory to expiratory time ratio, 186
- insulin
 - for beta-blocker overdose, **304**
 - glycemic control, 58, 60, 61
 - for hyperkalemia, 515
 - for organ donors, 297
 - pancreas transplantation criteria, 371
 - for sepsis, 409
 - total parenteral nutrition, 52
- Intensive Care Delirium Screening Checklist, **245**
- interdisciplinary meetings, palliative care, 69
- intermittent hemodialysis, 509–510
 - antibiotics, 434
- internal jugular vein (IJ), 18
 - thrombosis, 22
- international normalized ratio (INR), 561, 566–567
 - anticoagulant reversal, **557**
- interstitial fluid, fluid therapy on, 495
- interstitial lung disease, 207
 - idiopathic interstitial pneumonia, 453

- intestinal failure, 374
- intestinal mucosal atrophy, 54
- intestinal transplantation, 374–375, 376
- intoxication, 299–308
 - brain death vs, 295, 296
- intra-abdominal hypertension, 326–327, 328
- intra-abdominal pressure, 326, 327, 328
- intra-aortic balloon pumps, 91, 98
- intracellular fluid, fluid therapy on, 495
- intracellular sodium, 495
- intracerebral hemorrhage, 250, 252, 253, 263, 267
 - presentation, 254
 - score, 256
 - treatment, 258, 259
- intracortical electroencephalography, 289
- intracranial compliance, 284, 291
- intracranial pressure, 266, 291
 - acute hepatic failure, 362
 - children, 273
 - CNS infections, 473
 - management, 271, 272
 - mannitol, 258, 271, 286
 - monitoring, 271, 284–287
 - normal range, 284
- intracranial thrombophlebitis, septic, 473
- intraperitoneal free fluid, ultrasonography, 32
- intravascular devices, *see also* central venous access
 - blood stream infection, **430**
 - central line-associated (CLABSIs), 66–67, 415–417, 419, **430**
 - fever, 392–393
- intravascular volume
 - fluid therapy on, 495
 - IVC ultrasonography, 29–30
- intravenous immunoglobulin, pancreas transplantation, 372
- intravenous insulin, 58, 60, 61
- intrinsic PEEP *see* auto-PEEP
- intrinsic renal injury, 501, 502
- involuntary movements, 276
- ipratropium bromide, 225
- iron poisoning, **305**
- irradiated packed red blood cells, 545
- irradiated platelet products, 547
- irrigation, whole bowel, 303
- isavuconazole, **445**
- ischemia
 - delayed cerebral (DCI), monitoring, 289
 - mesenteric, 330–336, 394
 - traumatic brain injury, 266
- ischemic colitis, 312, 332, 424
- ischemic heart failure, 140
- ischemic stroke *see* acute ischemic stroke
- islet cell transplant, **372**
- isoflurane, 226
- isolation, CNS infections, 466
- isoniazid poisoning, **305**
- isoprenaline, 113
- itraconazole, **445**
- jellyfish sign, 31
- Josephson's sign, 111
- jugular venous oxygen saturation (SjvO₂), 287
- jugular venous pressure, 141, 142
- Kernig's sign, 468
- Kernohan's notch phenomenon, 270
- ketamine, 16, 226, **280**, 281
- kidneys, *see also* acute kidney injury; *entries beginning* renal. . .
 - transplantation, 368–370, 371, **372**, 376
- King's College criteria, acetaminophen, 361
- Klebsiella pneumoniae* carbapenemase, 443
- Kleihauer–Betke test, 381
- Kussmaul respiration, 170
- L-carnitine, **306**
- labetalol
 - aortic dissection, **105**
 - pre-eclampsia, **385**
- laboratory tumor lysis syndrome, 589
- lacosamide, **280**
- lactate, 150, 151, 157
 - acute mesenteric ischemia, 331
 - antiretroviral therapy, 486
 - arterial blood gases and, 176
 - sepsis, 397, 409
- lactate : pyruvate ratio (LPR), brain, 288, 289
- lactated Ringer's solution, 497
- lactic acidosis, HIV infection, 486
- lactulose, 362
- lanthanum, **520**
- laparotomies
 - decompressive, 328, 329
 - repeat, 323
- large curvilinear probes, ultrasonography, **27**, **34**
- laryngeal mask airways, 5
- laryngoscopes, 5
- laryngoscopy, 6–7
 - anesthesia, 6, 8
 - predictors of difficulty, 4
- laryngospasm, 8
- larynx, 3, 4
 - bronchoscopy, 39
- laser Doppler flowmetry, cerebral blood flow, 288
- lateral transtentorial herniation, 286
- lead poisoning, **305**
- leak test, endotracheal tubes, 8, 236
- leaks
 - cerebrospinal fluid, 466
 - NPPV, 200
- left ventricle
 - ejection fraction, acute coronary syndromes, 135
 - mechanical ventilation on function, 192
 - ultrasonography, 29
- left ventricular assist devices, 89, 93–97, **100**
 - complications, 96–97
 - Impella intraluminal catheter-based axial flow pump, 91

- monitoring parameters, 95
- troubleshooting, 95–96
- leg veins, ultrasonography, 32, 36
- Legionella pneumonia*, 453
 - antibiotics, 456
- leptospirosis, 474
- leukemia
 - arterial blood gases, 174, 176
 - tumor lysis syndrome, 586
- leukocyte count
 - Clostridium difficile*, 425, **431**
 - HIV infection, 483
 - shock, 152
- leukocyte esterase, 397
- leukocyte-reduced concentrates, 545
- leukocytes, thrombocytopenia, 568
- leukocytosis
 - acute mesenteric ischemia, 333
 - causes, **391–392**
 - differential diagnosis, 392
- leukoreduced platelet products, 547
- leukostasis *see* symptomatic leukostasis
- leukotriene-receptor antagonists, 226
- Leuven trial, glycemic control, 57–58, 59
- level 1 trauma facilities, transfer to, 342–343
- levetiracetam, **280**
- levofloxacin, **437, 456, 457**
- levothyroxine, thyroxine replacement protocol, 297
- LIDCO system, 76
- lidocaine
 - bedside tracheostomy, 45
 - bronchoscopy, 38
 - cardiac arrest, 161
 - monomorphic ventricular tachycardia, 115
- limit phase, mechanical ventilation, 187
- lincosamides, **438**
- linear array probes, ultrasonography, **27, 34**
- linezolid, **441**
 - methicillin-resistant *S. aureus*, **457**
 - stopping, 444
- lipids, TPN, **52, 53**
 - hypertriglyceridemia, 54
- lipophilic antibiotics, 434
- liposomal amphotericin B, **446, 520**
- Listeria meningitis, **430, 474**
 - pregnancy, 475
- lithium dilution cardiac output (LIDCO)
 - system, 76
- lithium poisoning, **305**
- liver, *see also* hepatic failure
 - steatosis, TPN, 54
 - transplantation, 362, 363, 365–368
 - with intestine, **374**
 - techniques, 367–368, 375
- liver function tests, 362
 - neutropenic fever, 589
 - thrombocytopenia, 570
- living donors
 - kidney, 368
 - pancreas, 371
- loading doses, antibiotics, 433
- local tissue transfer, burns, **353**
- locked-in syndrome, 293, 295
- log rolling, 340
- lorazepam, 14
 - bronchospasm, **225**
 - NPPV, 200
 - status epilepticus, **279, 281**
- low dose aspirin, 125, 254
- low dose catheter-directed thrombolysis, **582**
- low dose unfractionated heparin, 577
- low molecular weight heparin, 64, 553, 565
 - coagulation parameters, 567
 - pulmonary embolism, **582**
 - reversal, 557, **558**
 - thromboprophylaxis, 577
 - time to hemostasis after stopping, 559
- low pressure alarm, ventilators, 190
- lumbar puncture
 - CNS infections, 467, 469
 - cryptococcal meningitis, 487–488
 - fever, 398
 - HIV infection, 484
 - minimum platelet count, **571**
 - status epilepticus, 278
 - strokes, 256
- Lundberg A waves, 285
- lung flapping sign, 31
- lung point sign, 30
- lung sliding, 30, 31
- lungs, *see also* interstitial lung disease; pneumonia;
 - entries beginning pulmonary. . .*
 - cancer, SVC syndrome, 587
 - compliance, 179, 186, 209
 - examination, 170
 - injury, 206, 207, 395, 549–550
 - protective ventilation
 - HIV infection, 484
 - tidal volume setting, 185
 - transplantation, ARDS, 206
 - ultrasonography, 30–31, 181, 461, 462
- lupus anticoagulant (disorder), coagulation
 - parameters, 567
- lymphedema, 578
- lymphoproliferative disorder, post-transplant, 375
- MAC blade, laryngoscopy, 6
- MACOCHA score, **66**
- macrocytosis, 570
- macrolides, **438**
- macronutrients, 50
- mafenide acetate, **353, 354**
- magnesium
 - arrhythmias, 111
 - hypermagnesemia, 379

- magnesium therapy
 - anti-shivering protocol, 164
 - bronchospasm, **225**, 226
 - polymorphic ventricular tachycardia, 115
 - pre-eclampsia, **385**
- magnetic resonance imaging
 - acute mesenteric ischemia, 333
 - angiography
 - brain death, 296
 - strokes, 257
 - CNS infections, 469
 - heart failure, 143
 - pulse oximetry, 171
 - shock, 153
 - status epilepticus, 278
 - strokes, 257, 263
 - traumatic brain injury, 269
- mainstream technique, end-tidal CO₂, 172
- maintenance doses, antibiotics, 433–434
- malignant hypertension, 563
- malignant hyperthermia, 395
- Mallampati classification, 4, 9
- malnutrition, hypophosphatemia, 521
- mandatory ventilation, synchronized intermittent, 188
- manganese, liver failure, 53
- mannitol
 - acute hepatic failure, 362
 - intracranial pressure, 258, 271, 286, 287
- MASCC score, 589
- mask ventilation, 6
 - difficult airway, 7
 - predictors of difficulty, 4
- masks
 - nasal-oral, 196, 202
 - NPPV, 200, 202
- massive pulmonary embolism, 575, 579, 583, 585
- massive transfusion protocol, 547–548
- maximal inspiratory pressure, 180
- McConnell's sign, 29, 181
- mean airway pressure, 186
- mean arterial pressure (MAP), 82
 - acute hepatic failure, 362
 - brain death diagnosis, 295
 - intra-aortic balloon pumps, 91
 - pregnancy, **378**
 - traumatic brain injury, 266
- mechanical circulatory support, 89–101
- Meckel's diverticulum, 318
- MELD score, 361
- melena, 313
- meningeal irritation, 468
- meningitis, **405**, 463–479
 - antibiotics, **430**, 466, 470, 471–472
 - diagnosis, 467
 - HIV infection, 482, 487–488
 - prognosis, 476
- meningoencephalitis, viral, 465
- meningovascular syphilis, 474
- meperidine, anti-shivering protocol, 164
- Mercedes Benz signs
 - bronchoscopy, 39
 - echocardiography, 28
- meropenem, **457**
 - valproate and, **437**
- mesenteric ischemia, 330–336, 394
- mesenteric venous thrombosis (MVT), 331, 332, 333
- metabolic acidosis, 538, 539
 - delta ratio and delta-delta gap, 540
 - hyperchloremic, 498
 - presentation, 541
 - total parenteral nutrition, 53
 - treatment, 517, **542**
 - Winter's formula, 302
- metabolic alkalosis, 538, 539, 541, **542**
- metabolic crisis, brain, 289
- metabolic distress, brain, 289
- metabolic encephalopathy, 467
- methadone, 13
- methanol poisoning, **305**
- methemoglobinemia
 - arterial blood gases, 177
 - treatment, **305**
- methicillin-resistant *Staphylococcus aureus*
 - antibiotics, **422**, **457**
 - de-escalation, 444
- methylcellulose ionic silver dressing, **353**
- methylene blue, **305**
- methylprednisolone
 - acute exacerbation of COPD, 226
 - asthma, 222, **225**
 - immunosuppression, 370, 372, 375
 - pneumonia, **456**
- methylxanthine, 225
- metoprolol, 114, 382
- metronidazole, **441**
 - Clostridium difficile*, 425
- micafungin, **446**
- microangiopathies, thrombotic, 563
- microcirculatory failure, 149, 156
- microdialysis, cerebral, 288–289
- micronutrients, 49, 50
 - total parenteral nutrition, 52
- midazolam, 14
 - bronchoscopy, 38
 - bronchospasm, **225**
 - status epilepticus, **279**, **280**, 281
 - children, 282
- middle cerebral artery embolism, 264
- Miller blade, laryngoscopy, 6
- milrinone, **87**, 144, **155**, 383
- minimally conscious state, 293
- minute ventilation, 185
 - pregnancy, **378**
- mirror artifact, ultrasonography, 28
- mixed delirium, 243

- mixing studies, aPTT-based, 567–568
- mobilization, 233
- Mobitz types, AV block, 109, 112, 117
- modes of mechanical ventilation, 186, 187–190
- MOMENTUM trial, LVAD, 94
- monitoring
 - anticoagulants, 554, 564
 - arterial lines, 85
 - central nervous system, 284–292
 - electroencephalography, 276
 - glucose, 57, 58, 60, 61
 - hemodynamic, 63, 75–81, 85
 - intracranial pressure, 271, 284–287
 - left ventricular assist devices, 95
 - NPPV, 199–201
 - peak expiratory flow rate, 221
 - pregnancy, 380
 - renal replacement therapy, 511
 - respiratory, 169–183
 - respiratory muscle strength, 179–181
 - sedation and analgesia, 11–12
 - sodium, 65
 - total parenteral nutrition, 53
- monobactams, **437**
- monomorphic ventricular tachycardia, 110, 115
- Monro–Kellie doctrine, 284
- Morison's pouch with hemothorax view, 31, 32, 36
- morphine, 13
 - bronchospastic conditions, 224, **225**
- mortality
 - acute hepatic failure, 363
 - acute kidney injury, 506
 - amniotic fluid embolism, 384
 - ARDS, 213
 - bronchospastic conditions, 228
 - risk factors, 218
 - burns, 347, 355
 - CAUTIs, 419
 - CLABSIs, 417
 - Clostridium difficile* infection, 425
 - delirium, 248
 - eclampsia, 385
 - gastrointestinal bleeding, 318
 - HIV infection, 482, 488
 - mechanical ventilation, 185
 - meningitis, 476
 - pneumonia, 450, 459
 - pre-eclampsia, 385
 - premature extubation, 231
 - pulmonary embolism, 584
 - pregnancy, 384
 - renal replacement therapy, 510
 - sepsis, 402
 - shock, 157
 - strokes, 261
 - trauma, 337, 338
 - traumatic brain injury, 273
 - ventilator-associated pneumonia, 421
 - mosquitoes, diseases from, 466
 - motor response
 - Glasgow Coma Scale, 294
 - neurotrauma, 268
 - motor vehicle crashes, transfer to level 1 trauma facilities, 342
 - movements
 - brain death, 296
 - involuntary, 276
 - moxifloxacin, **437, 456**
 - MRC CRASH (calculator), 273
 - mucolytics, 226
 - Mucor* (spp.), 475
 - mucormycosis, brain abscess, 473
 - MUDPILES, anion gap, 540
 - multidrug-resistant organisms, 410, 416, 420
 - antibiotics, **422, 442–443**
 - pneumonia, 451
 - multifocal atrial tachycardia, 115
 - Multinational Association for Supportive Care in Cancer (MASCC) score, 589
 - multiorgan dysfunction syndrome, 344
 - multivisceral transplantation, 374–375, 376
 - mupirocin, **353**
 - murmurs
 - acute coronary syndromes, 126
 - aortic dissection, 104
 - Murphy's eye, 5
 - mycostatin, **353**
 - myocardial infarction, *see also* acute coronary syndromes
 - heart failure, 141
 - intra-aortic balloon pumps, 91
 - myocarditis, **405**
 - myoglobinuria, 354
 - myonecrosis, 394
 - myopathy, 63
 - N*-acetylcysteine
 - acute hepatic failure, 363
 - for contrast-induced nephropathy, 65
 - Naegleria fowleri*, 475
 - naftillin, **435**
 - naloxone
 - cardiac arrest, 161
 - opioid toxicity, **305**
 - overdose patients, 307
 - NAP1 strain, *Clostridium difficile*, 423
 - narrow complex tachyarrhythmias, 108, 110, 121
 - diagnostic algorithm, 112
 - treatment algorithm, 116
 - nasal cannula, 202, *see also* high flow oxygen therapy via nasal cannula
 - nasal-oral masks, 196, 202
 - nasal prongs, 196
 - nasogastric tubes, GI bleeding, 314
 - nasotracheal intubation
 - contraindications, 343
 - sinusitis, 393

- natriuretic factors, 494
- nebulized treatment, 225
 - epinephrine, 238
- neck
 - airway difficulty, 4
 - angiography, 341
 - bedside tracheostomy, 44, 47
 - CT angiography, 269
 - pounding sensation, 111
- necrotic bowel, 336
- necrotizing infections, 394
 - antibiotics, **432**
- Neelon and Champagne Confusion Scale, **245**
- Neisseria meningitidis*, 474
- nesiritide, **144**, 146
- neuroleptic malignant syndrome, **301**, **305**, 395
- neuromuscular blocking agents
 - ARDS, 210
 - brain death vs effect, 295
 - bronchospastic conditions, 222, 224
 - capnography waveform, **175**
 - cisatracurium, bronchospasm, **225**
 - pregnancy, 228
- neuromuscular disease
 - NPPV, 197
 - respiratory examination, 170
- neuron-specific enolase, 165
- neuropathies, metronidazole, **441**
- neurotrauma, 265–274
 - tracheostomy, 43, 271
- neutropenia
 - causes, **391–392**
 - fever, 389, 400
 - antibiotics, **431**
- neutropenic fever, 586, 587, 588, 589, 591, 592, 594
- neutrophils
 - absolute count (ANC), low, 400
 - ARDS, 204
- New Delhi metallo-beta-lactamase, 443
- nicardipine, aortic dissection, **105**
- NICE-SUGAR trial, 58, 59
- nifekalant, 161
- NIH Stroke Scale, 255–256
- nitrites, urine, 397
- nitrogen balance, total parenteral nutrition, 53
- nitroglycerin, **144**, 382
- nitroimidazoles, **441**
- nitroprusside, **144**
 - aortic dissection, **105**, 106
- NMDA receptors, status epilepticus, 276
- non-convulsive status epilepticus, 275, 276, 277, 289
- non-Hodgkin's lymphoma
 - superior vena cava syndrome, 587
 - tumor lysis syndrome, 586
- non-invasive positive pressure ventilation, 184, 194–202,
 - see also bilevel positive pressure ventilation
- ARDS, 210
- bronchospastic conditions, 223
 - for extubation failure, 238
- heart failure, 143, 199
- initiation, 199
- management, 199–201
- pneumonia, 455
- pregnancy, 228
- non-occlusive mesenteric ischemia (NOMI), 331, 332
 - papaverine, 331, **334**
- non-small cell lung cancer, SVC syndrome, 587
- non-ST elevation myocardial infarction, 123, 124, 134, 136
- nonsteroidal anti-inflammatory drugs
 - fever and, 400
 - gastrointestinal bleeding, 312, 313
- norepinephrine, **86**, **144**, **155**
 - acute hepatic failure, 362
 - angiotensin II on, 494
 - organ donors, 297
 - peri-partum cardiomyopathy, 383
- normal saline, 497
 - hyperchloremic metabolic acidosis, 498
 - for hyponatremia, 529
 - intracranial pressure management, 286
- nosocomial pneumonia, 449, 450
- nuclear medicine
 - CBF test, brain death, 296
 - fever, 398
 - gastrointestinal bleeding, 315, 316
 - hepatobiliary iminodiacetic acid (HIDA) scan, 398
 - V/Q scan, 580
- nucleoside reverse transcriptase inhibitors, 486
- numeric ranking scales, pain, 12
- Nursing Delirium Screening Scale, **245**
- nutritional support, 49–54, 64
 - acute hepatic failure, 363
 - ARDS, 211
 - burns, 350, 352
 - insulin, 58
 - mechanical ventilation, 233
 - neurotrauma, 271
- NYHA classification, heart failure, 142
- OASIS-5 trial, 136
- obesity
 - bedside tracheostomy, 44
 - endotracheal intubation, 7
 - risk of acute coronary syndromes, 124
 - total parenteral nutrition, 51
 - weaning from ventilation, 239
- objective pain scales, 12
- obstetric emergencies, 377–386
- obstructive shock, 83, 150–151, 158
 - ultrasonography, 33
- octreotide, **306**, **317**
- oculocephalic reflex, 295
- oculovestibular reflex, 295
- olanzapine, 247
- oliguria
 - abdominal compartment syndrome, 327

- sepsis, 409
- omeprazole, **317**
- oncologic emergencies, 586–595
- one hour bundle, Surviving Sepsis Campaign guidelines (2018), 410
- open cranial fractures, 271
- opioids, 12–13
 - addiction, cardiac arrest, 161
 - bronchospastic conditions, 224
 - pregnancy, 227
 - toxic syndromes, **301, 305**
- opportunistic infections, 398, 481, 484
- optimization (phase), treatment for shock, 154
- oral intake, aspiration risk, 4
- organ donors, 297–298
- organophosphates, cholinergic toxidrome, **301, 306**
- orthostatic hypotension, 314
- oseltamivir, **456**
- osmolal gap, 302, 540
- osmolality, 495
 - urine, 525, 528
- osmolarity, 495
- osmosis, renal replacement therapy, 509
- osteomyelitis, vertebral, 479
- outpatient care, trauma, 345
- overfeeding, 49, 52–53, 53
- OXA carbapenemases, 443
- oxacillin, **435**
- oxazolidinones, **441**
- oxygen, *see also* extracorporeal membrane oxygenation; fraction of inspired oxygen (FiO₂); peripheral capillary oxygen saturation (SpO₂); preoxygenation
 - brain, monitoring, 287
 - consumption
 - delivery vs, 149
 - pregnancy, **378**
 - hyperbaric, **304**
 - saturation gap, 302
 - pulse oximetry, 172
- oxygen saturation (SaO₂)
 - arterial blood gases vs pulse oximetry, 174
 - organ donors, 297
 - pulse oximetry, 171, 172
- oxygen tension (pO₂), 176, *see also* arterial oxygen pressure (PaO₂)
 - pulse oximetry, arterial blood gases vs, 174
- oxygen therapy, *see also* high flow oxygen therapy via nasal cannula
 - bronchospastic conditions, 222
 - heart failure, 143
- oxyhemoglobin curve, 174, 176
- P2Y12 inhibitors, **131**
- pacemakers, withdrawal, 71
- packed red blood cells, 545–546
- pain, *see also* chest pain
 - acute abdomen, 323
 - acute coronary syndromes, 126–127
 - acute mesenteric ischemia, 332
 - aortic dissection, 103
 - incidence, 10
 - pancreatitis, 394
 - scales, 12
 - palliative care, 68–71
 - NPPV, 198
 - shock, 154
 - pamidronate, **531**
 - pancreas, transplantation, 370–374
 - pancreas after kidney (PAK) transplantation, 371, **372**
 - pancreatitis, 394, 405
 - HIV infection, 486
 - hypocalcemia, 532
 - papaverine, non-occlusive mesenteric ischemia, 331, **334**
 - paracetamol *see* acetaminophen
 - paradoxical septal wall motion, ultrasonography, 29
 - paralytics *see* neuromuscular blocking agents
 - paraneoplastic encephalitis, 467
 - parapneumonic effusions *see* pleural effusions
 - parasternal positions, cardiac ultrasonography, 28, 34, 79
 - parathyroid hormone (PTH), 532
 - hypercalcemia, 530
 - hypocalcemia, 533
 - parenchymal hemorrhage, strokes, **251**
 - Parkland formula, 350
 - partial liver transplantation, 368
 - pathogens
 - CNS infections, 464, 466, 471, 474
 - colonization, 450
 - pneumonia, 458–459
 - patient positioning
 - airway management, 5, 7
 - bedside tracheostomy, 44, 45, 47
 - peak end-tidal CO₂, 173
 - peak expiratory flow rate, serial monitoring, 221
 - peak inspiratory pressure, 179, 183, 185
 - pentobarbital, **280**, 281, 287
 - penumbra, ischemic, 252
 - peptic ulcers, 312, 314, 316
 - percutaneous coronary intervention (PCI), 130, **132**
 - CABG vs, 137
 - cardiac arrest patients, 163
 - percutaneous dilational tracheostomy, bedside, 42–48
 - peri-partum cardiomyopathy, 156, 379, 381–383
 - pericardial effusion, 28–29, *see also* tamponade
 - pericarditis, 396
 - acute coronary syndromes vs, 126
 - dialysis, **511**
 - perimesencephalic non-aneurysmal SAH, 251
 - peripheral blood smear, thrombocytopenia, 568
 - peripheral capillary oxygen saturation (SpO₂), 172
 - cyanosis, 170, 174, 177
 - FiO₂ vs, 185
 - high flow nasal cannulae, 199
 - NPPV, 199
 - pregnancy, 378
 - pulmonary embolism, 583
 - spontaneous breathing trials, 236
 - peritonism, 333

- peritonitis, 321–325
 - antibiotics, **431**
- permissive hypercapnia, 224, 233–234
- pregnancy, 228
- persistent vegetative state, 293
- pH
 - blood, 176
 - plasma, 537, 538, 539
- pharmacodynamics, antibiotics, 433
- pharmacokinetics, 10–11
 - antibiotics, 433–434
- pharyngeal reflex, 295
- phased array probes, ultrasonography, **27, 34**
- phases, mechanical ventilation, 187
- phenobarbital, **280**
 - children, 282
- phenylephrine, 85, **87–88**
- phenytoin, status epilepticus, **280**
- phosphate
 - false high levels, 520
 - hyperphosphatemia, 518–520, 535
 - hypophosphatemia, 517, 521–523, 536
 - of potassium, **518, 522**
 - replacement, 522
 - renal failure, 523
 - restriction, 520
 - tumor lysis syndrome, 589
- phosphate binder therapy, 520
- phosphate clearance, 520
- phosphodiesterase 3 receptors, **84**
- physical therapy, 226, 227, 233
- physostigmine, **303**
- PiCCO system, 76
- piggyback technique, liver transplantation, 367, 375
- piperacillin/tazobactam, **435, 457**
- pitressin, organ donors, 297
- plaques, atheromatous, 123–124, 138
- plasma exchange, **572**
- plasma products, 546
 - fever, 550
 - sepsis from, 549
- plasma volume, 495
- plateau pressure, 179, 183, 186
 - ARDS, 210
 - bronchospastic conditions, 222, 224
 - high pressure alarm, 190
- plateau waves, 285
- platelet counts, 564, 566–567, *see also* thrombocytopenia
 - disseminated intravascular coagulation (DIC), 562
 - heparin-induced thrombocytopenia (HIT), 566
 - minimum for procedures, **571**
 - shock, 152
 - SOFA score, **407**
- platelet products, 547
 - fever, 550
 - sepsis from, 549
- platelet transfusion, 547, 564, 571
- platelets
 - blood smear, thrombocytopenia, 570
 - blood transfusion on levels, 547
 - lifespan, 554
- PLATO trial, 136
- pleural effusions
 - bacteriology, 397
 - clinical signs, 453
 - computed tomography, 462
 - pneumonia, 455, 457
 - ultrasonography, 31, 35
 - ventilator-associated events, 421
- pleural line, 18, 30
- pneumococcal meningitis, **430**
- pneumococcal pneumonia, antibiotics, **456**
- Pneumocystis jirovecii* pneumonia, 207, 450, 481, 483, 484, 487, 490
- pneumonia, 207, **405, 449–462, 578**
 - acute interstitial, 208
 - antibiotics, **432, 455–456**
 - HIV infection, **487**
 - ventilator-associated, 421, **422, 432, 456, 457**
 - chest X-ray, 398, 454, 461–462
 - Pneumocystis jirovecii*, 490
 - ventilator-associated, 421
 - cryptogenic organizing, 208, 453
 - diagnosis, 452–455
 - HIV infection, **487**
 - mechanical ventilation, 193
 - pathogens, 458–459
 - Pneumocystis jirovecii*, 207, 450, 481, 483, 484, 487, 490
 - risk factors, 451
 - sepsis, 404, 457
 - treatment, 455–459
 - ventilator-associated *see* ventilator-associated pneumonia
- pneumonia severity index, 454
- pneumothorax, 227, 578, *see also* barotrauma
 - acute coronary syndromes vs, 126
 - aortic dissection vs, 103
 - ARDS, 211
 - bedside tracheostomy, 43, 46
 - central venous access, 19
 - mechanical ventilation, 192
 - ultrasonography, 30, 35
- point-of-care glucose testing, 57
- poisoning, 299–308
 - acute hepatic failure, 360
- polydipsia, 524
- polyenes, **446**
- polymerase chain reaction (PCR), 398
- polymorphic ventricular tachycardia, 110, 111, 115
- polymyxin B, **439**
- polymyxins, 443
- polyneuropathy, 63
- portal drainage, pancreas transplantation, **373**

- portal reconstruction, liver transplantation, 367
- portal vein thrombosis, liver transplantation, 367
- posaconazole, **445**
- positive end-expiratory pressure (PEEP), 178, 185, 186
 - ARDS, 191, 210
 - bronchospastic conditions, 224
 - pregnancy, 378
 - ventilator-associated complications, 393
- positive pressure ventilation see non-invasive positive pressure ventilation
- possible ventilator-associated pneumonia, 450
- post-extubation respiratory failure, NPPV, 197–198
- post-infectious encephalitis, 467
- post-obstructive pneumonia, 452
- post-partum hemorrhage, 380–381
- post-renal AKI, 501
- post-thrombotic syndrome, 584
- post-transplant lymphoproliferative disorder, 375
- post-traumatic stress disorder, 10
- posterior enhancement, ultrasonography, 28
- posterior fossa hematoma, 271
- postoperative ARF, NPPV, 198
- postoperative fever, 393, 394
- postoperative hyperkalemia, 515
- potassium (level), tumor lysis syndrome, 589
- potassium
 - for hypernatremia, 529
 - removal, 515
- potassium acetate, **518**
- potassium chloride, 517, **518**
- potassium citrate, **518**
- potassium phosphate, **518**, 522
- potassium-sparing diuretics, 518
- pounding sensation, neck, 111
- power (parameter), LV assist devices, 95
- pralidoxime, **306**
- prasugrel, **131**, 136, 137
- pre-eclampsia, 384–385
- pre-renal acute kidney injury, 501
 - FeUrea, 503
- pre-shock, 152
- predicted body weight, 185
- prednisone
 - Pneumocystis jirovecii* pneumonia, **487**
 - pneumonia, **456**
- pregnancy, 377–386
 - acute abdomen, 324
 - acute coronary syndromes, 134
 - acute kidney injury, 505
 - aortic dissection, 106
 - ARDS, 213
 - bronchospastic conditions, 227–228
 - cardiac arrest, 165, 378–380
 - CNS infections, 475
 - coagulopathies, **572**
 - delirium, 247
 - neurotrauma, 272
 - physiology, **378**
 - pneumonia, 458
 - sepsis, 411
 - shock, 156
 - strokes, 260
 - trauma, 344
 - vascular effects, 344
 - venous thromboembolism, 383, 384, 584
- preload, 82
 - mechanical ventilation, 192
 - peri-partum cardiomyopathy, 383
- preoxygenation
 - airway management, 5–6, 66
 - NPPV for, 198
- pressure control, mechanical ventilation, 188
- pressure cycled ventilation, 187
- pressure limited ventilation, 188
- pressure microensors, intracranial pressure, 286
- pressure-regulated volume control, ventilation, 188
- pressure support ventilation, 188, 235
- pressure trigger, ventilation, 187
- pressure ulcer infection, 426–427
- pressure–volume loop, 178, 182
- probable ventilator-associated pneumonia, 450
- probes, ultrasonography, 27, 34
- procainamide, 115
- procalcitonin, 397, 444, 454, 456
- prone positioning, ARDS, 211
- propofol, 15
 - anti-shivering protocol, 164
 - bronchoscopy, 38
 - bronchospastic conditions, 224, **225**
 - children, 273, 282
 - pregnancy, 227
 - status epilepticus, **280**, 281
- propofol infusion syndrome, 15, 62
- PROPPR trial, 346
- propranolol, aortic dissection, **105**
- protamine sulfate, 557, **558**, 559
- protease inhibitors, 485
- protein
 - enteral nutrition, 50
 - acute hepatic failure, 363
 - loss, burns, 348
 - total parenteral nutrition, 51
 - energy requirements, **52**
 - renal failure, 53
- protein-bound antibiotics, 434
- prothrombin complex concentrates, 557, **558**, 559
- prothrombin time (PT), 561, 564, 566–567, 570
 - abnormal, 567–568
 - disseminated intravascular coagulation (DIC), 562
- protocols
 - anti-shivering, 164
 - massive transfusion, 547–548
 - thyroxine replacement, 297
 - weaning, 237

- proton pump inhibitors (PPIs), 64, 312, 313
 - acute hepatic failure, 362
- pseudomembranous colitis, 423
- Pseudomonas aeruginosa*, 421
 - pneumonia risk factors, 451
- pseudothrombocytopenia, 568
- PSI risk class, pneumonia, 454
- psychogenic status epilepticus, 278
- PTH-related protein, 530
- pulmonary artery catheters, monitoring via, 77–78, 81
- pulmonary artery pressure, 78
- pulmonary artery wedge pressure, 78
- pulmonary edema, 142, 452
 - ARDS vs, 207
 - NPPV, 197
 - ultrasonography, 31, 35
- pulmonary embolectomy, 581, **582**
 - removed emboli, 585
- pulmonary embolism, 219, 379, 453, 575–585
 - acute coronary syndromes vs, 126
 - aortic dissection vs, 103
 - cardiac ultrasonography, 181, 580
 - diagnosis, 578–581
 - peripartum, 156
 - pregnancy, 383–384
 - prognosis, 584
 - severity index (PESI), 579, 583
 - treatment, 581–584
 - ventilator-associated events vs, 420
- pulmonary hemorrhage, ventilator-associated events
 - vs, 421
- pulmonary hypertension, 579, 580
- pulmonary vascular resistance, 78, 192
- pulsatility, left ventricular assist devices, 95
- pulse contour analysis, 76, 80
- pulse oximetry, 171–172
 - abnormal hemoglobin, 177
 - arterial blood gases vs, 174
- pulse power analysis, 76–77
- pulseless electrical activity, 159, 160, 162
- pulses, acute hepatic failure, 361
- pupils
 - acute hepatic failure, 361
 - fixed dilated, 270
- pyelonephritis, complicated, **432**
- pyridoxine
 - ethylene glycol poisoning, **305**
 - isoniazid poisoning, **305**
- pyrogenic reactions, dialysis, **511**
- pyruvate, brain, 288
- QRS complex, arrhythmias, 108, 110, 111
- QT prolongation, 63, 115
- QTc intervals, 63
- quantitative waveform capnography, 172
- quick sequential organ failure assessment (qSOFA)
 - score, **403**
- rabbit ears, RSR complexes, 111
- rabies, 474
- raccoon eyes, 268
- radial artery, 21, 25, 75
 - arterial blood gases, 174
- radionuclide imaging see nuclear medicine
- ramipril, 382
- rapid dipstick test, urine, 397
- rapid sequence induction, 7
- rapid shallow breathing index (RSBI), 236
 - tidal volume on, 239
- Rapid Ultrasound in SHock (RUSH) exam, 33
- rasburicase, 591
- re-entry tachyarrhythmias, 109, 114
- rebleeding, gastrointestinal, 313, 314, 318
- rebound hypotension, **88**
- recombinant factor VIIa, 557, 559
- recovery phase, ARDS, 205
- red blood cells
 - blood smear, thrombocytopenia, 570
 - preparations, 545–546
 - fever, 550
 - sepsis from, 549
 - tagged RBC studies, GI bleeding, 315, 316
- refeeding syndrome, 54
- referral to specialists, heart failure, 144
- refractory respiratory failure, 226–227
- refractory status epilepticus, 278, 279, 281, 282
- regional citrate anticoagulation, 510
- rehabilitation, cardiac, **132**
- reintubation, 238
- REMATCH study, HeartMate™ devices, 94
- remifentanyl, NPPV, 200
- renal failure, see also acute kidney injury
 - abdominal compartment syndrome, 327
 - acute hepatic failure, 363
 - antibiotics, 434
 - diuretics, 144
 - energy requirements, **51**
 - hydroxyethyl starch, 498
 - neutropenic fever, 589
 - phosphate replacement, 523
 - total parenteral nutrition, 53
- renal perfusion pressure, 504
- renal replacement therapy, 507–512, see also dialysis
 - acute hepatic failure, 363
 - antibiotics, 434
 - complications, 510, **511**
 - lithium poisoning, **305**
 - monitoring, 511
 - procedure, 508
 - prognosis, 510
 - reasons to discontinue, 512
 - timing, 507–508
- renal views, ultrasonography, 31
- renin, 494
- renin–angiotensin–aldosterone axis, 494

- respiratory acidosis, 538, 539, 541
- respiratory alkalosis, 538, 539, 541, **542**
- respiratory cultures, 454, 455
- respiratory failure, 169–170, 232, *see also* acute respiratory distress syndrome
 - abdominal compartment syndrome, 327
 - acute hepatic failure, 360
 - gastrointestinal bleeding, 312
 - HIV infection, 482, 484
 - NPPV for, 197–198
 - refractory, 226–227
 - signs, 220
 - total parenteral nutrition, 52–53
- respiratory frequency, 185
- respiratory monitoring, 169–183
- respiratory muscles, monitoring strength, 179–181
- respiratory quotient, 53
- respiratory rate, 186
 - bronchospastic conditions, 224
- respiratory specimens, bacteriology, 397
- resting energy expenditure, burns, 348
- resuscitation
 - burns, 350
 - trauma, 343, 344
- resuscitative transesophageal echocardiography, 162
- reticular formation (ascending reticular activating system), 294
- reverberation artifact, ultrasonography, 28
- Revised Trauma Score, 340
- rewarming phase, 165
- rhabdomyolysis
 - hyperphosphatemia, 519
 - urine output for, 502
- ribavirin, 458
- Richmond Agitation-Sedation Scale, 11
- RIFLE criteria, acute kidney injury, 500, **501**
- right atrial pressure, 78, 79
- right heart catheterization, heart failure, 143
- right ventricle
 - Impella axial flow pump, 91, 99
 - LVAD troubleshooting, 95
 - pressure, 78
 - pulmonary hypertension, 579, 580
 - ultrasonography, 29, 35
- Riker Agitation-Sedation Scale, 11
- ring-down artifact, ultrasonography, 28
- Ringer's lactate, 497
- risperidone, 247
- rivaroxaban, **582**
- road traffic accidents, transfer to level 1 trauma facilities, 342
- Rockall score, 314, 318
- rocuronium, laryngoscopy, 7
- Roux-en-Y venting jejunostomy, pancreas transplantation, **373**
- RSBI (rapid shallow breathing index), 236
 - tidal volume on, 239
- RSR complexes, 111
- rule of nines, burns, 357
- Rumack–Matthew nomogram, 300
- RUSH (Rapid Ultrasound in SHock) exam, 33
- salicylates, toxicity, 302
- salvage treatment, shock, 154
- sampling, bronchoscopy, 39
- sarcoidosis, 467
- scalpels
 - in central venous access, 19
 - tracheostomy, 46
- scarring, burns, 355–356
- schistocytes, 570
- scintigraphy *see* nuclear medicine
- screening
 - abdominal aortic aneurysm, 103
 - acute coronary syndromes, 125, 162
 - ARDS, 206
 - blood pressure, 103
 - Clostridium difficile* infection, 423
 - CNS infections, 466
 - delirium, 245–246
 - heart failure, 140
 - HIV infection, 482
 - neurotrauma, 267
 - shock, 150
 - strokes, 253
 - venous thromboembolism, 577
- seashore sign, 30, 35, 181
- Seattle Heart Failure Model, 142
- secondary carina, 39
- secondary survey, trauma, 340
- sedation, 10–16, 62–63
 - ARDS, 210
 - bedside tracheostomy, 44, 45
 - bronchospastic conditions, 224
 - daily interruption, 63, 233
 - NPPV, 200
 - pregnancy, 227
- sedation vacations, 10, 63
- sedatives, toxic syndromes, **301**
- seizures, *see also* status epilepticus
 - cholinergic toxidrome, **306**
 - continuous EEG monitoring, 289
 - features, 254
 - hyponatremia, 526
 - strokes vs, 257
 - traumatic brain injury, 270
 - tumor lysis syndrome, **590**
- Seldinger technique, tracheostomy, 46
- sepsis, 389, 402–413
 - abdomen, 321–325
 - ARDS, 191, 206, 208–209
 - asplenia, 400
 - from blood transfusions, 549
 - definition, 402, **403**

- sepsis (*cont'd*)
 - diagnosis, 404–409
 - dobutamine, **86**
 - hemorrhage, treatment, **571**
 - hypophosphatemia, 521
 - lactate, 397, 409
 - line-related, 155
 - pneumonia, 404, 457
 - thrombocytopenia, 571, 573
 - treatment, 409–411
- Sepsis in Obstetrics Score, 411
- septal wall motion, ultrasonography, 29
- septic intracranial thrombophlebitis, 473
- septic shock, 323, 404, 406, 408
 - definition, **403**
 - incidence, 148
 - pneumonia, 457
 - pregnancy, 156
 - refractory, 324, 411
- sequential organ failure assessment (SOFA) score, 406, **407**
 - quick (qSOFA), **403**, 406
- serotonin syndrome, **301**, **306**, 395
- serum anion gap, 302
- sevelamer, **520**
- severe sepsis, 404, 406
 - definition, **403**
- sevoflurane, 226
- shadowing, ultrasonography, 28
- shivering, 164–165
- shock, 83–88, 148–158, *see also* hypovolemic shock;
 - septic shock
 - acute hepatic failure, 360
 - burns, 348
 - prognosis, 156–157
 - pulmonary embolism, 384, 579, 584
 - treatment, 84–88, 154–156
 - ultrasonography, 33
- SHOCK trial, 146
- short gut syndrome, 334
- shunting, ARDS, 205
- sidestream technique, end-tidal CO₂, 172
- signs of life, trauma, 340
- silicone collagen blend (Biobrane®), **353**
- silver nitrate, **353**, 354
- silver sulfadiazine, **353**, 354
- simultaneous deceased donor pancreas and living donor
 - kidney transplantation, **372**
- simultaneous pancreas–kidney (SPK) transplantation, 371, **372**, 376
- sinus node dysfunction, 109
- sinusitis, nasotracheal intubation, 393
- six hour bundle, Surviving Sepsis Campaign
 - guidelines, 410
- skin
 - breakdown, NPPV, 201
 - hypovolemia, 495–496
 - necrotizing infections, 394
 - sepsis, **405**
 - shock, 151
- skull, fractures, 268, 271
- small bowel transplantation, 374–375
- small cell lung cancer, SVC syndrome, 587
- smoking
 - COPD, 219
 - risk of acute coronary syndromes, 124, 125
- SOAP II trial, 148
- sodium, 495, 498
 - fractional excretion (FeNa), 503
 - levels, 523
 - organ donors, 297
 - overcorrection, 526, 529
 - reabsorption, 494
 - in therapeutic fluids, **497**
 - urine, 525
- sodium bicarbonate, **306**, **542**
 - for hyperkalemia, 515
- sodium nitrite, **304**
- sodium phosphate, 522
- sodium polystyrene sulfate, 65, 515
- sodium thiosulfate, **304**
- SOFA score, **403**, 406, **407**
- soft tissues
 - necrotizing infections, 394
 - sepsis, **405**
- sotalol, 115
- speed (parameter), LV assist devices, 95
- spinal abscess, 465, 473, 479
- spinal cord compression, hyperkalemia vs, 514
- spinal cord injury, 265, 266, 267
 - imaging, 269
 - surgery, 271
 - treatment, 271, 272
- spironolactone, 383
- splenorenal recess with hemothorax view, 31
- spondylitis, 467
- spontaneous breathing trials, 233, 234–237
- sputum
 - HIV infection, 483
 - pneumonia, 453
- ST elevation myocardial infarction, 123, 124, 134
 - electrocardiography, 129
- stabilization
 - trauma, 343
 - treatment for shock, 154
- staphylococcal scalded skin syndrome, 349
- static compliance, 186
- statins, 125
 - high intensity, **132**
- status asthmaticus, 191
 - mortality, 228
 - permissive hypercapnia, 224
- status epilepticus, 275–283
 - continuous EEG monitoring, 289
 - titration of therapy, 289
- steal syndrome, dialysis, **511**

- steatosis (hepatic), total parenteral nutrition, 54
- stents
 - acute coronary syndromes, 134
 - bronchial, **457**
 - strokes, 260
- sterility, central venous access, 18
- Stevens–Johnson syndrome, 349
- stool bacteriology, 397
 - Clostridium difficile*, 425
- stratosphere sign (barcode sign), 30, 35
- Streptococcus pneumoniae*
 - antibiotics, **422**
 - CNS infections, 474
 - pneumonia, 450, 451
 - pregnancy, 475
- stress
 - energy requirements, **51**
 - on gastrointestinal tract, 64
 - hyperglycemia, 55, 56, 57
 - risk of acute coronary syndromes, 124
- stress ulcers, 64
 - ARDS, 211
 - bleeding, 193, 312
- stridor, management, 238
- stroke volume, 76, 78, 82
- strokes, 250–264
 - complicating acute coronary syndromes, 133
 - left ventricular assist devices, 97
 - prognosis, 261
 - treatment, 258–260
- sub-massive pulmonary embolism, 575
- subarachnoid hemorrhage, 250, 251, 252, 263, 267, 467
 - delayed cerebral ischemia monitoring, 289
 - presentation, 254
 - severity scores, 256
 - tracheostomy, 43
 - treatment, 258, 260
- subclavian artery, 18
- subclavian lateral vein, 17–18
- subclavian medial vein, 17
- subcostal four chamber position, 79
- subcutaneous insulin, 58, 60
- subdural hematoma, 271, 467
- subfalcine herniation, 286
- subglottic secretion drainage ports, endotracheal tubes, 452
- substance abuse, fever, 395
- subtotal colectomy, 318
- subxiphoid position, cardiac ultrasonography, 28
- succimer, **305**
- succinylcholine
 - burns, 352
 - laryngoscopy, 7
- sucralfate, **317**
- suicide ventricle, phenylephrine for, **87**
- sulbactam, **435**
- sulfamethoxazole/trimethoprim, **438**
- sulfonamides, **438**
- sulfonyleureas, poisoning, **306**
- super-refractory status epilepticus, 278, 279
- superior vena cava syndrome, 586, 587, 588, 589
 - grading, **591**
 - imaging, 591
 - treatment, 592, 593
- supplements
 - enteral nutrition, 50
 - vitamin K, 565
- supraventricular tachycardia, 110
- surgery
 - acute mesenteric ischemia, **334**
 - burns, **353**
 - craniotomy, intracerebral hemorrhage, 258
 - energy requirements, **51**
 - gastrointestinal bleeding, 316, **317**
 - meningitis after, 463
 - NPPV, 198
 - pneumonia, **457**
 - site infections, 394
 - culture, 397
 - for traumatic brain injury, 270, 271
- Surviving Sepsis Campaign guidelines (2018), 410
- sympathomimetic toxidrome, **301**
- symptomatic leukostasis, 586, 587, 588, 589, 591, 592
 - probability from symptoms, **590**
- synchronized intermittent mandatory ventilation, 188
- syndrome of inappropriate ADH secretion (SIADH), 523, 524
- SYNERGY trial, 137
- SYNTAX trial, 137
- syphilis, meningovascular, 474
- systemic inflammatory response syndrome, 149, 348
- systemic vascular resistance (SVR), 78, 82
 - pregnancy, **378**
- systolic blood pressure, shock, 154
- systolic heart failure, 140, 144
- T-piece trials, 234, 235
- T-pieces, 240
- tabes dorsalis, 474
- tachyarrhythmias, 108, 109, *see also* broad complex
 - tachyarrhythmias; narrow complex
 - tachyarrhythmias
 - diagnostic algorithms, 112, 113
 - treatment, 114
- tachycardia, *see also specific types*
 - hypovolemia, 496
 - sepsis, 409
- tachyphylaxis, 85
- tacrolimus, 368
- Taenia solium*, brain, 475
- tagged RBC studies, GI bleeding, 315, 316
- tamponade (cardiac), 28–29, 151
 - aortic dissection vs, 103
 - with left ventricular assist devices, 96
- TandemHeart, 93, 100
- tazobactam, **435**
- technetium-99m sulfur colloid, GI bleeding, 315

- tedizolid, **441**
- telavancin, **440**
- temperature modulation *see* therapeutic temperature modulation
- tenecteplase, **132**
- tension pneumothorax, 46, 227
- terminal illness *see* palliative care
- tertiary survey, trauma, 340
- tetracyclines, **438**
- thawed plasma, 546
- therapeutic temperature modulation
 - acute hepatic failure, 362
 - cardiac arrest, 159, 163–165
 - intracranial pressure management, 287
 - traumatic brain injury and, 270
- thermal diffusion flowmetry, cerebral blood
 - flow, 288
- thermistors, 391
- thiamine, 279, **305**, 307
- thienopyridine, coagulation parameters, 367
- third space, 495
- thoracic aorta, transmural pressure, 192
- thoracic surgery, NPPV, 198
- thoraco-abdominal paradox, 170
- thoracotomy, trauma, 340
- Thoratec HeartMate™ devices, 94, 101
- three windows, 151
- thrombectomy, 264
- thrombocytopenia, 561–574, *see also* heparin-induced thrombocytopenia
 - investigations, 568–570
 - pre-eclampsia, 385
 - severity classification, 566
 - shock, 152
- thrombolysis
 - acute mesenteric ischemia, 333, 334
 - pregnancy, 584
 - pulmonary embolism, 581, **582**
 - strokes, 258
- thrombolytics, **132**, 133
- thrombophlebitis, 578
- thrombosis, 576, *see also* venous thromboembolism
 - acute mesenteric arterial, 330
 - atherosclerosis, 124
 - cavernous sinus, 473
 - cerebral venous sinuses, 252
 - internal jugular vein (IJ), 22
 - left ventricular assist devices, 95, 96, 97
 - leg veins, 32, 36
 - mesenteric venous (MVT), 331, 332, 333
 - portal vein, liver transplantation, 367
 - recombinant factor VIIa, 559
- thrombotic thrombocytopenic purpura, 563, 566, **572**
- thymoglobulin, 370, 375
- thyrotoxic periodic paralysis, exclusion, 517
- thyroxine replacement protocol, 297
- ticagrelor, **131**, 136
- ticks, diseases from, 466
- tidal volume, 185, 186
- ARDS, 203, 205, 206, 209, 210
 - bronchospastic conditions, 224
 - on rapid shallow breathing index, 239
- tigecycline, **438**
- time cycled ventilation, 187
- time-dependent activity, antibiotics, 433, 434
- time-gain compensation, ultrasonography, 27
- time trigger, mechanical ventilation, 187
- TIMI risk score, 127
- tinzaparin, 384
- tirofiban, **131**
- tissue perfusion, 150, 151
- tissue plasminogen activator (tPA)
 - pulmonary embolism, 384, **582**
 - stroke, 256, 260, 261
- tissue transfer (local), burns, **353**
- tobramycin, **440**, **457**
- tongue, airway difficulty, 4, 9
- torsades de pointes, 110, 115
- total artificial hearts, 89, 97
- total parenteral nutrition, 50–54
 - small bowel transplantation replacing, 374
- toxic epidermal necrolysis, 349
- toxic syndromes, 299–308
- toxins, *Clostridium difficile*, 422, 425
- Toxoplasma gondii*, encephalitis, 475
- trachea, 3
- tracheal reflex, 295
- tracheobronchomalacia, 219
- tracheostomy
 - bedside percutaneous dilational, 42–48
 - trauma, 343
 - traumatic brain injury, 43, 271, 346
 - weaning from ventilation, 238
- traffic accidents, transfer to level 1 trauma
 - facilities, 342
- tranexamic acid
 - CRASH-2 trial, 346
 - post-partum hemorrhage, 381
- transcatheter embolization, GI bleeding, **317**
- transcranial Doppler, brain death, 296
- transesophageal echocardiography
 - pulmonary embolism, 580
 - resuscitative, 162
- transfusion-associated circulatory overload, 550
- transfusion-associated dyspnea, 551
- transfusion-related acute lung injury (TRALI), 395, 549–550
- transient ischemic attack, 250, 258
- transjugular intrahepatic portosystemic shunt, liver
 - transplantation, 367
- transjugular liver puncture, minimum platelet
 - count, **571**
- transmural pressure, thoracic aorta, 192
- transplantation
 - hematopoietic stem cells, ARDS, 206
 - kidneys, 368–370, 371, **372**, 376
 - liver, 362, 363, 365–368
 - with intestine, **374**

- techniques, 367–368, 375
- lungs, ARDS, 206
- multivisceral, 374–375, 376
- pancreas, 370–374
- small bowel, 374–375
- transpulmonary thermomodulation cardiac output, 76
- transthoracic echocardiography *see* cardiac ultrasonography
- transverse myelitis, 267
- trauma, 337–346, *see also* burns
 - AMPLE (history), 339–340, 350
 - ARDS, 206
 - coagulopathy prevention, 565, 571
 - concomitant burns and, 352
 - from endotracheal tubes, 8
 - FAST exam, 26
 - laboratory diagnosis, 341
 - scoring systems, 340
- trauma centers, 339
- transfer to, 342–343
- traumatic brain injury, 265–274
 - incidence, 337
 - tracheostomy, 43, 271, 346
- tricyclic antidepressants, poisoning, **306**
- trigger phase, mechanical ventilation, 187
- trigger sensitivity, mechanical ventilation, 186
- trimethoprim, **438**
- trimethoprim-sulfamethoxazole
 - hyperkalemia, 514
 - Pneumocystis jirovecii* pneumonia, **487**
- TRITON-TIMI 38 trial, 136
- troponins, cardiac, 129, 580
- Trousseau sign, 533
- trypanosomiasis, 475
- tuberculosis, brain, 474
- tubular reabsorption of phosphate (TRP), 520
- tumor lysis syndrome, 396, 586, 588, 589
 - Cairo-Bishop grading, **590**
 - treatment, 591, 592
- ulcers *see* stress ulcers
- ultrasonography, *see also* bedside ultrasonography; cardiac ultrasonography
 - acute hepatic failure, 361
 - acute mesenteric ischemia, 333, 335
 - for airway management, 66
 - arterial access, 20
 - cardiac arrest, pregnancy, 379
 - for central venous access, 17–18, 23, 24, 65
 - diaphragmatic dysfunction, 180
 - equipment, 27
 - fever, 398
 - hypovolemia, 496
 - inferior vena cava, 29–30, 35, 79, 496
 - large curvilinear probes, **27, 34**
 - lungs, 30–31, 181, 461, 462
 - physics, **26, 27**
 - tracheostomy, 46
 - trauma, 341
 - venous, 580
 - uncal herniation, 286
 - unfractionated heparin, 553
 - acute coronary syndromes, **131**
 - coagulation parameters, 567
 - pulmonary embolism, 581, **582**
 - reversal, 557
 - thromboprophylaxis, 577
 - upper airway *see* airway (upper)
 - uremia, 503
 - coagulation parameters, 567
 - thrombocytopenia, 570
 - ureters, kidney transplantation, 369
 - uric acid, tumor lysis syndrome, 589
 - urinary catheter-associated bacteriuria, 393
 - urinary tract
 - infections, 67, **405**
 - bacteriology, 397
 - catheter-associated (CAUTIs), 67, 393, 397, 417–419
 - obstruction, 501, 505
 - ultrasonography, 32
 - urine
 - acute kidney injury, 503
 - anion gap, 541
 - culture, 417, 418–419
 - hypokalemia, 517
 - osmolality, 525, 528
 - rapid dipstick test, 397
 - sodium, 525
 - urine output
 - acute kidney injury, 500
 - electrical burns, 354
 - for rhabdomyolysis, 502
 - shock, 151
 - uteroplacental blood flow, **378**
 - uterus, atony, 381
 - V/Q scan, 580
 - pregnancy, 584
 - vaccinations
 - meningitis, 466
 - pneumonia prevention, 452
 - vacuum-assisted devices, abdominal closure, 343
 - vagus nerve, 3–4
 - valacyclovir, **447**
 - valganciclovir, **447**
 - valproate, 279, **280, 281**
 - meropenem and, **437**
 - poisoning, **306**
 - value-based decisions, palliative care, 69
 - valvular heart disease, 207
 - vancomycin, 416, 423, 425, **440**
 - dexamethasone and, 472
 - methicillin-resistant *S. aureus*, **457**
 - retention enema, **431**
 - stopping, 444
 - varices, gastroesophageal, 312, 313, 316
 - varsartan, 382

- vascular malformations
 - Dieulafoy, 312
 - strokes, 251
- vasculopathies, strokes, 251
- vasoactive drugs, 82–88, 144, **155**
- vasodilators
 - aortic dissection, **105**
 - heart failure, 143
- vasogenic edema, cerebral, 285
- vasopressin (medicine), **84, 88, 155**
 - cardiac arrest and, 161
- vasopressin (antidiuretic hormone), 494, 523–524,
 - see also* arginine vasopressin
- vasopressors, 83, 85
 - acute hepatic failure, 362
 - acute kidney injury, 504
 - organ donors, 297
 - pulmonary embolism, 581
- vasospasm, subarachnoid hemorrhage, 252
- vegetative state, 293
- veno-venous bypass, liver transplantation, 367
- veno-venous hemofiltration *see* continuous veno-venous hemofiltration
- venous blood gases, 77, 175–176
- venous drainage, pancreas transplantation, **373**
- venous return
 - mechanical ventilation, 192
 - pregnancy, **378**
- venous sinus thrombosis, cerebral, 252
- venous thromboembolism, 64, 575–585
 - ARDS, 211
 - diagnosis, 578–581
 - mechanical ventilation, 193
 - neurotrauma, 270
 - pregnancy, 383, 384, 584
 - prognosis, 584
 - prophylaxis, 577–578
 - recurrent, 578
 - stroke patients, 259
 - treatment, 581–584
- venous ultrasonography, 580
- ventilation, *see also* hyperventilation; non-invasive positive pressure ventilation
 - bilevel positive pressure, 188, 196
 - complications, 66
 - manual, 4, 6, 7, 160, 190
 - mechanical, 184–193
 - ARDS, 210
 - bronchospastic conditions, 223–224
 - capnography waveform, **175**
 - complications, 231
 - gastrointestinal bleeding, 312
 - loops, 178, 182
 - pregnancy, 228, 378
 - troubleshooting, 190
 - waveforms, 177, 182–183
 - weaning, 231–240
 - withdrawal, 70
 - protective, HIV infection, 484
- ventilator-associated condition, 419, 450
- ventilator-associated events, 193, 393, 419–422, 449
- ventilator-associated pneumonia, 67, 155, 393, 419–422, 449, 450
 - antibiotics, 421, **422, 432, 456, 457**
 - inhaled antibiotics, 442
 - prevention, 452
- ventilator ‘bundles,’ 193
- ventilators
 - problems, 239
 - setting, 186–190
- ventricular fibrillation, 159
 - recurrent, 115
 - transesophageal echocardiography, 162
- ventricular tachycardia, 110, 111, 115, 117, 159
- ventriculitis, infectious, 286
- verapamil, 113, 114–115
- verbal response, Glasgow Coma Scale, 294
- vertebral osteomyelitis, 479
- video laryngoscopes, 5
- Virchow's triad, 331
- viruses, *see also specific types*
 - encephalitis, 465, 471, 474
 - meningitis, 464, 471, 474
 - pneumonia, 450
 - thrombocytopenia, 570
- visual analog scale, pain, 12
- vital capacity, 179
 - diaphragmatic dysfunction, 180
- vitamin K, 553, 556, **557, 558, 562, 571**
 - deficiency, coagulation parameters, 567
 - dietary, 565
- vitamin K antagonists, 553
 - major hemorrhage rates, **553**
 - monitoring, 554
 - reversal, 556, **558**
 - time to hemostasis after stopping, 559
- vocal cords
 - abnormal motion, 219
 - bronchoscopy, 39
 - paralysis, 8
- volume cycled ventilation, 187–188
 - bronchospastic conditions, 224
- volume overload, 497
 - left ventricular assist devices, 96
 - total parenteral nutrition, 53
- volume reduced platelet products, 547
- volume-reduced red blood cells, 545
- volume support, mechanical ventilation, 188
- vomiting, endotracheal intubation, 66
- von Willebrand disease, coagulation parameters, 567
- von Willebrand factor cleaving protease, 563
- voriconazole, **445**

- warfarin, pregnancy, 584
- washed packed red blood cells, 545
- washed platelet products, 547
- water
 - body, 494–495
 - total parenteral nutrition, 52
- water balance, 495
- water deficit, 528
- waveforms
 - arterial access, 25
 - capnography, 172, **175**
 - intracranial pressure, 285
 - mechanical ventilation, 177, 182–183
- weaning
 - mechanical ventilation, 231–240
 - total parenteral nutrition, 52
- weight reduction, 125
- Wenckebach block, 109
- West Haven Hepatic Encephalopathy Scale, 361
- West Nile virus encephalitis, 469
- white cell count *see* leukocyte count
- whole bowel irrigation, 303
- wide complex tachyarrhythmias *see* broad complex tachyarrhythmias
- Wilson's disease, 360
- Winnipeg criteria, COPD, 226
- Winter's formula, 302, **539**
- work of breathing, 184
 - agitation *vs*, 239
 - bronchospastic conditions, 218
 - T-piece trials, 235
- wound infections, 394
 - culture, 397
- zolendronic acid, **531**

